



# Life Quality of Pediatric Patients with Central Nervous System Infections: A 1-Year Follow-up

Hamid Rahimi Hajiabadi <sup>1</sup>, Zahra Pourmoghaddas <sup>1</sup>, Omid Yaghini <sup>2</sup>, Tayebbeh Mohammadi <sup>3</sup>, Sharareh Moghim <sup>4</sup>, Maryam Khalili <sup>4</sup>, Hamidreza Roohafza <sup>5</sup>, Rana Saleh <sup>6,\*</sup>

<sup>1</sup> Infection Division, Department of Pediatric, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>2</sup> Neurology Division, Department of Pediatric, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>3</sup> Neurology Division, Department of Pediatric, School of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

<sup>4</sup> Department of Microbiology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>5</sup> Cardiac Rehabilitation Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>6</sup> Department of Pediatric, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

\*Corresponding Author: Department of Pediatric, Imam Hossein Children's Hospital, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran. Email: ranasaleh.medical.university@gmail.com

Received: 2 July, 2023; Revised: 27 September, 2024; Accepted: 5 October, 2024

## Abstract

**Background:** Central nervous system (CNS) infections can lead to long-term motor and cognitive complications in children.

**Objectives:** We aimed to investigate the prevalence of attention deficit-hyperactivity disorder (ADHD) and assess the quality of life in children with CNS infections.

**Methods:** A longitudinal study was conducted on 94 children (aged 6 - 15 years) with CNS infection symptoms who were admitted to referral hospitals in Isfahan, Iran. Parents completed questionnaires assessing quality of life and neurological deficits at 6 and 12 months after discharge. All patients were evaluated by the corresponding physician for a final assessment.

**Results:** Attention deficit-hyperactivity disorder symptoms were diagnosed in 30 patients, with viral encephalitis being the most common infection. The quality of life in children with CNS infections was significantly lower compared to that of healthy children.

**Conclusions:** Central nervous system infections negatively impact the quality of life in children. Screening for ADHD and managing affected children can help improve outcomes. Further research and interventions are needed to mitigate long-term complications.

**Keywords:** Central Nervous System, Encephalitis, Pediatrics, Quality of Life, Attention Deficit-hyperactivity Disorder

## 1. Background

Central nervous system (CNS) infections refer to infections and inflammation of the brain, with a higher prevalence observed in pediatric populations (1-3). Children can experience a wide range of outcomes, from rapid recovery to severe dysfunction, depending on various risk factors (4, 5). Previous studies have shown a higher prevalence of attention deficit-hyperactivity disorder (ADHD) in children with CNS infections, including those who have fully recovered (6, 7). Risk factors for long-term neurological complications include focal neurological defects, abnormal

neuroimaging, infectious causes, and prolonged hospitalization (8). Children with CNS infections often require regular neuropsychological assessments, and rehabilitation services may be necessary for those with neuropsychological complications (3). However, there is limited understanding of the factors influencing outcomes in pediatric patients with CNS infections, making it difficult to accurately predict patient outcomes (9, 10).

In addition to traditional medical outcome measures such as mortality, the concept of quality of life has gained significant importance, especially in pediatric populations (11, 12). Quality of life encompasses physical,

emotional, and social aspects and serves as a key indicator of outcomes in children (13). The Pediatric Quality of Life Inventory Version 4.0 Generic Core Scales (PedsQL) is a standardized questionnaire used to assess quality of life in children, evaluating domains such as physical, emotional, social, and school functioning. Low PedsQL scores can indicate challenges such as depression, fatigue, decreased school attendance, and social pressures, which are critical when assessing the long-term effects of CNS infections (11).

## 2. Objectives

In this study, we examined the prevalence of ADHD and quality of life in pediatric patients with CNS infections admitted to the largest referral hospital and the main children's hospital in Isfahan, Iran.

## 3. Methods

### 3.1. Study Population

In this prospective longitudinal study, we analyzed medical records and clinical evaluation reports of patients suspected of having CNS infections, admitted to referral hospitals affiliated with Isfahan University of Medical Sciences between 2019 and 2020. The study focused on infants presenting with fever and seizures, children with fever and suspicious CNS symptoms, positive meningeal signs, and acute loss of consciousness. The inclusion criteria involved patients aged 6 to 16 years without underlying conditions such as endocrinopathies, metabolic disorders, genetic disorders, or previous nervous system conditions. We excluded patients with incomplete clinical, laboratory, and imaging data, as well as those diagnosed with non-infectious causes like autoimmune encephalitis. Additionally, patients who declined to participate or withdrew from the study were also excluded.

The diagnosis of CNS infection was confirmed by pediatric infectious disease and pediatric neurology experts (H.R., Z.P., O.Y.) after a thorough review of medical records and data. Initially, 108 patients were suspected of having a CNS infection, but 14 were excluded due to underlying conditions. Therefore, a total of 94 patients were included in the final analysis. Demographic information, clinical features, and laboratory data were collected anonymously using a research team-designed form. All eligible patients participated in the study after being informed about its

objectives, and parental consent was obtained to ensure the confidentiality of their information. The study protocol received ethical approval from the Isfahan University of Medical Sciences' ethics committee (IR.MUI.MED.REC.1399.344).

### 3.2. Study Procedure

The study was conducted in four steps (Figure 1).

T1: Upon the patient's admission, an initial assessment was conducted to gather demographic information and obtain their medical history. Cerebrospinal fluid (CSF) analysis, laboratory data, and neuroimaging were obtained.


T2: During hospitalization, CSF samples were collected for nucleic acid analysis using polymerase chain reaction (PCR) with the SinaPure ONE kit (SinaClon Co., Iran) according to the manufacturer's protocol. Based on laboratory results and medical records, patients were categorized into three groups: Bacterial infection, viral infection, and negative infection. Patients with CNS symptoms but no evidence of infection were classified as negative.

T3: Six months after being discharged from the hospital, patients were contacted through phone calls, and data collection forms were filled out regarding their current condition and well-being.

T4: One year after discharge, patients underwent a comprehensive examination. The SNAP-IV checklist was used for ADHD diagnosis, and the PedsQL™ 4.0 Generic Core Scale (Pediatric Quality of Life Inventory™) was used to measure health-related quality of life (HRQOL) in patients. The maximum age on the checklist was adjusted to 15 years old to match the inclusion criteria (6 - 16 years old). Data from the study groups, including self-assessment checklists, PedsQL scores, and ADHD scores from the final clinic visit, were analyzed to compare ADHD symptom presentation among the groups. Parental participation was sought to complete the required items in the data collection forms.

### 3.3. Statistical Analysis

IBM SPSS Statistics software (version 25.0, IBM Corp.) was used for the data analysis. Qualitative variables were presented as percentages, and quantitative variables as mean  $\pm$  standard deviation (SD). The chi-square test was employed for categorical variables, while the Kruskal-Wallis or Mann-Whitney U test was used for continuous



**Figure 1.** Study procedure

variables. Univariate logistic regression analysis was performed to assess the association of positive ADHD, higher PedsQL scores, seizures, and abnormal neurological exams between the CNS infection-positive and CNS infection-negative groups. A significance level of  $P < 0.05$  was considered statistically significant.

## 4. Results

### 4.1. Clinical Features and Characteristics of Patients

The study included 94 children with a mean age of  $13.29 \pm 6.73$  years (ranging from 6 to 16 years old). Among the participants, 42.6% were female and 57.4% were male. Fever was the most common presenting complaint in their medical history. Of all cases, 91 patients (96.8%) were discharged without neurological complications, while 3 patients (3.2%) died during hospitalization. Patients were categorized into three groups: Bacterial infection (11 patients, 11.7%), viral infection (37 patients, 39.4%), and negative for CNS infection (46 patients, 48.9%). Sixteen patients (17%) required admission to the intensive care unit (ICU). Additional clinical features and patient characteristics are detailed in Table 1.

### 4.2. Blood and Cerebrospinal Fluid Analysis

Significant differences were observed between the CNS infection-positive and CNS infection-negative groups regarding white blood cell (WBC) count in blood analysis ( $P = 0.008$ ). In the CSF analysis, significant differences were found between the groups in terms of WBC count, polymorphonuclear neutrophils (PMNs), lymphocytes, protein, and glucose levels ( $P < 0.05$ ). Detailed results of the blood and CSF analyses are presented in Table 2.

### 4.3. Molecular Analysis

Molecular analysis revealed that Enterovirus (43.2%), Herpes simplex virus (27.0%), and gram-positive cocci (45.5%) were the most frequently detected microorganisms in patients with positive CNS infection (Table 3).

### 4.4. Imaging Studies

Neuroimaging was performed on 84 out of the 94 subjects, with abnormal results found in 21 cases. Of these, 36.4% were in the bacterial group, and 45.9% were

**Table 1.** The Clinical Features and Characteristic of Patients in Different Groups <sup>a</sup>

| Variables                         | Total (n = 94) | Positive CNS Infection |                | Negative CNS Infection (n = 46) | P-Value           |
|-----------------------------------|----------------|------------------------|----------------|---------------------------------|-------------------|
|                                   |                | Bacterial (n = 11)     | Viral (n = 37) |                                 |                   |
| (A) Patients' characteristics     |                |                        |                |                                 |                   |
| Gender                            |                |                        |                |                                 | 0.49              |
| Male                              | 54 (57.4)      | 5 (45.5)               | 20 (54.0)      | 29 (63.0)                       |                   |
| Feeding in the first-year         |                |                        |                |                                 | 0.87              |
| Breastfeeding                     | 72 (76.6)      | 9 (81.8)               | 29 (78.4)      | 34 (74.0)                       |                   |
| Formula                           | 5 (5.3)        | 1 (9.1)                | 2 (5.4)        | 2 (4.3)                         |                   |
| Both                              | 17 (18.1)      | 1 (9.1)                | 6 (16.2)       | 10 (21.7)                       |                   |
| Attending school                  | 17 (18.1)      | 1 (9.1)                | 9 (24.3)       | 7 (15.2)                        | 0.41              |
| Exposed to sick person            | 10 (10.6)      | 0                      | 9 (24.3)       | 1 (2.2)                         | 0.03 <sup>b</sup> |
| Secondhand smoke                  | 18 (19.1)      | 3 (27.3)               | 11 (29.7)      | 4 (8.7)                         | 0.38              |
| Maternal education                |                |                        |                |                                 | 0.34              |
| Illiterate                        | 9 (9.6)        | 0 (0)                  | 5 (13.5)       | 4 (8.7)                         |                   |
| School                            | 56 (59.6)      | 7 (63.6)               | 22 (59.5)      | 27 (58.7)                       |                   |
| University                        | 29 (30.8)      | 4 (36.4)               | 10 (27.0)      | 15 (32.6)                       |                   |
| (B) Signs of patients             |                |                        |                |                                 |                   |
| Tachy/bradycardia                 | 36 (38.3)      | 4 (36.4)               | 18 (48.6)      | 14 (30.4)                       | 0.23              |
| Tachy/bradypnea                   | 36 (38.3)      | 5 (45.5)               | 16 (43.24)     | 15 (32.6)                       | 0.51              |
| CRT > 2 Seconds                   | 2 (2.1)        | 0                      | 1 (2.7)        | 1 (2.2)                         | 0.86              |
| Hypoxia                           | 14 (14.9)      | 3 (27.3)               | 5 (13.5)       | 6 (13.0)                        | 0.54              |
| Neck rigidity, Kernig, Brudzinski | 25 (26.6)      | 6                      | 12             | 7                               | 0.13              |
| Weakness, fatigue                 | 8 (8.5)        | 1 (9.1)                | 6 (16.2)       | 1 (2.2)                         | 0.08              |
| Skin rash                         | 22 (23.4)      | 2 (18.2)               | 6 (16.2)       | 14 (30.4)                       | 0.28              |
| Respiratory tract infection signs | 12 (12.76)     | 2 (18.2)               | 5 (13.5)       | 5 (10.9)                        | 0.77              |

Abbreviations: CNS, central nervous system; CRT, cardiac resynchronization therapy.

<sup>a</sup> Values are expressed as No. (%).<sup>b</sup> Indicates significant difference. Use of the chi-square statistical test.

in the viral group, showing a statistically significant difference ( $P < 0.001$ ).

#### 4.5. Patient Monitoring

Significant differences were observed between the CNS infection-positive and CNS infection-negative groups regarding the PedsQL total score and abnormal neurological examination ( $P < 0.05$ ). Additionally, bacterial and viral infections showed significant differences in terms of both PedsQL total score and abnormal neurological examination ( $P < 0.05$ ). After one year of follow-up, 30 patients were diagnosed with ADHD, and 35 patients experienced seizures (Table 4). The PedsQL score was significantly associated with positive CNS infection ( $P = 0.014$ ; OR = 1.270; 95% CI = 1.050 - 1.534), and abnormal neurological examination was also significantly associated with positive CNS

infection ( $P = 0.012$ ; OR = 3.799; 95% CI = 1.345 - 10.729) (Table 4).

#### 5. Discussion

The present study contributes to the existing body of research on CNS infections in pediatrics by emphasizing their association with complications such as mental disabilities and difficulties in learning and school activities. Our findings show a significant decline in the quality of life among patients with CNS infections, aligning with previous studies that have also reported reduced quality of life in pediatric patients suffering from inflammatory brain diseases (11). For instance, Liu et al. identified a statistically significant correlation between seizures or cognitive dysfunction at presentation and diminished quality of life in children with inflammatory brain illnesses (14). Similarly, Rao et al. reported significantly lower PedsQL scores in children diagnosed with acute encephalitis (15). Giger et

**Table 2.** Blood and Cerebrospinal Fluid Analysis of Patients in Different Groups<sup>a</sup>

| Variables          | Total (n = 94)  | Positive CNS Infection |                 | Negative CNS Infection (n = 46) | P-Value              |
|--------------------|-----------------|------------------------|-----------------|---------------------------------|----------------------|
|                    |                 | Bacterial (n = 11)     | Viral (n = 37)  |                                 |                      |
| (A) Blood analysis |                 |                        |                 |                                 |                      |
| WBC                | 11391 ± 5635    | 16936 ± 6753           | 10480 ± 4793    | 10778 ± 5324                    | 0.008 <sup>b</sup>   |
| PMN                | 53.98           | 59.63                  | 51.44           | 54.63                           | 0.520                |
| Lymphocytes        | 37.43           | 30.81                  | 39.69           | 37.25                           | 0.442                |
| Platelets          | 324944 ± 148461 | 352818 ± 83537         | 326200 ± 143183 | 317323 ± 165167                 | 0.541                |
| ESR                | 31.83 ± 26.50   | 49.00 ± 36.48          | 30.62 ± 24.94   | 28.65 ± 23.90                   | 0.147                |
| (B) CSF analysis   |                 |                        |                 |                                 |                      |
| WBC                | 164 ± 688       | 1168 ± 1754            | 66 ± 133        | 2.5 ± 8.5                       | < 0.001 <sup>b</sup> |
| PMN                | 25              | 71                     | 25              | 15                              | < 0.001 <sup>b</sup> |
| Lymphocytes        | 20              | 28                     | 34              | 6.4                             | < 0.001 <sup>b</sup> |
| Protein            | 40.30 ± 53.14   | 89.00 ± 58.48          | 50.90 ± 65.09   | 19.21 ± 23.04                   | < 0.001 <sup>b</sup> |
| Glucose            | 54.61 ± 17.62   | 39.00 ± 21.21          | 54.33 ± 18.18   | 58.56 ± 14.21                   | 0.01 <sup>b</sup>    |

Abbreviations: WBC, white blood cells; PMNs, polymorphonuclear neutrophils; ESR, erythrocyte sedimentation rate; CRP, c-reactive protein; CSF, cerebrospinal fluid.

<sup>a</sup> Values are expressed as (%) or mean ± SD.

<sup>b</sup> Indicates significant difference. Use of the Mann-Whitney test.

**Table 3.** The Frequency of Detected Viruses and Bacteria by Molecular Analysis<sup>a</sup>

| Viral Types            | Frequency (n = 37) | Bacterial Types         | Frequency (n = 11) |
|------------------------|--------------------|-------------------------|--------------------|
| Enterovirus            | 16 (43.2)          | Gram-negative cocci     | 2 (18.1)           |
| Herpes simplex virus   | 10 (27.0)          | Gram-positive cocci     | 5 (45.5)           |
| Mumps                  | 2 (5.4)            | <i>Burkholderia</i>     | 1 (9.1)            |
| Varicella zoster virus | 3 (8.1)            | <i>Salmonella typhi</i> | 1 (9.1)            |
| Epstein-Barr virus     | 6 (16.2)           | Brucellosis             | 1 (9.1)            |
| Adenovirus             | 0                  | <i>Citrobacter</i>      | 1 (9.1)            |
| Influenza viruses      | 0                  | -                       | -                  |

<sup>a</sup> Values are expressed as No. (%).

al. also demonstrated that children with post-hemorrhagic hydrocephalus had lower PedsQL scores compared to their healthy counterparts (16).

Moreover, ADHD has been reported with increased frequency among patients with CNS infections. International studies have highlighted a significantly higher prevalence of ADHD (50%) and learning disabilities (20%) in children with CNS infections, such as encephalitis, compared to the general population, where rates of ADHD range from 5 - 10% (17, 18). In our study, 30 patients (31.9%) out of the 94 hospitalized children exhibited ADHD symptoms within the year following hospitalization. Although ADHD can be identified in patients with brain infections after six

years of follow-up, our study's relatively small sample size is due to its short-term, one-year follow-up (17).

An important factor in understanding the clinical outcomes of different types of CNS infections, such as encephalitis, is recognizing the specific infectious agents and prognostic factors present in patients (15). In a study by Salih et al., the outcomes of 18 patients with herpes simplex encephalitis were examined over 13 years. The study found that 39% of the patients experienced moderate to severe sequelae, while 61% had either no sequelae or only minor ones (19). In the present study, abnormal neurological examination findings were observed in 25.5% (24 cases) of the patients after one year, with 8.4% in the bacterial group, 40.5% in the viral group, and 15.2% in the negative CNS infection

**Table 4.** The Monitoring of Patients in Different Groups for One Year<sup>a</sup>

| Variables                                   | Positive CNS Infection |                | Negative CNS Infection (n = 46) | P-Value <sup>b</sup> | P-Value <sup>c</sup> |
|---|------------------------|----------------|---------------------------------|----------------------|----------------------|
|   | Bacterial (n = 11)     | Viral (n = 37) |                                 |                      |                      |
| (A) Comparison Analysis                     |                        |                |                                 |                      |                      |
| PedsQL total                                | 16.54 ± 21.54          | 22.48 ± 21.98  | 11.82 ± 15.94                   | 0.047 <sup>d</sup>   | 0.020 <sup>d</sup>   |
| ADHD positive                               | 5 (45.5)               | 12 (32.4)      | 13 (28)                         | 0.553                | 0.251                |
| Abnormal neurological examination           | 2 (18.2)               | 15 (40.5)      | 7 (15.2)                        | 0.028 <sup>d</sup>   | 0.033 <sup>d</sup>   |
| Seizure                                     | 4 (36.4)               | 18 (48.6)      | 13 (28.3)                       | 0.161                | 0.077                |
| Abnormal imaging                            | 4 (36.4)               | 17 (45.9)      | 0 (0)                           | < 0.001 <sup>d</sup> | < 0.001 <sup>d</sup> |
| (B) Univariate Logistic Regression Analysis |                        |                |                                 |                      |                      |
| Variables and Group                         | OR                     |                | CI                              | P-Value              |                      |
| ADHD positive                               |                        |                |                                 |                      |                      |
| Negative CNS (ref)                          | 1                      |                | -                               | -                    |                      |
| Viral                                       | 1.066                  |                | 0.865 - 1.313                   | 0.547                |                      |
| Bacterial                                   | 1.180                  |                | 0.867 - 1.607                   | 0.293                |                      |
| Positive CNS                                | 1.093                  |                | 0.900 - 1.327                   | 0.369                |                      |
| PedsQL (upper 25th percentile)              |                        |                |                                 |                      |                      |
| Negative CNS (ref)                          | 1                      |                | -                               | -                    |                      |
| Viral                                       | 1.270                  |                | 1.050 - 1.534                   | 0.014 <sup>d</sup>   |                      |
| Bacterial                                   | 1.270                  |                | 0.928 - 1.737                   | 0.136                |                      |
| Positive CNS                                | 3.800                  |                | 1.300 - 11.110                  | 0.015 <sup>d</sup>   |                      |
| Seizure                                     |                        |                |                                 |                      |                      |
| Negative CNS (ref)                          | 1                      |                | -                               | -                    |                      |
| Viral                                       | 2.405                  |                | 0.968 - 5.972                   | 0.059                |                      |
| Bacterial                                   | 1.451                  |                | 0.363 - 5.801                   | 0.599                |                      |
| Positive CNS                                | 1.148                  |                | 0.912 - 5.060                   | 0.080                |                      |
| Abnormal neurological examination           |                        |                |                                 |                      |                      |
| Negative CNS (ref)                          | 1                      |                | -                               | -                    |                      |
| Viral                                       | 3.799                  |                | 1.345 - 10.729                  | 0.012 <sup>d</sup>   |                      |
| Bacterial                                   | 1.238                  |                | 0.219 - 6.988                   | 0.809                |                      |
| Positive CNS                                | 3.055                  |                | 1.126 - 8.293                   | 0.028                |                      |

Abbreviations: ADHD, attention deficit-hyperactivity disorder; CNS, central nervous system.

<sup>a</sup> Values are expressed as No. (%), mean ± SD unless otherwise indicated.<sup>b</sup> Comparison of groups viral, bacterial, and negative CNS.<sup>c</sup> Comparison of groups positive and negative CNS.<sup>d</sup> Indicate significant difference. Use of statistical test.

group. Patients with positive CNS infections had a higher frequency of abnormal neurological examination findings compared to those with negative CNS infections.

Rao et al. conducted a study on 142 patients over 10 years and found that the presence of seizures at admission was associated with ongoing seizure disorders during follow-up. Additionally, abnormal MRI findings and the number of abnormalities at initial presentation were linked to lower quality-of-life scores (15, 20). In our study, 35 out of 94 patients experienced seizures, with rates of 36.4% in the bacterial group, 48.6%

in the viral group, and 28.3% in the negative CNS infection group.

Neuroimaging was performed on 84 subjects, revealing abnormal findings in 21 cases. Among these, 80.1% were in the bacterial group, while 19% were in the viral group. The difference between patients with positive and negative CNS infections was statistically significant. Early diagnosis and treatment initiation are crucial, as delayed treatment increases the risk of severe complications. Empirical treatment with acyclovir is recommended if viral encephalitis cannot be ruled out within six hours of admission (20). Focal cortical



parenchymal abnormalities observed on MRI have also been shown to predict poorer long-term neurological outcomes (21).

While our study suggests that CNS infections lead to a decrease in the quality of life among pediatric patients, it is important to acknowledge its limitations. The small sample size and short follow-up period are critical.

### 5.1. Conclusions

Our study revealed that children hospitalized due to CNS infections require early diagnosis and timely, comprehensive treatment. Additionally, a higher incidence of mental function impairments was observed in children one year after the onset of the disease. These patients exhibited lower PedsQL scores, indicating a reduced quality of life. It is essential that all patients with encephalitis undergo long-term follow-up evaluations, and neuropsychological rehabilitation should be provided for these individuals. Further research in this area is necessary to enhance knowledge and potentially improve patient care and outcomes.

### Footnotes

**Authors' Contribution:** Study concept and design: H. R., Z. P., and O. Y.; conducting laboratory work: S. M. and M. K.; acquisition of data: R. S.; analysis and interpretation of data: R. S.; drafting of the manuscript: R. S.; critical revision of the manuscript for important intellectual content: H. R., Z. P., and R. S.; statistical analysis: R. S. and H. R.; administrative, technical, and material support: H. R. and Z. P.; study supervision: H. R. and Z. P.; major Revision of the manuscript: H. R. and Z. P.; minor revision of the manuscript: H. R., Z. P., and R. S. All authors read and approved the final manuscript.

**Conflict of Interests Statement:** All the authors declare no support from any organization for the submitted work, no financial relationships with any organizations that might have an interest in the submitted work in the previous 5 years, and no other relationships or activities that could appear to have influenced the submitted work.

**Data Availability:** The data used to support the findings of this study are available from the corresponding author upon request.

**Ethical Approval:** The study protocol was approved by the deputy of Research and Technology of Isfahan University of medical sciences, Isfahan, Iran (IR.MUI.MED.REC.1399.344 ).

**Funding/Support:** This research was not financially supported.

**Informed Consent:** Written informed consent was obtained from the participant.

### References

1. Michaeli O, Kassis I, Shachor-Meyouhas Y, Shahar E, Ravid S. Long-term motor and cognitive outcome of acute encephalitis. *Pediatrics*. 2014;**133**(3):e546-52. [PubMed ID: 24534397]. <https://doi.org/10.1542/peds.2013-3010>.
2. Desena A, Graves D, Warnack W, Greenberg BM. Herpes simplex encephalitis as a potential cause of anti-N-methyl-D-aspartate receptor antibody encephalitis: report of 2 cases. *JAMA Neurol*. 2014;**71**(3):344-6. [PubMed ID: 24473671]. <https://doi.org/10.1001/jamaneurol.2013.4580>.
3. Thompson C, Kneen R, Riordan A, Kelly D, Pollard AJ. Encephalitis in children. *Arch Dis Child*. 2012;**97**(2):150-61. [PubMed ID: 21715390]. <https://doi.org/10.1136/archdischild-2011-300100>.
4. Falcchek SJ. Encephalitis in the pediatric population. *Pediatr Rev*. 2012;**33**(3):122-33. quiz 33. [PubMed ID: 22383515]. <https://doi.org/10.1542/pir.33-3-122>.
5. Barari Sawadkahi R, Ahmadpour-Kacho M. Clinical Manifestations, Laboratory Findings and Outcomes of Children with Herpetic Encephalitis in Amirkola Children Hospital, Northern Iran. *Iran Red Crescent Med J (IRCMJ)*. 2010;**12**(5):568-71.
6. Asadi-Pooya AA, Simani L. Central nervous system manifestations of COVID-19: A systematic review. *J Neurol Sci*. 2020;**413**:116832. [PubMed ID: 32299017]. [PubMed Central ID: PMC7151535]. <https://doi.org/10.1016/j.jns.2020.116832>.
7. Pruitt AA. Chapter 13 - Central nervous system infections in cancer patients. In: Newton HB, Malkin MG, editors. *Neurological Complications of Systemic Cancer and Antineoplastic Therapy (Second Edition)*. Cambridge, Massachusetts: Academic Press; 2022. p. 195-222. <https://doi.org/10.1016/B978-0-12-821976-8.00014-1>.
8. Lucas MJ, Brouwer MC, van de Beek D. Neurological sequelae of bacterial meningitis. *J Infect*. 2016;**73**(1):18-27. [PubMed ID: 27105658]. <https://doi.org/10.1016/j.jinf.2016.04.009>.
9. Sweeney MD, Sagare AP, Zlokovic BV. Blood-brain barrier breakdown in Alzheimer disease and other neurodegenerative disorders. *Nat Rev Neurol*. 2018;**14**(3):133-50. [PubMed ID: 29377008]. [PubMed Central ID: PMC5829048]. <https://doi.org/10.1038/nrneurol.2017.188>.
10. Lee MR, Sheng WH, Hung CC, Yu CJ, Lee LN, Hsueh PR. Mycobacterium abscessus Complex Infections in Humans. *Emerg Infect Dis*. 2015;**21**(9):1638-46. [PubMed ID: 26295364]. [PubMed Central ID: PMC4550155]. <https://doi.org/10.3201/2109.141634>.
11. Varni JW, Burwinkle TM, Seid M, Skarr D. The PedsQL 4.0 as a pediatric population health measure: feasibility, reliability, and validity. *Ambul Pediatr*. 2003;**3**(6):329-41. [PubMed ID: 14616041]. [https://doi.org/10.1367/1539-4409\(2003\)003<0329:tpaapp>2.0.co;2](https://doi.org/10.1367/1539-4409(2003)003<0329:tpaapp>2.0.co;2).

12. Eiser C, Morse R. A review of measures of quality of life for children with chronic illness. *Arch Dis Child*. 2001;**84**(3):205-11. [PubMed ID: 11207164]. [PubMed Central ID: PMC1718699]. <https://doi.org/10.1136/adc.84.3.205>.
13. Hossain MA, Shafin R, Ahmed MS, Rana MS, Walton LM, Raigangar V, et al. Health-related quality of life and coping strategies adopted by COVID-19 survivors: A nationwide cross-sectional study in Bangladesh. *PLoS One*. 2022;**17**(11). e0277694. [PubMed ID: 36383545]. [PubMed Central ID: PMC9668139]. <https://doi.org/10.1371/journal.pone.0277694>.
14. Liu E, Twilt M, Tyrrell PN, Dropol A, Sheikh S, Gorman M, et al. Health-related quality of life in children with inflammatory brain disease. *Pediatr Rheumatol Online J*. 2018;**16**(1):73. [PubMed ID: 30458827]. [PubMed Central ID: PMC6245877]. <https://doi.org/10.1186/s12969-018-0291-4>.
15. Rao S, Elkon B, Flett KB, Moss AF, Bernard TJ, Stroud B, et al. Long-Term Outcomes and Risk Factors Associated With Acute Encephalitis in Children. *J Pediatric Infect Dis Soc*. 2017;**6**(1):20-7. [PubMed ID: 26553786]. <https://doi.org/10.1093/jpids/piv075>.
16. Gigi M, Roth J, Eshel R, Constantini S, Bassan H. Health-related quality of life after post-haemorrhagic hydrocephalus in children born preterm. *Dev Med Child Neurol*. 2019;**61**(3):343-9. [PubMed ID: 30187913]. <https://doi.org/10.1111/dmcn.14012>.
17. Chou IC, Lin CC, Kao CH. Enterovirus Encephalitis Increases the Risk of Attention Deficit Hyperactivity Disorder: A Taiwanese Population-based Case-control Study. *Med (Baltimore)*. 2015;**94**(16). e707. [PubMed ID: 25906098]. [PubMed Central ID: PMC4602682]. <https://doi.org/10.1097/MD.0000000000000707>.
18. Lin CH, Lin WD, Chou IC, Lee IC, Hong SY. Epilepsy and Neurodevelopmental Outcomes in Children With Etiologically Diagnosed Central Nervous System Infections: A Retrospective Cohort Study. *Front Neurol*. 2019;**10**:528. [PubMed ID: 31156546]. [PubMed Central ID: PMC6529952]. <https://doi.org/10.3389/fneur.2019.00528>.
19. Salih MA, El Khashab HY, Hassan HH, Kentab AY, Al Subaei SS, Zeidan RM, et al. A Study on Herpes Simplex Encephalitis in 18 Children, Including 3 Relapses. *Open Pediatr Med J*. 2009;**3**(1):48-57. <https://doi.org/10.2174/1874309900903010048>.
20. Costa BKD, Sato DK. Viral encephalitis: a practical review on diagnostic approach and treatment. *J Pediatr (Rio J)*. 2020;**96** Suppl 1(Suppl 1):12-9. [PubMed ID: 31513761]. [PubMed Central ID: PMC9431993]. <https://doi.org/10.1016/j.jped.2019.07.006>.
21. Wang IJ, Lee PI, Huang LM, Chen CJ, Chen CL, Lee WT. The correlation between neurological evaluations and neurological outcome in acute encephalitis: a hospital-based study. *Eur J Paediatr Neurol*. 2007;**11**(2):63-9. [PubMed ID: 17240177]. <https://doi.org/10.1016/j.ejpn.2006.09.010>.