



# Investigation of the Most Common Specimens in the Pediatric Hospital in Kirkuk and the Epidemiology of Predominant Bacteria Using Multiple Antibiotic Resistance Indices

Abbas Hameed Sh. Al-Wandawy <sup>1,\*</sup>, Luma Abdulhady Zwain <sup>2</sup>

<sup>1</sup> Directorate of Education of Kirkuk, Ministry of Education, Kirkuk, Iraq

<sup>2</sup> Department of Biology, Collage of Education Pure Science Ibn Al-Haitham, University of Baghdad, Baghdad, Iraq

\*Corresponding Author: Directorate of Education of Kirkuk, Ministry of Education, Kirkuk, Iraq. Email: [abbas.wandawy@uokirkuk.edu.iq](mailto:abbas.wandawy@uokirkuk.edu.iq)

Received: 12 December, 2023; Revised: 16 November, 2024; Accepted: 12 December, 2024

## Abstract

**Background:** Urinary tract infections (UTIs) are among the most prevalent infectious diseases in children and are a leading cause of antibiotic use and pediatric hospitalization.

**Objectives:** This study aimed to investigate the most common specimens and determine the frequency of Multiple Antibiotic Resistance Indexes (MARI) for predominant bacteria in the pediatric hospital of Kirkuk.

**Methods:** A total of 299 different samples were collected from the pediatric hospital in Kirkuk province between May 1st and October 1st. The samples were cultured on blood agar, mannitol agar, and MacConkey agar and incubated for 24 hours at 37°C. Identification was based on morphological and microscopic examination, as well as the API kit.

**Results:** The results showed that most of the samples were urine, with 221 samples collected. Of these, 66 (28.50%) showed positive growth, and 155 (71.49%) showed negative growth. Blood samples were the second most common, with 51 samples collected, of which 4 (9.61%) were positive, and 47 (90.38%) were negative. Stool samples amounted to 13, with 2 (15.38%) showing positive growth, and 11 (84.61%) showing negative growth. Cerebrospinal fluid (CSF), throat swab, and vaginal swab samples amounted to 11, 2, and 1, respectively, with no bacterial growth observed in them. The majority of isolates were from urine, with *Escherichia coli* being the most common species (28 isolates, 42.42%). Other species isolated included *Klebsiella pneumoniae* (13 isolates, 19.69%), *Staphylococcus* spp. (12 isolates, 19.05%), *Enterobacter* spp. (7 isolates, 10.61%), and *Pseudomonas* spp. (4 isolates, 6.06%). *Proteus* spp. and *Streptococcus* spp. each accounted for 1 isolate (1.25%). The bacteria isolated from blood were *Acinetobacter* spp., *Enterobacter* spp., *Pseudomonas* spp., and *Klebsiella* spp., each representing 1 isolate (1.25%) of the total. Two (100%) *E. coli* isolates were obtained from stool samples. *E. coli* isolates from urine were resistant to ampicillin (Am), tetracycline (Te), amoxicillin/clavulanic acid (AMC), and doxycycline (Do). *Klebsiella* spp. were resistant to Am, *Staphylococcus* spp. to erythromycin (E) and azithromycin (AZM), *Enterobacter* spp. to amikacin (AK) and AMC, and *Pseudomonas aeruginosa* to gentamicin (CN), all exhibiting 100% resistance. The MARI was greater than 0.2 in 27 isolates of *E. coli* (96.42%), 13 isolates of *Klebsiella* spp. (100%), 11 isolates of *Staphylococcus* spp. (91.66%), 7 isolates of *Enterobacter* spp. (100%), and 4 isolates of *P. aeruginosa* (100%).

**Conclusions:** The current study concludes that urinary tract infection is the most common pathological condition among the young age group. Urine is one of the most frequently examined samples, with *E. coli* being the most predominant pathogen. The study also highlights the prevalence of antibiotic-resistant bacteria, as evidenced by the elevated MARI in isolates of *E. coli*, *K. pneumoniae*, *Enterobacter* spp., and *Pseudomonas* spp.

**Keywords:** Predominant Bacteria, Multiple Antibiotic Resistance Index, Clinical Specimens

## 1. Background

Urinary tract infections (UTIs) are among the most prevalent infectious diseases in children and are a leading cause of antibiotic use and pediatric

hospitalization (1, 2). Urinary tract infections also represent a significant challenge for pediatricians, healthcare providers, and the management of children in emergency departments (1, 3, 4). Bacteria are the primary cause of UTIs, with fungi and some viruses

being rare contributors. Uropathogenic *Escherichia coli* is the most frequent cause of UTIs, followed by *Staphylococcus* species, *Klebsiella pneumoniae*, *Proteus mirabilis*, and *Enterococcus faecalis* (4).

*Escherichia coli* strains that cause UTIs have defense mechanisms, including a glycosylated polysaccharide capsule that protects against phagocytosis and the immune complement system (2). *Escherichia coli* is one of the most significant causative agents of UTIs, cholecystitis, bacteremia, meningitis, and traveler's diarrhea (5). More than 40% of the bacteria responsible for UTIs are resistant to some of the antibiotics used, leading to patient relapse, treatment failure, progression to more severe diseases (such as bacteremia), which requires hospitalization, and, ultimately, death (3).

Gram-negative bacteremia remains a leading cause of mortality and morbidity, with its incidence rising globally (6). The presence of resistant genes on extrachromosomal DNA, primarily found in the *Enterobacteriaceae* family, is mainly responsible for the occurrence of antibiotic resistance in gram-negative bacteria that cause infections in hospitals and the community (7).

The second most common cause of death among young children is diarrhea, often resulting from pathogens transmitted from the mouth to the feces (8). *Escherichia coli* (enteropathogenic, enteroinvasive, and enteroaggregative strains) is one of the causes of diarrhea in children under five years of age (9).

## 2. Objectives

Given the increasing number of pediatric patients admitted to the children's hospital in Kirkuk, especially those with UTIs who fail to respond to treatment, the study aimed to identify the most frequent specimens and investigate the epidemiology of predominant bacteria using the Multiple Antibiotic Resistance Index (MARI).

## 3. Methods

### 3.1. Sample Collection

A total of 299 samples were collected from males and females aged less than one to 15 years at the pediatric hospital in Kirkuk Governorate, during the period from May 1, 2022, to October 1, 2023.

### 3.2. Culturing Samples

The samples were cultured and examined under a microscope using the API20E and API Staph systems after being incubated for 24 hours at 37°C on blood agar, mannitol salt agar, and MacConkey agar.

### 3.3. Bacterial Sensitivity Study

Bacterial sensitivity was determined using the Kirby-Bauer disk diffusion method, where the inhibition zone was measured using a ruler (in millimeters).

### 3.4. Multiple Antibiotic Resistance Index

Multiple Antibiotic Resistance Index was calculated and interpreted as described by Krumperman (10), applying the formula  $a/b$ , where "b" represents the total number of antibiotics tested and "a" represents the number of antibiotics to which a single isolate is resistant (11).

### 3.5. Ethical Approval

Ethical approval was obtained from the Scientific Research Ethics Committee of the Ministry of Health - Kirkuk Health Department under Order 1263, dated January 2022.

## 4. Results

A total of 299 samples were collected from suspected patients, as diagnosed by the examining physician, at the pediatric hospital in Kirkuk Governorate, between May 1, 2022, and October 1, 2023. As shown in Table 1, most of the samples were urine, with a total of 221 samples, of which 66 (28.50%) showed positive growth and 155 (71.49%) showed negative growth. Blood samples followed, with 51 samples, 4 (9.61%) of which were positive growth and 47 (90.38%) were negative growth. Stool samples accounted for 13 samples, 2 (15.38%) of which were positive growth and 11 (84.61%) were negative growth. The numbers of Cerebrospinal fluid (CSF), throat swab, and vaginal swab samples were 1, 2, and 11, respectively, with no bacterial growth observed in any of them (Table 2).

### 4.1. Resistance of the Most Isolated Bacteria from Urine to Antibiotics

Figure 1 shows the antibiotic resistance profile of *E. coli* bacteria isolated from urine. The bacteria exhibited 100% resistance to the antibiotics ampicillin (Am), tetracycline (Te), amoxicillin/clavulanic acid (AMC), and doxycycline (Do). The resistance percentages for the following antibiotics were as follows: Azithromycin (AZM) (76.47%), ciprofloxacin (CIP) (28.57%),

**Table 1.** Percentage of Bacterial Growth According to Sources of Isolation

Isolation Source	Total Number	Bacterial Growth <sup>a</sup>	
		Positive Growth	Negative Growth
Urine	221	66 (28.50)	155 (71.49)
Blood	51	4 (9.61)	47 (90.38)
Stool	13	2 (15.38)	11 (84.61)
CSF	11	-	11 (100)
Throat swab	2	-	2 (100)
Vaginal swab	1	-	1 (100)

Abbreviation: CSF, cerebrospinal fluid.

<sup>a</sup> Values are expressed as No. (%).**Table 2.** Percentage of Isolates According to Sources of Isolation

Source of Isolation (No.) and Bacterial Species	No. (%)
<b>Urine (66)</b>	
<i>Escherichia coli</i>	28 (42.42)
<i>Klebsiella pneumoniae</i>	13 (19.696)
<i>Staphylococcus</i> spp.	12 (19.047)
<i>Enterobacter</i> spp.	7 (10.606)
<i>Pseudomonas</i> spp.	4 (6.06)
<i>Proteus</i> spp.	1 (1.515)
<i>Streptococcus</i> spp.	1 (1.515)
No identification	6 (9.09)
<b>Blood (4)</b>	
<i>Acinitobacter</i> spp.	1 (25)
<i>Enterobacter</i> spp.	1 (25)
<i>Pseudomonas</i> spp.	1 (25)
<i>Klebsiella</i> spp.	1 (25)
<b>Stool (2)</b>	
<i>E. coli</i>	2 (100)
<b>CSF (0)</b>	-
<b>Throat swab (0)</b>	-
<b>Vaginal swab (0)</b>	-

Abbreviation: CSF, cerebrospinal fluid.

trimethoprim-sulfamethoxazole (SXT) (78.57%), cephalexin (CN) (52.63%), amikacin (AK) (66.66%), cefixime (CFM) (69.23%), levofloxacin (LEV) (25%), ceftazidime (CAZ) (75%), aztreonam (ATM) (56%), cefepime (FEP) (81.48%), nalidixic acid (NA) (58.33%), nitrofurantoin (F) (26.66%), and imipenem (IPM) (7.14%).

The bacteria showed resistance to AZM, CIP, SXT, AM, AMC, CAZ, ATM, FEP, DO, and F, with resistance percentages of 91.66%, 14.28%, 5%, 100%, 91.66%, 81.81%, 72.72%, 72.72%, 85.71%, and 55.55%, respectively (Figure 2).


The isolates showed 100% resistance to erythromycin (E) and AZM, and resistance to CIP, CN, DO, vancomycin

(VA), and rifampin (RA) at percentages of 37.5%, 70%, 87.5%, 60%, and 14.28%, respectively (Figure 3).


The isolates exhibited 100% resistance to AK and AMC, and resistance to CIP, SXT, CFM, CAZ, ATM, and cefepime (FEP) at percentages of 25%, 16.66%, 60%, 57.14%, 42.85%, and 14.28%, respectively (Figure 4).

The isolates were resistant to CN by 100%, CIP, CAZ, FEP, and IPM by 75%, and AZM, SXT, AK, LEV, and F by 50% (Figure 5).

#### 4.2. Multiple Antibiotic Resistance Index for Bacterial Isolates from Urine



**Figure 1.** Resistance to antibiotics in *Escherichia coli* isolated from urine. AZM, azithromycin; CIP, ciprofloxacin; SXT, trimethoprim-sulfamethoxazole; AM, ampicillin; CN, cephalixin; AK, amikacin; TE, tetracycline; CFM, cefixime; LEV, levofloxacin; AMC, amoxicillin-clavulanate; CAZ, ceftazidime; ATM, aztreonam; FEP, cefepime; NA, nalidixic acid; DO, doxycycline; F, nitrofurantoin; IPM, imipenem.



**Figure 2.** Resistance to antibiotics in *Klebsiella* spp. isolated from urine. AZM, azithromycin; CIP, ciprofloxacin; SXT, trimethoprim-sulfamethoxazole; AM, ampicillin; AMC, amoxicillin-clavulanate; CAZ, ceftazidime; ATM, aztreonam; FEP, cefepime; DO, doxycycline; F, nitrofurantoin.


Table 3 shows the MARI for bacterial isolates from urine. The MARI was greater than 0.2 in 27 isolates of *E. coli* (96.42%), 13 isolates of *Klebsiella* spp. (100%), 11 isolates of *Staphylococcus* spp. (91.66%), 7 isolates of *Enterobacter* spp. (100%), and 4 isolates of *Pseudomonas aeruginosa* (100%).

## 5. Discussion


Overprescription of antibiotics is common in primary healthcare facilities. To reduce the likelihood of

antibiotic resistance developing, health authorities should strictly regulate or outright prohibit the overprescription of antibiotics (12).

It is reported that *E. coli* (64.5%), *Klebsiella* species (11.6%), and *Enterococcus* species (6.1%) were the most common bacteria isolated from urine samples (13). This was supported by Luo et al. (14), which indicated that gram-negative bacteria were the most frequently isolated species in UTIs. A study highlighted the predominance of *E. coli* bacteria in patients suffering from UTIs (15).



**Figure 3.** Resistance to antibiotics in *Staphylococcus* spp. isolated from urine. AZM, azithromycin; CIP, ciprofloxacin; E, erythromycin; CN, cephalixin; DO, doxycycline; VA, vancomycin; RA, rifampin.




**Figure 4.** Resistance to antibiotics in *Enterobacter* spp. isolated from urine. CIP, ciprofloxacin; SXT, trimethoprim-sulfamethoxazole; AK, amikacin; CFM, cefixime; AMC, amoxicillin-clavulanate; CAZ, ceftazidime; ATM, aztreonam; FEP, cefepime.

The results indicated that bacteria isolated from urine samples exhibited resistance to many antibiotics, classifying them as multidrug-resistant (MDR) bacteria. This was confirmed by a study (16), who recorded the resistance of gram-negative bacteria to each of the following antibiotics: Gentamicin, Am/sulbactam, CIP, IM, FEP, LEV, ATM, and AK, with resistance rates of 69.56%, 56.52%, 43.47%, 0%, 52.17%, 82.60%, 60.87%, and 39.13%, respectively.

Similarly, Islam et al. (17) found that *E. coli* and *Klebsiella* species were resistant to a range of antibiotics, including sulfonamides (56%, 41%), fluoroquinolones (69%, 53%), third-generation cephalosporins (69%, 58%), macrolides (70%, 76%), and penicillin (85%, 95%), respectively.

According to a study (18), a significant proportion of antibiotic resistance was observed in *K. pneumoniae* and *E. coli*. *K. pneumoniae* demonstrated high resistance to



**Figure 5.** *Enterobacter* species isolated from urine that are resistant to antibiotics. AZM, azithromycin; CIP, ciprofloxacin; SXT, trimethoprim-sulfamethoxazole; CN, cephalixin; AK, amikacin; LEV, levofloxacin; AMC, amoxicillin-clavulanate; CAZ, ceftazidime; FEP, cefepime; F, nitrofurantoin; IPM, imipenem.

**Table 3.** Multiple Antibiotic Resistance Index of Bacterial Isolates from Urine <sup>a</sup>

MARI	<i>Escherichia coli</i>	<i>Klebsiella</i> Species	<i>Staphylococcus</i> Species	<i>Enterobacter</i> Species	<i>Pseudomonas aeruginosa</i>
< 0.2	1 (3.57)	0 (0)	1 (8.33)	0 (0)	0 (0)
0.2 - 0.6	18 (64.28)	8 (61.53)	6 (50)	6 (85.71)	0 (0)
0.7 - 0.9	7 (25)	5 (38.46)	3 (25)	0 (0)	4 (100)
1 <	2 (7.14)	0 (0)	2 (16.66)	1 (14.28)	0 (0)

<sup>a</sup> Values are expressed as No. (%).

streptomycin (88%), E (88%), cloxacillin (96%), while *E. coli* exhibited resistance to all three antibiotics.

The antibiotics IM (95%), STX (69.8%), ATM (60.5%), chloramphenicol (45.3%), and meropenem (27.9%) were more effective against *Pseudomonas* isolates (19).

Regarding the resistance of *Pseudomonas* bacteria, the results of a study (19) indicated that *Pseudomonas* isolates exhibited greater resistance to IM (95%), STX (69.8%), ATM (60.5%), chloramphenicol (45.3%), and meropenem (27.9%).

In contrast, Mapipa et al. (20) reported that the antibiotic ceftazidime had the highest resistance rate (63%) against *P. aeruginosa* isolates, with resistance to other antibiotics ranging between 7% and 35%.

Regarding the antibiotic resistance of *Staphylococcus* bacteria, a study (21) indicated that all *Staphylococcus*

isolates were 100% resistant to AZM and E, and 95.56% resistant to cefixime, 50% resistant to Am, and 95% resistant to amoxicillin. Meanwhile, another study (22) indicated that all *Staphylococcus aureus* isolates were 100% resistant to Am and penicillin, 97.6% resistant to AK, and 90% resistant to ciprofloxacin and gentamicin.

According to a study (23), the development of quinolone resistance in *Enterobacteriaceae* is a complex and multifaceted process. The primary resistance mechanisms include one or more genetic mutations at the target site that alter the drug's affinity for binding to target enzymes, overexpression of the AcrAB-TolC MDR efflux pumps, and decreased expression of porins and plasmid-coded resistance proteins, such as the protection protein Qnr.

A study assessed gene sequencing in *E. coli* bacteria and indicated the presence of the genes orf00490,



orf00819, orf001916, and orf01200, which regulate the expression of the enzyme fumarate reductase subunit D (frdD) as well as the cell division protein FtsI (penicillin-binding protein 3) (24). Additionally, the outer membrane porin protein OmpD is controlled by the genes orf00490, orf00819, and orf001916.

The occurrence of several mutations in these genes leads to bacterial resistance to many antibiotics, while the gene orf04094 expresses histidine kinase, orf02235 expresses multidrug resistance, and orf03479 expresses valine-glycine repeat G, which is excreted through the type VI secretion system (T6SS), one of the extracellular substances that contribute to antibiotic resistance.

The resistance of *P. aeruginosa* to cephalosporins, carbapenems, aminoglycosides, and fluoroquinolones is primarily due to horizontal gene transfer, involving integrons, plasmids, and transposons (25).

It is noted from the results that the MARI of bacteria isolated from urine was mostly greater than 0.2, indicating the extent of the epidemic of each of the bacteria (*E. coli*, *K. pneumoniae*, *Enterobacter* spp., *Pseudomonas* spp.).

This is supported by Ayandele et al. (18), which found that *E. coli* had the highest MAR index, reaching 1.00 in 17 isolates that showed resistance to 14 antibiotics. Meanwhile, another study (26) indicated in their study that 44% of *Staphylococcus* spp. isolates and 50% of *E. coli* isolates had MAR indices greater than 0.2.

In a study, the MARI of *Pseudomonas* was 0.85 (27), while a study by Ayandele et al. (18) showed in their study that the MARI in *Pseudomonas* isolates ranged between 0.0 and 0.8.

As for *Enterococcus* bacteria, a study (28), indicated in their study that the MARI for *Enterococcus* ranged between 0.08 and 0.83, while another study (20) showed that the MARI for *P. aeruginosa* ranged between 0.23 and 0.38.

Multiple Antibiotic Resistance Index is a useful tool for assessing the vulnerability of humans to the risk of resistant bacteria, as well as the extent of environmental danger due to antibiotic-resistant bacteria (10). This is especially important because antibiotic-resistant environmental bacteria can evolve into pathogens through genetic association (29) and phenotypic diversity between environmental and clinical bacteria (30). Additionally another study (31) indicated that a rise in MARI to 0.82, 0.73, and 0.64 signifies significant contamination with *Vibrio parahaemolyticus* bacteria. The MARI for *Pseudomonas putida* isolated from fish was 0.76, indicating high antibiotic use in fish farms (32). The rise of MARI above 0.2 signifies the extensive use of antibiotics in the aquatic environment. This variation in

the MARI of isolated bacteria is attributed to mobile genetic elements, particularly Class 1 integrons, which provide a significant opportunity for the spread of antibiotic resistance among fish.

While Mishra et al. (33) indicated that *Enterobacter* with an MARI above 0.3 was observed to possess outer membrane proteins as well as other virulence factors, such as Type 1 fimbriae, biofilm production, and serum resistance.

This is supported by a study (34), which noted that there is a difference in MARI between bacterial genera. *Pseudomonas aeruginosa* had a higher MARI than *Klebsiella* spp. and *Proteus* spp., which in turn had a higher MARI than *Enterobacter* spp. The study also pointed out that 25% of the isolates posed a significant risk to humans and animals, with the majority being *P. aeruginosa*, which poses a higher threat as a potential pathogen compared to *Enterobacter* spp., based on MARI and virulence factors.

It was indicated that bacterial isolates with an MARI greater than 0.3 and a virulence factor above 0.5 pose a significant threat, while isolates with an MARI less than 0.3 and a virulence factor less than 0.5 represent a medium risk (34). Isolates with low risk have both an MARI less than or equal to 0.3 and a virulence factor greater than or equal to 0.5. Non-risk isolates are those with an MARI greater than 0.3 and a virulence factor greater than 0.5.

It is noted from the results that the values of MARI vary between genera as well as between the same bacterial species. This may be due to many factors, as indicated by (35-37), who found that differences in MARI values depend on the source of the sample, geographical location, and the method of testing.

### 5.1. Conclusions

According to the current study, UTIs are the most common pathological cases among young people, and urine is one of the most frequently examined samples. *Escherichia coli* is the most common bacterium, and antibiotic-resistant bacteria spread through the increasing MARI of isolated strains, including *Enterobacter* species, *Pseudomonas* species, *E. coli*, and *K. pneumoniae*.

### Footnotes

**Authors' Contribution:** A. H. Sh. Al-W.: Visualization, reviewing, editing, conceptualization, and methodology; L. A. Z: Writing, investigation, software, validation, and original draft preparation.

**Conflict of Interests Statement:** The authors declare that they have no known competing financial interests or personal relationships that could have influenced the work reported in this paper.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication.

**Ethical Approval:** Ethical approval was obtained from the Scientific Research Ethics Committee of the Ministry of Health, Kirkuk Health Department, by Order 1263 in January 2022.

**Funding/Support:** This project was not financially supported.

## References

- Autore G, Bernardi L, La Scola C, Ghidini F, Marchetti F, Pasini A, et al. Management of Pediatric Urinary Tract Infections: A Delphi Study. *Antibiotics (Basel)*. 2022;**11**(8). [PubMed ID: 36009990]. [PubMed Central ID: PMC9404756]. <https://doi.org/10.3390/antibiotics11081122>.
- Daniel M, Szymanik-Grzelak H, Sierdzinski J, Podsiadly E, Kowalewska-Mlot M, Panczyk-Tomaszewska M. Epidemiology and Risk Factors of UTIs in Children-A Single-Center Observation. *J Pers Med*. 2023;**13**(1). [PubMed ID: 36675799]. [PubMed Central ID: PMC9865477]. <https://doi.org/10.3390/jpm13010138>.
- McCowan C, Bakhshi A, McConnachie A, Malcolm W, Barry SJ, Santiago VH, et al. E. coli bacteraemia and antimicrobial resistance following antimicrobial prescribing for urinary tract infection in the community. *BMC Infect Dis*. 2022;**22**(1):805. [PubMed ID: 36307776]. [PubMed Central ID: PMC9621144]. <https://doi.org/10.1186/s12879-022-07768-7>.
- Mancuso G, Midiri A, Gerace E, Marra M, Zummo S, Biondo C. Urinary Tract Infections: The Current Scenario and Future Prospects. *Pathogens*. 2023;**12**(4). [PubMed ID: 3711509]. [PubMed Central ID: PMC10154414]. <https://doi.org/10.3390/pathogens12040623>.
- Tanba C, Bandaru S, Haas CJ. Double Trouble: Escherichia Coli Bilateral Empyema. *Am J Respir Crit Care Med*. 2023;**203**:A2110.
- Lee IR, Tong SYC, Davis JS, Paterson DL, Syed-Omar SF, Peck KR, et al. Early oral stepdown antibiotic therapy versus continuing intravenous therapy for uncomplicated Gram-negative bacteraemia (the INVEST trial): study protocol for a multicentre, randomised controlled, open-label, phase III, non-inferiority trial. *Trials*. 2022;**23**(1):572. [PubMed ID: 35854360]. [PubMed Central ID: PMC9295110]. <https://doi.org/10.1186/s13063-022-06495-3>.
- Kundar R, Gokarn K. CRISPR-Cas System: A Tool to Eliminate Drug-Resistant Gram-Negative Bacteria. *Pharmaceuticals (Basel)*. 2022;**15**(12). [PubMed ID: 36558949]. [PubMed Central ID: PMC9781512]. <https://doi.org/10.3390/ph15121498>.
- Simanjuntak DF, Kusumawati RL, Bader O, Luder CGK, Zimmermann O, Gross U. A comparative pilot study on Gram-negative bacteria contaminating the hands of children living in urban and rural areas of Indonesia versus Germany - A suitable monitoring strategy for diarrhea risk assessment? *Front Microbiol*. 2023;**14**:1152411. [PubMed ID: 37077245]. [PubMed Central ID: PMC10106674]. <https://doi.org/10.3389/fmicb.2023.1152411>.
- Shrestha SK, Shrestha J, Mason CJ, Sornsakrin S, Dhakhwa JR, Shrestha BR, et al. Etiology of Acute Diarrheal Disease and Antimicrobial Susceptibility Pattern in Children Younger Than 5 Years Old in Nepal. *Am J Trop Med Hyg*. 2023;**108**(1):174-80. [PubMed ID: 36509064]. [PubMed Central ID: PMC9833095]. <https://doi.org/10.4269/ajtmh.21-1219>.
- Krumperman PH. Multiple antibiotic resistance indexing of Escherichia coli to identify high-risk sources of fecal contamination of foods. *Appl Environ Microbiol*. 1983;**46**(1):165-70. [PubMed ID: 6351743]. [PubMed Central ID: PMC239283]. <https://doi.org/10.1128/aem.46.1.165-170.1983>.
- Mir R, Salari S, Najimi M, Rashki A. Determination of frequency, multiple antibiotic resistance index and resistotype of Salmonella spp. in chicken meat collected from southeast of Iran. *Vet Med Sci*. 2022;**8**(1):229-36. [PubMed ID: 34597476]. [PubMed Central ID: PMC8788964]. <https://doi.org/10.1002/vms3.647>.
- Nafea Abed Aijalii RW, Wadi Algawwam HG. Evaluation of over prescription of antibiotics for upper respiratory tract infections in children < 12 years old in Kirkuk, Iraq. *Iraq Med J*. 2019;**3**(3).
- Wanke-Rytt M, Sobierajski T, Lachowicz D, Seliga-Gasior D, Podsiadly E. Analysis of Etiology of Community-Acquired and Nosocomial Urinary Tract Infections and Antibiotic Resistance of Isolated Strains: Results of a 3-Year Surveillance (2020-2022) at the Pediatric Teaching Hospital in Warsaw. *Microorganisms*. 2023;**11**(6). [PubMed ID: 37374940]. [PubMed Central ID: PMC10301861]. <https://doi.org/10.3390/microorganisms11061438>.
- Luo XM, Li YY, Qi XM, Wu YG. Risk factors and etiological characteristics of urinary tract infection in hospitalized continuous ambulatory peritoneal dialysis patients. *Eur Rev Med Pharmacol Sci*. 2023;**27**(9):3837-45. [PubMed ID: 37203808]. [https://doi.org/10.26355/eurrev\\_202305\\_32289](https://doi.org/10.26355/eurrev_202305_32289).
- Wali MR, Zwain LA, Al-Wandawy AH. Study of Antibiotic-resistant Bacteria Isolated from Children with Urinary Tract Infection. *Int J Drug Delivery Technol*. 2023;**13**(1):150-7. <https://doi.org/10.25258/ijddt.13.1.23>.
- Li Y, Jiang L, Luo S, Hu D, Zhao X, Zhao G, et al. Analysis of Characteristics, Pathogens and Drug Resistance of Urinary Tract Infection Associated with Long-Term Indwelling Double-J Stent. *Infect Drug Resist*. 2023;**16**:2089-96. [PubMed ID: 37063938]. [PubMed Central ID: PMC10094401]. <https://doi.org/10.2147/IDR.S392857>.
- Islam MA, Islam MR, Khan R, Amin MB, Rahman M, Hossain MI, et al. Prevalence, etiology and antibiotic resistance patterns of community-acquired urinary tract infections in Dhaka, Bangladesh. *PLoS One*. 2022;**17**(9). e0274423. [PubMed ID: 36107878]. [PubMed Central ID: PMC9477272]. <https://doi.org/10.1371/journal.pone.0274423>.
- Ayandele AA, Oladipo EK, Oyeibisi O, Kaka MO. Prevalence of Multi-Antibiotic Resistant Escherichia coli and Klebsiella species obtained from a Tertiary Medical Institution in Oyo State, Nigeria. *Qatar Med J*. 2020;**2020**(1):9. [PubMed ID: 32280610]. [PubMed Central ID: PMC7118460]. <https://doi.org/10.5339/qmj.2020.9>.
- Meng L, Liu H, Lan T, Dong L, Hu H, Zhao S, et al. Antibiotic Resistance Patterns of Pseudomonas spp. Isolated From Raw Milk Revealed by Whole Genome Sequencing. *Front Microbiol*. 2020;**11**:1005. [PubMed ID: 32655503]. [PubMed Central ID: PMC7326020]. <https://doi.org/10.3389/fmicb.2020.01005>.
- Mapipa Q, Digban TO, Nnolim NE, Nwodo UU. Antibioqram profile and virulence signatures of Pseudomonas aeruginosa isolates recovered from selected agrestic hospital effluents. *Sci Rep*. 2021;**11**(1):11800. [PubMed ID: 34083705]. [PubMed Central ID: PMC8175747]. <https://doi.org/10.1038/s41598-021-91280-6>.
- Urmi MR, Ansari WK, Islam MS, Sobur MA, Rahman M, Rahman MT. Antibiotic resistance patterns of Staphylococcus spp. isolated from fast foods sold in different restaurants of Mymensingh, Bangladesh. *J Adv Vet Anim Res*. 2021;**8**(2):274-81. [PubMed ID: 34395598]. [PubMed Central ID: PMC8280991]. <https://doi.org/10.5455/javar.2021.h512>.



22. Yusuf ST, Kwaga JKP, Okolocha EC, Bello M. Phenotypic occurrence of methicillin-resistant *Staphylococcus aureus* in camels slaughtered at Kano abattoir, Kano, Nigeria. *Sokoto J Veterinary Sci.* 2017;**15**(2). <https://doi.org/10.4314/sokjvs.v15i2.4>.
23. Azargun R, Gholizadeh P, Sadeghi V, Hosainzadegan H, Tarhriz V, Memar MY, et al. Molecular mechanisms associated with quinolone resistance in Enterobacteriaceae: review and update. *Trans R Soc Trop Med Hyg.* 2020;**114**(10):770-81. [PubMed ID: 32609840]. <https://doi.org/10.1093/trstmh/traa041>.
24. Li M, Liu Q, Teng Y, Ou L, Xi Y, Chen S, et al. The resistance mechanism of *Escherichia coli* induced by ampicillin in laboratory. *Infect Drug Resist.* 2019;**12**:2853-63. [PubMed ID: 31571941]. [PubMed Central ID: PMC6750165]. <https://doi.org/10.2147/IDR.S221212>.
25. Nguyen L, Garcia J, Gruenberg K, MacDougall C. Multidrug-Resistant *Pseudomonas* Infections: Hard to Treat, But Hope on the Horizon? *Curr Infect Dis Rep.* 2018;**20**(8):23. [PubMed ID: 29876674]. <https://doi.org/10.1007/s11908-018-0629-6>.
26. Sombie JIN, Kagira J, Maina N. Prevalence and Antibigram of *Escherichia coli* and *Staphylococcus* spp. Isolated from Cattle Milk Products Sold in Juja Sub-County, Kenya. *J Trop Med.* 2022;**2022**:5251197. [PubMed ID: 36452460]. [PubMed Central ID: PMC9705080]. <https://doi.org/10.1155/2022/5251197>.
27. Puia D, Gheorghinca S, Priscop C. The Antimicrobial Resistance Index and Fournier Gangrene Severity Index of Patients Diagnosed with Fournier's Gangrene in a Tertiary Hospital in North Eastern Romania. *Med (Kaunas).* 2023;**59**(4). [PubMed ID: 37109603]. [PubMed Central ID: PMC10144816]. <https://doi.org/10.3390/medicina59040643>.
28. Roy K, Islam MS, Paul A, Ievy S, Talukder M, Sobur MA, et al. Molecular detection and antibiotyping of multi-drug resistant *Enterococcus faecium* from healthy broiler chickens in Bangladesh. *Vet Med Sci.* 2022;**8**(1):200-10. [PubMed ID: 34786882]. [PubMed Central ID: PMC8788975]. <https://doi.org/10.1002/vms3.669>.
29. Martins VV, Pitondo-Silva A, Manco Lde M, Falcao JP, Freitas Sdos S, da Silveira WD, et al. Pathogenic potential and genetic diversity of environmental and clinical isolates of *Pseudomonas aeruginosa*. *APMIS.* 2014;**122**(2):92-100. [PubMed ID: 23879442]. <https://doi.org/10.1111/apm.12112>.
30. Grosso-Becerra MV, Santos-Medellin C, Gonzalez-Valdez A, Mendez JL, Delgado G, Morales-Espinosa R, et al. *Pseudomonas aeruginosa* clinical and environmental isolates constitute a single population with high phenotypic diversity. *BMC Genomics.* 2014;**15**:318. [PubMed ID: 24773920]. [PubMed Central ID: PMC4234422]. <https://doi.org/10.1186/1471-2164-15-318>.
31. Ashrafudoulla M, Mizan MFR, Park H, Byun KH, Lee N, Park SH, et al. Genetic Relationship, Virulence Factors, Drug Resistance Profile and Biofilm Formation Ability of *Vibrio parahaemolyticus* Isolated From Mussel. *Front Microbiol.* 2019;**10**:513. [PubMed ID: 30949142]. [PubMed Central ID: PMC6435529]. <https://doi.org/10.3389/fmicb.2019.00513>.
32. Preena PG, Dharmaratnam A, Raj NS, Kumar TVA, Raja SA, Swaminathan TR. Antibiotic susceptibility pattern of bacteria isolated from freshwater ornamental fish, guppy showing bacterial disease. *Biologia.* 2019;**74**(8):1055-62. <https://doi.org/10.2478/s11756-019-00261-8>.
33. Mishra M, Panda S, Barik S, Sarkar A, Singh DV, Mohapatra H. Antibiotic Resistance Profile, Outer Membrane Proteins, Virulence Factors and Genome Sequence Analysis Reveal Clinical Isolates of *Enterobacter* Are Potential Pathogens Compared to Environmental Isolates. *Front Cell Infect Microbiol.* 2020;**10**:54. [PubMed ID: 32154188]. [PubMed Central ID: PMC7047878]. <https://doi.org/10.3389/fcimb.2020.00054>.
34. Singh SK, Ekka R, Mishra M, Mohapatra H. Association study of multiple antibiotic resistance and virulence: a strategy to assess the extent of risk posed by bacterial population in aquatic environment. *Environ Monit Assess.* 2017;**189**(7):320. [PubMed ID: 28589461]. <https://doi.org/10.1007/s10661-017-6005-4>.
35. Tunung R, Margaret S, Jeyaletchumi P, Chai LC, Tuan Zainazor TC, Ghazali FM, et al. Prevalence and quantification of *Vibrio parahaemolyticus* in raw salad vegetables at retail level. *J Microbiol Biotechnol.* 2010;**20**(2):391-6. [PubMed ID: 20208446].
36. Lesley MB, Velnetti L, Cheah YK, Son R, Kasing A, Samuel L, et al. Antibiotic resistance and plasmid profiling of *Vibrio parahaemolyticus* isolated from cockles (*Anadara granosa*) at Tanjung Karang, Kuala Selangor. *Int Food Res J.* 2011;**18**(3).
37. Robert-Pillot A, Guenole A, Lesne J, Delesmont R, Fournier JM, Quilici ML. Occurrence of the *tdh* and *trh* genes in *Vibrio parahaemolyticus* isolates from waters and raw shellfish collected in two French coastal areas and from seafood imported into France. *Int J Food Microbiol.* 2004;**91**(3):319-25. [PubMed ID: 14984780]. <https://doi.org/10.1016/j.ijfoodmicro.2003.07.006>.