



Plyometric Training Enhances Proprioception: A Systematic Review

Razieh Yousefian Molla ^{1,*}, Taian Martins Vieira^{2,3}

¹ Department of Sports Biomechanics, Central Tehran Branch, Islamic Azad University, Tehran, Iran

² Laboratory for Engineering of the Neuromuscular System, Department of Electronics and Telecommunications, Politecnico di Torino, Torino, Italy

³ PolitoBIOMed Lab-Biomedical Engineering Lab, Department of Electronics and Telecommunications, Politecnico di Torino, Torino, Italy

*Corresponding Author: Department of Sports Biomechanics, Central Tehran Branch, Islamic Azad University, Tehran, Iran. Email: raziehyousefianmolla@gmail.com

Received: 10 January, 2024; Revised: 12 July, 2024; Accepted: 24 July, 2024

Abstract

Context: This systematic review investigated the effects of plyometric exercise interventions on proprioception.

Methods: An electronic search was conducted in October 2022 across eight databases: PubMed, Scopus, ProQuest, Science Direct, Google Scholar, DOAJ, IEEE Xplore, and Cochrane. To be included in the review, studies had to meet all the established inclusion criteria. Twelve studies were included in the final analysis, and their quality was assessed using the Downs and Black checklist.

Results: The review found that the included studies considered various plyometric training protocols and methods for assessing proprioception. Despite differences in training regimens and assessment metrics, the data consistently showed a positive effect of plyometric training on proprioception.

Conclusions: The review concluded that plyometric exercises improve proprioception, regardless of the specific type of training or assessment method used. A primary limitation identified was the scarcity of randomized controlled trials (RCTs) that focus on plyometric exercises and their impact on proprioception.

Keywords: Stretch-Shortening Cycle, Joint Position Sense, Kinesthesia, Skeletal Muscle, Performance

1. Context

Proprioception is defined as the ability to perceive body position and movement in space through the integration of sensory signals from various mechanoreceptors (1, 2). It is well established that proprioception plays a crucial role in joint position awareness, balance control, and other motor functions (3-5). Previous research has demonstrated that proprioception also significantly influences athletic performance, making it a determinant of success in sports (6, 7). Proprioception supports the motor system by maintaining tendon tonicity (8) and enhances strength, accuracy, and movement control, which are vital in both sports and general movement (3).

In the existing literature, there is some debate regarding the effect of physical exercise on proprioception (3). Exercises such as core stabilization (9), strength training (10), tai chi (11, 12), isokinetic

training (13, 14), sensorimotor exercises (15), neuromuscular training (16), vibration therapy (17), resistance training (18), combined training programs (19), and Pilates (20) have been reported to improve joint position discrimination. However, some studies have shown no significant effect of exercise on proprioception (21-23).

One specific category of exercise whose effect on proprioception has yet to be fully assessed is plyometric exercise. Plyometric exercises involve rapid stretching of a muscle and its connective tissue (eccentric activity), followed by a forceful concentric contraction (the stretch-shortening cycle) (24). These exercises are primarily used to enhance mechanical power during successive movements by relying on the elastic energy stored in the muscles and the stretch reflex (24, 25). Evidence suggests that plyometric training improves various physical attributes such as explosive power, jumping ability, sprinting speed, agility (26, 27), and

endurance (28). Additionally, plyometric exercises are thought to benefit the musculoskeletal system by increasing bone mass, enhancing the stiffness of muscle-tendon complexes, and improving overall muscle function (28-30). Given these benefits, it is plausible to hypothesize that plyometric exercises may also improve proprioception.

To test this hypothesis, it is necessary to conduct experiments under various conditions, as there are multiple variants of plyometric exercises and approaches for assessing proprioception. Any study that attempts to establish a causal relationship between plyometric training and proprioception would be limited to the specific experimental protocol employed.

2. Objectives

To provide more general insights, this systematic review aims to assess the influence of plyometric exercises on proprioception, either directly or indirectly, by reviewing the relevant literature.

3. Methods

This systematic review was conducted following the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines (31) (Figure 1). The review also adhered to the PICO (Participants, Intervention, Comparison, Outcome, and Study Design) framework for assessing the effect of plyometric training on proprioception (32). The study has been registered with the international online registry PROSPERO under the code [CRD42024563346](#).

3.1. Search Strategy

A comprehensive digital search was conducted in October 2022 across multiple databases, including PubMed, Scopus, ProQuest, Science Direct, Google Scholar, DOAJ, IEEE Xplore, and Cochrane. The search covered studies from 1990 to October 2022, and terms based on Medical Subject Headings (MeSH) were used. These search terms included “proprioception,” “kinesthesia,” “kinesthesia,” “kinesthetic sense,” “sense of self-movement,” “body position,” “sixth sense,” “position movement sensation,” “sense of locomotion,” “limb position,” “joint position,” “muscle sense,” “proprioceptive,” “proprioceptors,” “joint afferent,” “mechanosensation,” “mechanosensitive,” “muscle sensation,” “muscle sensibility,” “muscle sense,”

“muscular sensibility,” “myaesthesia,” “myesthesia,” “sense of equilibrium,” “sense of balance,” “interception,” “vestibular,” “myoaesthesia,” “myoesthesia,” “plyometric,” “jump training,” “plyos,” “shock method,” “jump exercise,” “jump workout,” “Stretch-Shortening Cycle,” and “Ballistic resistance training.”

3.2. Inclusion and Exclusion Criteria

To be included in this systematic review, articles had to meet the following inclusion criteria: They had to be randomized controlled trials (RCTs), have full-text articles available in English, use valid and reliable assessment tools, and report the effects of plyometric training on proprioception. The exclusion criteria included studies that were not in English, as well as book chapters, conference papers, conference posters, and dissertations.

3.3. Study Selection and Quality Assessment

No restrictions were placed on age, gender, participants, or statistical methods due to the limited number of studies available for review. Instead, the review focused on identifying the effects of plyometric exercises on proprioception. If disagreements between reviewers arose, they discussed the relevant abstracts to reach a consensus, and a third reviewer was consulted if necessary. After this process, 12 articles were selected for inclusion in the final review.

The quality of each study's methodology was assessed using a modified 27-item version of the Downs and Black Checklist (33). Items were scored as either yes = (1), no or unable to determine = (0), except for item 5, where yes = (2), partially = (1), and no = (0). The maximum possible score for each study was 28. Based on their scores, studies were classified into quality levels: Excellent (26 - 28), good (20 - 25), fair (15 - 19), and poor (≤ 14) (33). The risk of bias was independently evaluated by two reviewers, focusing on six key areas: (1) bias in selecting participants, (2) bias in classifying interventions, (3) bias due to deviations from intended interventions, (4) bias from missing data, (5) bias in measuring outcomes, and (6) bias in selective reporting of outcomes. Studies with a high risk of bias across all categories were excluded from the review (34).

4. Results




Figure 1. PRISMA flow chart

A total of 386 participants, both control and intervention groups, were included across the 12 studies assessed. Nine of these studies involved healthy individuals, while three focused on people with specific conditions: (1) multiple sclerosis (MS) (35), (2) functional ankle instability (FAI) (36), and (3) those with grade I or II unilateral inversion ankle sprain (37).

The length of interventions varied, ranging from eight days to 12 weeks, with six weeks being the most common. Most interventions were conducted with a frequency of two to three sessions per week, with three sessions being the most common. Plyometric exercises across studies included jump training, throwing, hops, dynamic lunges, and exercises like kick butt.

According to the data presented in Table 1, ten studies reported positive effects of plyometric exercises on

proprioception (35-44), while two studies found no significant effects (45, 46).

Table 2 provides the results from the Downs and Black checklist, which was used to assess the quality of the included studies.

Significant effects of proprioception on both upper and lower limbs were reported across the studies assessed (Table 1). Three studies specifically explored the effects of plyometric training on upper extremity proprioception (46) and rotator cuffs (39, 43). The first study by Heiderscheit (1996) (46), which involved plyometric training using a one-handed overhead throw on a trampoline, did not show any significant change in kinesthetic scores. In contrast, the studies by Swanik et al. (2002) (39) and Shemy and Battecha (2017) (43)

Table 2. Results of the Downs and Black Checklist ^a

Article	Reporting (n = 10)	External Validity (n = 3)	Internal Validity		Power (n = 1)	Total (n = 27)
			Bias (n = 7)	Confounding (n = 6)		
Zhou et al., (2022) (38)	6	1	4	2	0	13
Swanik et al., (2002) (39)	7	2	4	3	0	16
Alikhani et al., (2019) (40)	7	1	5	3	0	16
Ozer et al., (2011) (45)	2	1	5	3	0	11
Sokhangu et al., (2022) (35)	7	3	5	3	0	18
Seo et al., (2010) (41)	7	1	4	4	0	16
Waddington et al., (2000) (42)	3	1	5	4	0	13
Heiderscheit et al., (1996) (46)	5	1	5	2	0	13
Shemy and Battecha, (2017) (43)	8	3	4	6	1	22
Ismail et al., (2010) (37)	7	1	7	3	0	18
Park and Kim, (2019) (36)	4	1	5	3	0	13
Saran et al. (2022), (44)	7	1	5	5	0	18

^a Downs and Black score ranges: Excellent (26 - 28); good (20 - 25); fair (15 - 19); and poor (≤ 14) (33).

demonstrated that plyometric training significantly improved the proprioception of the rotator cuffs.

Two other studies (36, 38) focused on lower limbs and examined the effects of plyometric training using stable and unstable supporting surfaces. One study (38) found that combining plyometric and balance training on unstable surfaces increased proprioception in both dominant and non-dominant legs. Another study (36) showed that jump plyometric training yielded significant improvements only in plantar flexion range of motion (ROM) sense, specifically in the ankle joint position sense.

A study (45) on weighted and non-weighted rope exercises reported improvements in proprioception in both groups, although there was no significant difference between the proprioception scores of the weighted rope jump group and the control group.

Additionally, a combination of plyometric and neuromuscular training studied by Sokhangu et al. (2022) (35) indicated enhanced proprioception in the intervention group compared to the control group. Other studies (37, 40-42), which investigated the effects of plyometric training through jumping and hopping exercises or using ballistic six plyometric exercises (44), also reported significant increases in proprioception in the intervention groups.

Lower limb plyometric training mainly involved exercises such as squat jumps, tuck jumps, box jumps, side-to-side hops, and cone hops (35-38, 40-42), with one

study (45) using only rope exercises. For upper limb training, a wider variety of exercises were used, including throwing (46), throwing and catching (39), and ballistic movements (43, 44).

Proprioception was assessed through various methods. Five studies (35, 37, 39, 41, 43) employed a Biodex isokinetic dynamometer device. Other studies used different tools, including the Charles Bell tool (45), the AMEDA device (42), and the LIDO Active Multi-Joint II Isokinetic system (46). Simpler methods, such as photography (39), electro-goniometer (36), inclinometer (44), and force plate (38), were also utilized.

Excluding the study by Heiderscheit et al. (1996) (46), which focused solely on throwing exercises for the upper limbs, and the study by Ismail et al. (2010) (37), which combined jump and hop exercises for the lower limbs, significant improvement in proprioception was observed in the remaining studies.

5. Discussion

The present study aimed to systematically review the effects of plyometric exercises on proprioception. Based on the 12 studies included in this review, it was observed that various types of plyometric training and different methods of proprioception assessment were considered. Overall, the results suggest that plyometric training has a generally positive effect on proprioception, regardless of the specific training protocol or assessment metric used.

In most of the studies involving lower limb plyometric exercises (35-38, 40-42), the exercises included different types of jumping and hopping, such as squat jumps, tuck jumps, box jumps, side-to-side hops, and cone hops. Only one study (45) focused on rope exercises. While all studies reported significant effects of plyometric exercises on proprioception, some did not pass the full requirements of the Downs and Black checklist for study quality. Notably, four of the eight studies on lower limb plyometric exercises (36, 38, 42, 45) did not fully describe their outcome measurements, which is critical for determining the quality of the findings. In contrast, four studies (35, 37, 40, 41) used isokinetic machines for proprioception assessment, with three reporting positive effects of plyometric training on proprioception.

Regarding study reliability, the majority of the lower limb studies had consistent intervention parameters and participant numbers. The most common duration for interventions was six weeks, and the average number of participants was 19.28, with one study involving 44 subjects. Most studies scored well in the external and internal validity sections of the checklist, although some unknown or non-reported details about randomized intervention assignments affected the internal validity scores in a few studies (38, 40-42, 45).

In terms of upper limb plyometric exercises, four studies were reviewed (39, 43, 44, 46). Three of these studies (39, 44, 46) received fair or poor quality scores based on the checklist, with only one study (43) achieving a good score. Despite variations in assessment methods and training tools, all four studies generally indicated positive effects of plyometric exercises on proprioception, involving a total of 207 subjects. Thus, it can be concluded that plyometric exercises improve proprioception for both upper and lower limbs, although there is variability in study quality.

5.1. Limitations

One of the main limitations of this review is the limited number of randomized controlled trials with control groups. Additionally, most of the reviewed studies had small sample sizes, and gender differences were not accounted for in most cases. Another limitation was the variety of plyometric exercises used across studies, which may have influenced the effect on proprioception. Furthermore, the use of different tools

for assessing proprioception could have impacted the consistency and accuracy of the results.

5.2. Conclusions

Based on the studies reviewed, it is evident that plyometric exercises likely enhance proprioception. Although there are variations in the types of plyometric training and assessment methods used, the overall findings support the potential of plyometric exercises to improve proprioception in both upper and lower limbs. Future research with larger sample sizes, more standardized assessment methods, and better study designs is recommended to further substantiate these findings.

Footnotes

Authors' Contribution: Study concept and design: R. Y.; acquisition of data: R. Y.; analysis and interpretation of data: R. Y. and T. M.; drafting of the manuscript: R. Y.; critical revision of the manuscript for important intellectual content: T. M.; statistical analysis: R. Y.; administrative, technical, and material support: T. M.; study supervision: T. M.

Conflict of Interests Statement: The authors declare no conflict of interests.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Funding/Support: The authors declare no fundings.

References

1. Han J, Waddington G, Adams R, Anson J, Liu Y. Assessing proprioception: A critical review of methods. *J Sport Health Sci.* 2016;5(1):80-90. [PubMed ID: 30356896]. [PubMed Central ID: PMC6191985]. <https://doi.org/10.1016/j.jshs.2014.10.004>.
2. Goble DJ. Proprioceptive acuity assessment via joint position matching: from basic science to general practice. *Phys Ther.* 2010;90(8):1176-84. [PubMed ID: 20522675]. <https://doi.org/10.2522/ptj.20090399>.
3. Ogard W. Proprioception in Sports Medicine and Athletic Conditioning. *Strength Condition J.* 2011;33:111-8. <https://doi.org/10.1519/SSC.0b013e31821bf3ae>.
4. Nagai T, Sell TC, House AJ, Abt JP, Lephart SM. Knee proprioception and strength and landing kinematics during a single-leg stop-jump task. *J Athl Train.* 2013;48(1):31-8. [PubMed ID: 23672323]. [PubMed Central ID: PMC3554030]. <https://doi.org/10.4085/1062-6050-48.1.14>.

5. Han J, Anson J, Waddington G, Adams R, Liu Y. The Role of Ankle Proprioception for Balance Control in relation to Sports Performance and Injury. *Biomed Res Int.* 2015;2015:842804. [PubMed ID: 26583139]. [PubMed Central ID: PMC4637080]. <https://doi.org/10.1155/2015/842804>.
6. Lee D, Aronson E. Visual proprioceptive control of standing in human infants. *Percept Psychophys.* 1974;15:529-32. <https://doi.org/10.3758/BF03199297>.
7. Davids K, Williams AM, Williams JG. *Visual perception and action in sport.* 1st ed. London: Routledge; 2005. <https://doi.org/10.4324/9780203979952>.
8. Ljubojevic A, Popović B, Bijelic S, Jovanović S. Proprioceptive training in dance sport: effects of agility skills. *Turkish J Kinesiolo.* 2020;6:109-17. <https://doi.org/10.31459/turkjin.742359>.
9. Hlaing SS, Puntumetakul R, Rhine EE, Boucaut R. Effects of core stabilization exercise and strengthening exercise on proprioception, balance, muscle thickness and pain related outcomes in patients with subacute nonspecific low back pain: a randomized controlled trial. *BMC Musculoskelet Disord.* 2021;22(1):998. [PubMed ID: 34847915]. [PubMed Central ID: PMC8630919]. <https://doi.org/10.1186/s12891-021-04858-6>.
10. Lu CC, Yao HI, Fan TY, Lin YC, Lin HT, Chou PP. Twelve Weeks of a Staged Balance and Strength Training Program Improves Muscle Strength, Proprioception, and Clinical Function in Patients with Isolated Posterior Cruciate Ligament Injuries. *Int J Environ Res Public Health.* 2021;18(23). [PubMed ID: 34886588]. [PubMed Central ID: PMC8657930]. <https://doi.org/10.3390/ijerph182312849>.
11. Hu X, Lai Z, Wang L. Effects of Taichi exercise on knee and ankle proprioception among individuals with knee osteoarthritis. *Res Sports Med.* 2020;28(2):268-78. [PubMed ID: 31524502]. <https://doi.org/10.1080/15438627.2019.1663520>.
12. Xu D, Hong Y, Li J, Chan K. Effect of Taichi exercise on proprioception of ankle and knee joints in old people. *Br J Sports Med.* 2004;38(1):50-4. [PubMed ID: 14751946]. [PubMed Central ID: PMC1724726]. <https://doi.org/10.1136/bjsm.2002.003335>.
13. Sekir U, Yildiz Y, Hazneci B, Ors F, Aydin T. Effect of isokinetic training on strength, functionality and proprioception in athletes with functional ankle instability. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(5):654-64. [PubMed ID: 16770637]. <https://doi.org/10.1007/s00167-006-0108-8>.
14. Hanci E, Sekir U, Gur H, Akova B. Eccentric Training Improves Ankle Evertor and Dorsiflexor Strength and Proprioception in Functionally Unstable Ankles. *Am J Phys Med Rehabil.* 2016;95(6):448-58. [PubMed ID: 26745222]. <https://doi.org/10.1097/phm.0000000000000421>.
15. Lim C. Multi-Sensorimotor Training Improves Proprioception and Balance in Subacute Stroke Patients: A Randomized Controlled Pilot Trial. *Front Neurol.* 2019;10:157. [PubMed ID: 30881333]. [PubMed Central ID: PMC6407432]. <https://doi.org/10.3389/fneur.2019.00157>.
16. Ghaderi M, Letafatkar A, Almonroeder TG, Keyhani S. Neuromuscular training improves knee proprioception in athletes with a history of anterior cruciate ligament reconstruction: A randomized controlled trial. *Clin Biomech (Bristol, Avon).* 2020;80:105157. [PubMed ID: 32871397]. <https://doi.org/10.1016/j.clinbiomech.2020.105157>.
17. Trans T, Aaboe J, Henriksen M, Christensen R, Bliddal H, Lund H. Effect of whole body vibration exercise on muscle strength and proprioception in females with knee osteoarthritis. *Knee.* 2009;16(4):256-61. [PubMed ID: 19147365]. <https://doi.org/10.1016/j.knee.2008.11.014>.
18. Moghadasi A, Ghasemi G, Sadeghi-Demneh E, Etemadifar M. The Effect of Total Body Resistance Exercise on Mobility, Proprioception, and Muscle Strength of the Knee in People With Multiple Sclerosis. *J Sport Rehabil.* 2020;29(2):192-9. [PubMed ID: 30676232]. <https://doi.org/10.1123/jscr.2018-0303>.
19. Daman M, Shiravani F, Hemmati L, Taghizadeh S. The effect of combined exercise therapy on knee proprioception, pain intensity and quality of life in patients with hypermobility syndrome: A randomized clinical trial. *J Bodyw Mov Ther.* 2019;23(1):202-5. [PubMed ID: 30691753]. <https://doi.org/10.1016/j.jbmt.2017.12.012>.
20. Suner-Keklik S, Numanoglu-Akbas A, Cobanoglu G, Kafa N, Guzel NA. An online pilates exercise program is effective on proprioception and core muscle endurance in a randomized controlled trial. *Ir J Med Sci.* 2022;191(5):2133-9. [PubMed ID: 34716884]. [PubMed Central ID: PMC8556804]. <https://doi.org/10.1007/s11845-021-02840-8>.
21. Holm I, Fosdahl MA, Friis A, Risberg MA, Myklebust G, Steen H. Effect of neuromuscular training on proprioception, balance, muscle strength, and lower limb function in female team handball players. *Clin J Sport Med.* 2004;14(2):88-94. [PubMed ID: 15014342]. <https://doi.org/10.1097/00042752-200403000-00006>.
22. Schmitt H, Kuni B, Sabo D. Influence of professional dance training on peak torque and proprioception at the ankle. *Clin J Sport Med.* 2005;15(5):331-9. [PubMed ID: 16162992]. <https://doi.org/10.1097/01.jsm.0000181437.41268.56>.
23. Lai Z, Zhang Y, Lee S, Wang L. Effects of strength exercise on the knee and ankle proprioception of individuals with knee osteoarthritis. *Res Sports Med.* 2018;26(2):138-46. [PubMed ID: 29366340]. <https://doi.org/10.1080/15438627.2018.1431541>.
24. Lesinski M, Prieske O, Chaabene H, Granacher U. Seasonal Effects of Strength Endurance vs. Power Training in Young Female Soccer Athletes. *J Strength Cond Res.* 2021;35(Suppl 12):S90-S96. [PubMed ID: 32149876]. <https://doi.org/10.1519/jsc.00000000000003564>.
25. Stanganelli LC, Dourado AC, Oncken P, Mançan S, da Costa SC. Adaptations on jump capacity in Brazilian volleyball players prior to the under-19 World Championship. *J Strength Cond Res.* 2008;22(3):741-9. [PubMed ID: 18438245]. <https://doi.org/10.1519/JSC.0b013e31816a5c4c>.
26. Moran J, Sandercock G, Ramirez-Campillo R, Clark CCT, Fernandes JFT, Drury B. A Meta-Analysis of Resistance Training in Female Youth: Its Effect on Muscular Strength, and Shortcomings in the Literature. *Sports Med.* 2018;48(7):1661-71. [PubMed ID: 29626334]. <https://doi.org/10.1007/s40279-018-0914-4>.
27. Ramirez-Campillo R, Sanchez-Sanchez J, Romero-Moraleda B, Yanci J, García-Hermoso A, Manuel Clemente F. Effects of plyometric jump training in female soccer player's vertical jump height: A systematic review with meta-analysis. *J Sports Sci.* 2020;38(13):1475-87. [PubMed ID: 32255389]. <https://doi.org/10.1080/02640414.2020.1745503>.
28. Markovic G, Mikulic P. Neuro-musculoskeletal and performance adaptations to lower-extremity plyometric training. *Sports Med.* 2010;40(10):859-95. [PubMed ID: 20836583]. <https://doi.org/10.2165/11318370-000000000-00000>.
29. Behm DG, Faigenbaum AD, Falk B, Klentrou P. Canadian Society for Exercise Physiology position paper: resistance training in children and adolescents. *Appl Physiol Nutr Metab.* 2008;33(3):547-61. [PubMed ID: 18461111]. <https://doi.org/10.1139/h08-020>.
30. Lloyd R, Meyers R, Oliver J. The Natural Development and Trainability of Plyometric Ability During Childhood. *Strength Condition J.* 2011;33:23-32. <https://doi.org/10.1519/SSC.0b013e3182093a27>.

31. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;**6**(7): e1000097. [PubMed ID: [19621072](#)]. [PubMed Central ID: [PMC2707599](#)]. <https://doi.org/10.1371/journal.pmed.1000097>.
32. Higgins J. *Cochrane handbook for systematic reviews of interventions. Version 5.1. Cochrane Collab.* 2011, [updated March 2011]. Available from: <https://training.cochrane.org/handbook/archive/v5.1/>.
33. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health.* 1998;**52**(6):377-84. [PubMed ID: [9764259](#)]. [PubMed Central ID: [PMC1756728](#)]. <https://doi.org/10.1136/jech.52.6.377>.
34. Silva AF, Clemente FM, Lima R, Nikolaidis PT, Rosemann T, Knechtle B. The Effect of Plyometric Training in Volleyball Players: A Systematic Review. *Int J Environ Res Public Health.* 2019;**16**(16). [PubMed ID: [31426481](#)]. [PubMed Central ID: [PMC6720263](#)]. <https://doi.org/10.3390/ijerph16162960>.
35. Sokhangu MK, Rahnama N, Etemadifar M, Rafeii M, Saberi A. Effect of Neuromuscular Exercises on Strength, Proprioceptive Receptors, and Balance in Females with Multiple Sclerosis. *Int J Prev Med.* 2021;**12**:5. [PubMed ID: [34084302](#)]. [PubMed Central ID: [PMC8106275](#)]. https://doi.org/10.4103/ijpvm.IJPVM_525_18.
36. Park C, Kim BG. Effect of Jumping Exercise on Supporting Surface on Ankle Muscle Thickness, Proprioception and Balance in Adults with Functional Ankle Instability. *J Int Acad Physic Ther Res.* 2019;**10**:1756-62. <https://doi.org/10.20540/JIAPTR.2019.10.2.1756>.
37. Ismail MM, Ibrahim MM, Youssef EF, El Shorbagy KM. Plyometric training versus resistive exercises after acute lateral ankle sprain. *Foot Ankle Int.* 2010;**31**(6):523-30. [PubMed ID: [20557819](#)]. <https://doi.org/10.3113/fai.2010.0523>.
38. Zhou L, Gong W, Wang S, Guo Z, Liu M, Chuang S, et al. Combined balance and plyometric training enhances knee function, but not proprioception of elite male badminton players: A pilot randomized controlled study. *Front Psychol.* 2022;**13**:947877. [PubMed ID: [36017428](#)]. [PubMed Central ID: [PMC9396213](#)]. <https://doi.org/10.3389/fpsyg.2022.947877>.
39. Swanik KA, Lephart SM, Swanik CB, Lephart SP, Stone DA, Fu FH. The effects of shoulder plyometric training on proprioception and selected muscle performance characteristics. *J Shoulder Elbow Surg.* 2002;**11**(6):579-86. [PubMed ID: [12469083](#)]. <https://doi.org/10.1067/mse.2002.127303>.
40. Alikhani R, Shahrjerdi S, Golpaigany M, Kazemi M. The effect of a six-week plyometric training on dynamic balance and knee proprioception in female badminton players. *J Can Chiropr Assoc.* 2019;**63**(3):144-53. [PubMed ID: [31988535](#)]. [PubMed Central ID: [PMC6973753](#)].
41. Seo BD, Shin HS, Yoon JD, Han DW. The Effect of Lower Extremity Plyometric Training on the Proprioception and Postural Stability of Collegiate Soccer Players with Postural Instability. *Korean J Sport Biomechanics.* 2010;**20**:1-12. <https://doi.org/10.5103/KJSB.2010.20.1.001>.
42. Waddington G, Seward H, Wrigley T, Lacey N, Adams R. Comparing wobble board and jump-landing training effects on knee and ankle movement discrimination. *J Sci Med Sport.* 2000;**3**(4):449-59. [PubMed ID: [11235009](#)]. [https://doi.org/10.1016/s1440-2440\(00\)80010-9](https://doi.org/10.1016/s1440-2440(00)80010-9).
43. Shemy S, Battecha K. EFFECT OF ISOKINETIC VERSUS PLYOMETRIC TRAINING ON SHOULDER PROPRICEPTION AND EXTERNAL TO INTERNAL ROTATORS STRENGTH RATIO IN SWIMMER CHILDREN. *Int J Physiother Res.* 2017;**5**:2133-43. <https://doi.org/10.16965/ijpr.2017.161>.
44. Saran M, Pawaria S, Kalra S. Kinesio taping with ballistic six plyometric training on speed, accuracy, target and joint proprioception in fast bowlers with glenohumeral instability. *Comparative Exerc Physiol.* 2022;**18**:1-8. <https://doi.org/10.3920/CEP220008>.
45. Ozer D, Duzgun I, Baltaci G, Karacan S, Colakoglu F. The effects of rope or weighted rope jump training on strength, coordination and proprioception in adolescent female volleyball players. *J Sports Med Phys Fitness.* 2011;**51**(2):211-9. [PubMed ID: [21681154](#)].
46. Heiderscheit BC, McLean KP, Davies GJ. The effects of isokinetic vs. plyometric training on the shoulder internal rotators. *J Orthop Sports Phys Ther.* 1996;**23**(2):125-33. [PubMed ID: [8808515](#)]. <https://doi.org/10.2519/jospt.1996.23.2.125>.

Table 1. Intervention Characteristics

Study	Participants		Intervention				Outcome
	n	Characteristics	Plyometric Intervention	Frequency	Duration	Control	
Zhou et al., (2022) (38)	16	Healthy elite male badminton players	Combined training: 40 minutes of plyometric training and then 20 min of balance training on an unstable support (e.g., BOSU ball, Swiss ball, and Balance pad).	1 hour, 3 sessions Per weeks	6 weeks	Plyometric training: Participants completed 40 min of plyometric training (e.g., depth jump and lateral barrier jump) and then 20 min of balance training on a stable support.	Both groups led to significant improvements in proprioception
	24	Female division I swimmers	Regular swimming training program with additional plyometric training, focused on strengthening the internal rotators of the shoulder (exercises with elastic tubing and the Pitchback System)	Three sets of 15 repetitions were performed 2 days a week	6 weeks	Regular swimming training program	Plyometric training resulted in significant improvement in proprioception kinesthesia. The plyometric group improved significantly more than the control group in 5 out of 6 proprioceptive tests (active reproduction of passive positioning)
Swanik et al., (2002) (39)	22	Healthy be-ginner female badminton players,	Plyometric training which consisted of jumping exercises (e.g., wall jumps, squat jumps, board jumps, box jumps, etc.), performed in three levels of difficulty	20 minutes with 10 minutes Warm up and cool down, three times per week	6 weeks	Continued their usual badminton training and practice,	The knee joint angle reconstruction absolute error significantly improved in the intervention group compared to controls after plyometric training
	25	Female experienced-volleyball players for at least 2 years	Group I received a technical volleyball training program with a weighted rope. Group II was given the same training program but with a normal rope.	3 sessions a week	12 weeks	Followed only a technical volleyball training program for the same duration,	Proprioception improved in both intervention groups, no significant difference was found between the weighted rope jump groups and controls.
Alikhani et al., (2019) (40)	20	Female volunteers with relapsing remitting MS	Neuromuscular exercises, and in the last 2 weeks, plyometric exercises (Dynamic lunge, Side step up, Vertical jump, Pair jump, Hopping) were added with different intensities	60 min (10 min warm up, 45 exercise, 5 min cool down), 3 sessions a week	8 weeks	They were asked to maintain normal daily activities during the 8-week intervention.	The proprioceptive error during evaluating angular reconstruction decreased significantly in the experimental group, but not in the control group.
	10	Male collegiate soccer players	Plyometric training (Squat Jump, Tuck jump, Box Jump Up, Box Jump Down, Horizontal Jump, Butt Kick)	Training Intensity was increased	6 weeks	Postural stability training	Plyometric training significantly improved proprioception and postural stability ($P < 0.05$)
Ozer et al., (2011) (45)	44	Male subjects playing in three teams in the Victorian Football League (VFL)	Normal football training plus jump-landing	10 minutes, 3 sessions	8 weeks	Normal training	Knee and Ankle discrimination improved overall pre to post test in normal training plus jump-landing group compared to normal training group.
	78	Sedentary, college-aged females	Plyometric training, (On the Plyoback System and from the center of the trampoline, throwing the weighted balls at it using a one-handed overhead throw of the dominant arm)	2 sessions per week	8 weeks	Control group	None of plyometric training group and control group exhibited a significant change in kinesthetic score
Sokhangu et al., (2022) (35)	75	Swimmer boys	Study group (1) received plyometric training for the rotator cuff muscles	2 sessions per week	8 weeks	Control group (2) had no strength training, Study group (3) received isokinetic training, - Three groups swam for four hours per week in the eight weeks through- out the study.	Plyometric group showed significant differences in all measured variables with no significant changes being observed in the control group.
	22	Athletes with grade I or II unilateral inversion ankle sprain	Plyometric Training (Side to side ankle hops, standing jump and reach, Front cone hops, Standing long jump, Lateral jump over barrier, Cone hops with 180 Degree turn, Hexagon drill, etc.)	2 sessions per week	6 weeks	Resistance Training	Both plyometric and resistive training improved isokinetic evtor and invertor peak torques and functional performance of athletes $P < 0.05$. There were no significant differences between groups concerning peak torque/body weight for invertors and evtors at both speeds measured $P > 0.05$. The functional test measures of the plyometric group were significantly higher than that of resistance training group.

Study	Participants		Intervention				Outcome
	n	Characteristics	Plyometric Intervention	Frequency	Duration	Control	
Seo et al., (2010) (41)	20	Adults with FAI	Stable supporting surface jump group	30 minutes, 3 sessions	8 weeks	Unstable supporting jump group	In comparison between the groups, a significant difference in the plantar flexion range of the joint position sense after exercise was observed.
	30	Male semi-professional fast bowlers	Ballistic six plyometric training with conventional upper extremity workouts	60 minutes, 3 sessions	Weeks	Kinesio-taping along with ballistic six plyometric training with conventional upper extremity workouts	Ballistic six plyometric training group showed significant difference for joint proprioception in comparison to control group.
Waddington et al., (2000) (42)	Participants		Intervention	Outcome			
	n	Characteristics	Plyometric Intervention	Frequency	Duration	control	
Heiderscheit et al., (1996) (46)	16	Healthy elite male badminton players	Combined training: 40 minutes of plyometric training and then 20 min of balance training on an unstable support (e.g., BOSU ball, Swiss ball, and Balance pad).	1 hour, 3 sessions per weeks	6 weeks	Plyometric training: Participants completed 40 min of plyometric training (e.g., depth jump and lateral barrier jump) and then 20 min of balance training on a stable support.	Both groups led to significant improvements in proprioception
	24	Female division I swimmers	Regular swimming training program with additional plyometric training, focused on strengthening the internal rotators of the shoulder (exercises with elastic tubing and the Pitchback System)	Three sets of 15 repetitions were performed 2 days a week	6 weeks	Regular swimming training program	Plyometric training resulted in significant improvement in proprioception kinesthesia. The plyometric group improved significantly more than the control group in 5 out of 6 proprioceptive tests (active reproduction of passive positioning)
Shemy and Battecha, (2017) (43)	22	Healthy be- ginner female badminton players	Plyometric training which consisted of jumping exercises (e.g., wall jumps, squat jumps, board jumps, box jumps, etc.), performed in three levels of difficulty	20 minutes with 10 minutes warm up and cool down, three times per week	6 weeks	Continued their usual badminton training and practice,	The knee joint angle reconstruction absolute error significantly improved in the intervention group compared to controls after plyometric training
	25	Female experienced- volleyball players for at least 2 years	Group I received a technical volleyball training program with a weighted rope. Group II was given the same training program but with a normal rope.	3 sessions a week	12 weeks	Followed only a technical volleyball training program for the same duration,	Proprioception improved in both intervention groups, no significant difference was found between the weighted rope jump groups and controls.
Ismail et al., (2010) (37)	20	Female volunteers with relapsing remitting MS	Neuromuscular exercises, and in the last 2 weeks, plyometric exercises (Dynamic lunge, Side step up, Vertical jump, Pair jump, Hopping) were added with different intensities	60 min (10 min warm up, 45 exercise, 5 min cool down), 3 sessions a week	8 weeks	They were asked to maintain normal daily activities during the 8- week intervention.	The proprioceptive error during evaluating angular reconstruction decreased significantly in the experimental group, but not in the control group.
	10	Male collegiate soccer players	Plyometric training (Squat Jump, Tuck jump, Box Jump Up, Box Jump Down, Horizontal Jump, Butt Kick)	Training intensity was increased	6 weeks	Postural stability training	Plyometric training significantly improved proprioception and postural stability($P < 0.05$)
Park and Kim, (2019) (36)	44	Male subjects playing in three teams in the Victorian Football League (VFL)	Normal football training plus jump-landing	10 minutes, 3 sessions	8 weeks	Normal training	Knee and Ankle discrimination improved overall pre to post test in normal training plus jump-landing group compared to normal training group.
	78	Sedentary, college-aged females	Plyometric training, (On the Plyoback System and fom the center of the trampoline, throwing the weighted balls at it using a one- handed overhead throw of the dominant arm)	2 sessions per week	8 weeks	Control group,	None of plyometric training group and control group exhibited a significant change in kinesthetic score
Saran et al., (2022) (44)	75	Swimmer boys,	Study group (1) received plyometric training for the rotator cuff muscles	2 sessions Per week	8 weeks	Control group (2) had no strength training, Study group (3) received isokinetic training, -Three groups swam for four hours per week in the eight weeks through- out the study.	Plyometric group showed significant differences in all measured variables with no significant changes being observed in the control group.