



# Isometric Exercise for Blood Pressure and Endothelial Function in Metabolic Syndrome: A Review

Mohammad Bagher Afshar Naseri <sup>1, \*</sup>

<sup>1</sup> Physical Education and Sports Science Department, North Tehran Branch, Islamic Azad University, Tehran, Iran

\* **Corresponding Author:** Physical Education and Sports Science Department, North Tehran Branch, Islamic Azad University, Tehran, Iran. Email: mafsharnaseri@gmail.com

**Received:** 20 August, 2024; **Revised:** 20 October, 2024; **Accepted:** 15 November, 2024

## Abstract

**Context:** Metabolic syndrome (MetS) is characterized by a cluster of metabolic disorders, including hypertension, insulin resistance, and dyslipidemia, which collectively heighten the risk of cardiovascular diseases (CVDs). The global prevalence of MetS is steadily increasing, paralleling the rise in obesity and sedentary lifestyles. Pharmacological treatments for MetS often face challenges such as inadequate blood pressure (BP) control and limited improvements in cardiovascular outcomes. Consequently, there is a growing emphasis on non-pharmacological interventions, particularly exercise. Among various exercise modalities, isometric resistance training (IRT), which involves static muscle contractions without joint movement, has emerged as a promising strategy for effectively lowering BP and enhancing endothelial function, both of which are critical for cardiovascular health.

**Evidence Acquisition:** This comprehensive literature review synthesizes existing research on the effects of isometric exercise on BP regulation and endothelial function in individuals diagnosed with MetS. The review examines studies sourced from databases such as PubMed, Scopus, and Web of Science, focusing on randomized controlled trials, meta-analyses, and observational studies published in peer-reviewed journals. Key discussion areas include the underlying physiological mechanisms of IRT, comparisons with other exercise modalities, clinical implications, and recommendations for future research. Articles were selected based on qualitative and descriptive reviews.

**Results:** Evidence indicates that isometric resistance training effectively reduces both systolic and diastolic BP, achieving results comparable to other exercise forms like aerobic and dynamic resistance training. Furthermore, IRT has significantly improved endothelial function, particularly among populations with MetS. These physiological benefits are attributed to mechanisms such as enhanced nitric oxide bioavailability, reduced oxidative stress, and improved autonomic regulation.

**Conclusions:** The review supports the integration of isometric exercise into hypertension management protocols, especially for individuals with MetS. Isometric resistance training offers a viable non-pharmacological alternative or adjunct to traditional treatments, with the added advantages of accessibility and adaptability for diverse populations. However, further research is necessary to optimize exercise protocols, determine long-term outcomes, and fully elucidate the molecular mechanisms underlying the cardiovascular benefits of IRT.

**Keywords:** Isometric Exercise, Blood Pressure, Endothelium, Metabolic Syndrome, Cardiovascular Health, Public Health

## 1. Background

Hypertension is a primary component of metabolic syndrome (MetS) and one of the leading causes of cardiovascular morbidity and mortality worldwide (1, 2). Metabolic syndrome, on the other hand, comprises a cluster of metabolic disorders, including abdominal obesity, dyslipidemia, insulin resistance, and hypertension, posing a significant threat to public health. The prevalence of MetS has been steadily

increasing, paralleling the global rise in obesity and sedentary lifestyles (3-5). Despite advances in pharmacological treatments, there are instances where these approaches fail to fully control blood pressure (BP) or improve cardiovascular outcomes. As a result, there is an increasing emphasis on non-pharmacological interventions, particularly exercise (6, 7).

The coexistence of hypertension and MetS can present complex challenges, especially for at-risk

populations diagnosed with one or both conditions. Addressing these challenges through tailored exercise regimens may significantly benefit both conditions (4).

Isometric resistance training (IRT), which involves static muscular contractions without joint movement, has emerged as a powerful non-pharmacological strategy for lowering blood pressure and enhancing endothelial function (8, 9). Unlike dynamic exercises, IRT involves applying constant pressure to muscles rather than continuous movement, eliciting distinct cardiovascular responses (8, 10).

This review explores the impact of isometric exercise on BP and endothelial function in individuals with MetS. It provides a comprehensive overview of previous studies, highlighting key findings and offering valuable insights into potential therapeutic applications.

Hypertension is a prevalent and serious cardiovascular condition, defined as a persistent elevation of systolic blood pressure above 140 mm Hg and/or diastolic blood pressure above 90 mm Hg (10). It is a major risk factor for cardiovascular diseases (CVDs), including stroke, myocardial infarction, and heart failure (11). The pathophysiology of hypertension is complex, involving a combination of genetic, environmental, and behavioral factors. In the context of metabolic syndrome, hypertension is often accompanied by insulin resistance, dyslipidemia, and central obesity, all of which contribute to endothelial dysfunction and increased cardiovascular risk (12, 13). The rising prevalence of metabolic syndrome is largely attributed to lifestyle factors, particularly sedentary behavior and unhealthy dietary habits (14). Addressing these risk factors through lifestyle modifications, including increased physical activity, is crucial for mitigating the adverse health effects associated with metabolic syndrome (14, 15).

Non-pharmacological interventions play a critical role in the management of hypertension, particularly in individuals with MetS, where pharmacological treatments alone may not be sufficient (15-17). Lifestyle modifications, including dietary changes, weight loss, and increased physical activity, are essential components of hypertension management (18-20). Exercise, in particular, has been shown to have a direct impact on BP regulation and cardiovascular health (20-22). Various forms of exercise, including aerobic, resistance, and isometric training, have been studied for their effects on BP (15, 21, 23). Aerobic exercises such as walking, running, and cycling are widely recommended for lowering BP and enhancing cardiovascular health (24, 25). However, some recent studies have highlighted the potential benefits of both resistance and isometric

exercises, especially among individuals with hypertension and MetS (4, 10, 15, 26). Notably, isometric exercise has been shown to reduce both systolic and diastolic BP, with effects comparable to those achieved through aerobic exercise (10). Isometric exercise involves the contraction of muscles without any movement of the muscle filaments at the intracellular level or movement at the joint. Common examples include planks, wall sits, and handgrip exercises. Isometric exercise improves cardiovascular health by increasing nitric oxide bioavailability, enhancing autonomic balance, boosting parasympathetic activity, reducing sympathetic outflow, and decreasing vascular resistance through sustained intramuscular pressure (10, 12, 27, 28). Unlike dynamic exercises, which involve a full range of motion, isometric exercises are performed by maintaining a static position, which places continuous tension on the muscles. Isometric training contributes to reduced oxidative stress and improved vascular health, further supporting cardiovascular stability and reducing the risk of cardiovascular events (6, 8, 25, 29). This unique aspect of isometric exercise contributes to its distinct physiological effects, particularly in relation to BP and vascular function. The sustained increase in intramuscular pressure leads to a reduction in blood flow to the active muscles, resulting in a localized increase in vascular resistance. This triggers a compensatory increase in cardiac output and a subsequent rise in systemic BP. However, with regular isometric training, these acute responses are attenuated, leading to long-term reductions in resting BP (6, 10).

## 2. Methods

### 2.1. Evidence Acquisition

This review provides a complete analysis of the current evidence regarding the impact of isometric exercise on the regulation of blood pressure and endothelial function in patients who have been diagnosed with metabolic syndrome. The review provides an in-depth examination of papers obtained from several sources such as PubMed, Scopus, and Web of Science. The primary focus is on randomized controlled trials, meta-analyses, and observational studies that have been published in peer-reviewed journals. The search approach used the use of specific keywords like "isometric exercise," "blood pressure," "endothelial function," "metabolic syndrome," and "cardiovascular health."

### 2.2. Study Selection

Studies were included if they met the following criteria: (1) Involved adult participants diagnosed with metabolic syndrome or related cardiovascular risk factors; or (2) investigated the effects of isometric exercise on BP and/or endothelial function; and/or (3) provided quantitative measures of outcomes. Studies that focused on other exercise modalities without comparison to isometric exercise, or those that did not report on BP or endothelial function, were excluded.

### 2.3. Data Extraction and Synthesis

Data on study design, participant characteristics, exercise protocols, duration, and outcomes were extracted and synthesized to provide a comprehensive overview of the impact of isometric exercise on BP and endothelial function. The review also compares isometric resistance training with other exercise modalities to evaluate its relative effectiveness and discusses potential mechanisms underlying the observed effects.

### 2.4. Limitations

It is acknowledged that the heterogeneity of study designs, variations in exercise protocols, and participant characteristics may influence the outcomes. Additionally, the review is limited to studies published in English, which may introduce language bias. Based on existing literature, this review aims to synthesize and represent more application-based studies for a more straightforward investigation. Another consideration is the potential for publication bias, as studies showing positive results are more likely to be published than those with null findings. Moreover, the review was restricted to English-language publications, which may exclude relevant studies in other languages and contribute to language bias.

## 3. Results

The articles included in this review consist of descriptive reviews (1, 3-5, 8, 10, 11, 13-17, 20, 22-24, 26, 30-37) and practical studies (12, 21, 25, 27, 28, 38) (Table 1). These studies focus on MetS, hypertension, and practical non-pharmacological interventions. Numerous studies have investigated the effects of isometric exercise on blood pressure, consistently showing significant reductions in both systolic and diastolic BP (6). A meta-analysis of randomized controlled trials by Carlson et al. demonstrated that isometric handgrip exercise led to an average reduction of 6.77 mm Hg in systolic BP and 3.96 mm Hg in diastolic BP among hypertensive individuals (6). These reductions are significant and suggest that

isometric exercise could serve as a complementary therapy for hypertension management, with effects comparable to those achieved with antihypertensive medications (10).

Further research by Millar et al. examined the impact of different intensities of isometric exercise on blood pressure reduction, finding that moderate-intensity exercises, conducted at 30 - 40% of maximal voluntary contraction, were particularly effective (8). The study showed significant decreases in both systolic and diastolic blood pressure, positioning isometric exercise as a viable non-pharmacological approach for managing hypertension. The flexibility of these exercises allows them to be adapted to individual fitness levels, making them accessible and effective for a wide range of people. This adaptability highlights the potential for isometric training to be widely used in both clinical and personal health settings to improve cardiovascular outcomes (8).

Endothelial dysfunction is a critical factor in the pathogenesis of hypertension and metabolic syndrome, contributing to increased vascular stiffness and impaired vasodilation. Evidence suggests that isometric exercise, particularly isometric handgrip training, can significantly improve endothelial function by enhancing nitric oxide production and reducing oxidative stress (11, 28). For instance, a study by Seals et al. demonstrated that 8 weeks of isometric handgrip training led to substantial improvements in endothelial function in older adults with prehypertension (11). These improvements were reflected in increased flow-mediated dilation and reduced levels of endothelin-1, a potent vasoconstrictor (28). The enhancements in endothelial function are especially beneficial for individuals with metabolic syndrome, where endothelial dysfunction is often worsened by insulin resistance and chronic inflammation. By improving endothelial health, isometric exercise may play a crucial role in mitigating the heightened cardiovascular risks associated with metabolic syndrome, as supported by findings that link these exercise-induced vascular benefits to a reduction in blood pressure and improved autonomic regulation (11, 15). While aerobic exercise has long been considered the gold standard for cardiovascular health, with well-documented benefits for BP regulation and endothelial function, isometric exercise offers unique advantages, particularly for individuals who may have difficulty performing dynamic exercises due to mobility issues or other health concerns (10, 27, 28, 37, 39).

Physiological benefits suggest that isometric exercise can be effectively integrated into standard hypertension management protocols. As a non-pharmacological

Table 1. Practical Studies Extracted

Author(s), Year	Study Design	Population	Intervention	Comparison/Control	Duration	Outcomes Measured	Main Findings	Quality/Risk of Bias
1. Cornelissen and Smart, 2013 (12)	Systematic review and meta-analysis	5 223 adults (1 401 exercise;1822 control) from 93 RCTs	Endurance, dynamic resistance, combined endurance & resistance, isometric resistance training	Control groups (no exercise)	≥ 4 weeks	SBP and DBP	Isometric resistance training led to the largest SBP reduction (-10.9 mmHg) compared to endurance (-3.5 mmHg) and dynamic resistance (-1.8 mmHg). DBP reductions were significant across all modalities, with isometric training showing substantial decreases (-6.2 mmHg).	A high number of studies included robust meta-analytic methods. Potential heterogeneity due to varying exercise protocols.
2. Cornelissen et al., 2010 (38)	Randomized crossover study	Sedentary adults ≥ 55 years old	Endurance training at low (33% HR reserve) and high intensity (66% HR reserve)	Sedentary period (no exercise)	Ten weeks per training intensity	SBP, HR, HRV	Both low and high-intensity training significantly reduced SBP similarly. High-intensity training had more pronounced effects on HR but no significant impact on HRV.	Randomized crossover design reduces inter-subject variability. Limited sample size may affect generalizability.
3. Correia et al., 2023 (27)	Systematic review and meta-analysis of RCTs	253 hypertensive adults from 14 studies	Strength Training (various protocols)	Control groups (no exercise)	≥ 8 weeks	SBP, DBP	Significant reductions in SBP and DBP, especially with moderate to vigorous intensity (> 60% 1RM), ≥ 2 times/week, and ≥ 8 weeks duration.	Cochrane methodology enhances reliability. A limited number of studies on isometric training specifically.
4. Decaux et al., 2022 (28)	Randomized sham-controlled study	30 physically inactive adults (15 males, 15 females)	Isometric exercise training (IET)-Wall squats at 95% peak HR	Sham group (wall squats < 75% peak HR), No-intervention control	Four weeks	Resting and continuous BP, cardiac autonomic modulation HRV	IET significantly decreased SBP, DBP, and mean BP compared to sham and control. Improved HRV indicators (↓ LFnu, ↑ HFnu) in the IET group.	Sham-controlled design strengthens causal inferences. A small sample size may limit statistical power.
5. Park et al., 2020 (25)	Pilot study (randomized controlled trial)	20 obese older men (10 EXP, 10 CON)	Combined exercise: Elastic-band resistance training + aerobic (walking/running, cycling) at 60 -70% HR max, 3 days/week	Control group (no exercise)	12 weeks	Body composition, cardiometabolic risk factors, bp, arterial stiffness, physical functions	Significant decreases in body weight, BMI, % body fat, BP, arterial stiffness, LDL-C, and epinephrine. Increases in VO <sub>2</sub> peak and grip strength in the EXP group.	A pilot study with a small sample size; results are promising but need replication in larger trials.
6. Pedralli et al., 2020 (21)	Randomized clinical trial	42 individuals with prehypertension or hypertension (average age 54 ± 11 years)	Aerobic training, RT, combined training	Each exercise modality compared to others	8 weeks	Endothelial Function FMD, Ambulatory BP SBP, DBP	All three modalities reduced BP and improved FMD similarly. CT showed the greatest improvement in FMD (+6.8%).	Randomized allocation enhances internal validity. Variability in BP response suggests individualized effects.

Abbreviations: SBP, systolic blood pressure; DBP, diastolic blood pressure; HR, heart rate; HRV, heart rate variability; 1RM, one-repetition maximum; FMD, flow-mediated dilation; LFnu, low-frequency normalized units; HFnu, high-frequency normalized units; EXP, exercise group; CON, control group.

intervention, it provides a viable alternative or complement to traditional pharmacological treatments, particularly for patients who do not respond adequately to medication alone or who wish to minimize drug dependency (16, 33). Furthermore, its versatility and adaptability make it accessible to a broad range of

individuals, including those with mobility limitations or other health concerns that may restrict their ability to engage in more dynamic forms of exercise. Integrating isometric exercise into clinical practice could offer a comprehensive approach to blood pressure management, potentially leading to improved



**Table 2.** Sample Recommendations for Implementing Isometric Exercise in Hypertension Management

Exercise Type	Intensity (%MVC <sup>a</sup> )	Duration per Session (min)	Frequency (per Week)	Expected Outcome	Additional Notes
Handgrip	30 - 40	2 - 3	3 - 5	Moderate BP reduction	Ideal for beginners; portable and easy to perform
Wall sits	30 - 50	1 - 2	3 - 5	Significant BP reduction	Effective for lower body strength and endurance
Planks	35 - 45	1 - 2	3 - 4	Improved endothelial function	Engages core muscles, beneficial for overall stability
Leg press	40 - 50	2 - 3	3	Enhanced vascular health	Suitable for those with access to gym equipment
Squat hold	30 - 45	1 - 2	3 - 4	Moderate BP reduction, improved muscle endurance	Requires proper form to avoid knee strain
Arm curl hold	30 - 40	1 - 2	3 - 4	Improved upper body strength, moderate BP reduction	Can be performed with resistance bands or light weights

Abbreviation: BP, systolic blood pressure.

<sup>a</sup> Percentage of maximum voluntary contraction.

cardiovascular outcomes and a reduction in the overall burden of hypertension-related complications. Therefore, further research and clinical trials are needed to optimize isometric exercise protocols and fully incorporate them into hypertension treatment strategies (32-34).

To maximize the benefits of isometric exercise, it is essential to establish clear guidelines for its implementation. Current evidence-based recommendations suggest that isometric exercises should be performed 3 - 5 times per week (Table 2), with each session lasting 10 - 15 minutes (10, 19, 32, 38). Exercises should be performed at a moderate intensity, equivalent to 30 - 40% of maximal voluntary contraction, to achieve optimal blood pressure reductions (Table 2). Additionally, healthcare providers should take into account individual patient characteristics—such as fitness level, comorbidities, and treatment goals—when prescribing isometric exercise programs (12, 19, 38). Table 2 summarizes guidelines for individuals and professionals to use when developing a training plan based on their specific needs. It is important to note that this sample plan is likely to be more generalized, and advanced clinicians should approach its practical application with caution.

The findings of this review support the efficacy of isometric exercise in reducing BP and improving endothelial function in individuals with MetS (11, 19, 21, 25). The observed reductions in BP are comparable to those achieved with antihypertensive medications, underscoring the potential of isometric exercise as a non-pharmacological intervention for hypertension

management. The improvements in endothelial function further highlight the cardiovascular benefits of isometric exercise, particularly in populations at high risk for CVDs (34, 40). While aerobic and dynamic resistance exercises are widely recognized for their cardiovascular benefits, isometric exercise offers unique advantages, particularly in improving vascular function. The findings suggest that isometric exercise may be especially beneficial for individuals who are unable or unwilling to engage in more dynamic forms of exercise (12, 19, 38). However, further research is needed to determine the optimal combination of exercise modalities for maximizing cardiovascular health outcomes.

The clinical significance of these findings lies in the potential for isometric exercise to be integrated into routine clinical practice for managing hypertension and metabolic syndrome. Healthcare providers should consider prescribing isometric exercise as part of a comprehensive lifestyle intervention program for patients with elevated BP and cardiovascular risk factors (19, 30, 31, 33, 34). Given its accessibility and ease of implementation, isometric exercise could be a valuable tool for improving public health outcomes, particularly in underserved populations with limited access to traditional exercise facilities (20, 30).

#### 4. Discussion

This literature review provides a synthetic analysis of existing research on the effects of isometric exercise on BP and endothelial function in individuals with

metabolic syndrome. The findings demonstrate that isometric exercise significantly reduces both systolic and diastolic BP, with its impact being comparable to, and in some cases exceeding, those achieved through traditional aerobic or dynamic resistance training. This conclusion aligns with the findings of Edwards et al. (10). Additionally, isometric exercise enhances endothelial function, a critical factor in cardiovascular health, by improving nitric oxide bioavailability and simultaneously reducing oxidative stress.

The reductions in BP observed across multiple studies are clinically significant, with consistent results reported within the same category. For instance, Carlson et al. reported average decreases of 6.77 mm Hg in systolic BP and 3.96 mm Hg in diastolic BP following isometric handgrip training (6). Such reductions are noteworthy because even modest decreases in BP can substantially lower the risk of cardiovascular events, such as stroke and myocardial infarction. This is a crucial outcome for individuals at heightened risk, particularly those with underlying conditions like MetS. The findings of Millar et al. further reinforce the efficacy of moderate-intensity isometric exercise (30 - 40% of maximal voluntary contraction) in achieving optimal BP reductions (8).

The improvement in endothelial function triggered by isometric exercise is particularly relevant for individuals with metabolic syndrome, who often exhibit endothelial dysfunction due to factors such as insulin resistance and chronic inflammation. Enhanced endothelial function, as evidenced by increased flow-mediated dilation and decreased levels of endothelin-1, suggests that isometric exercise may help reverse some of the vascular impairments characteristics of MetS (11). This could potentially lead to reduced arterial stiffness and improved overall vascular health, thereby increasing longevity and overall well-being.

When comparing isometric exercise to other modalities, such as aerobic exercise and resistance training, several unique advantages emerge that underline the beneficial aspects of isometric training. While aerobic exercise remains the gold standard for cardiovascular health, improving both blood circulation and cardiac output, isometric exercise offers a viable alternative in certain cases. As argued by Edwards, isometric exercise may even surpass the potential benefits of aerobic exercise, particularly for individuals who have limitations that prevent them from engaging in dynamic activities—limitations often caused by aging or a sedentary lifestyle, which are common contributors to metabolic syndrome. For example, individuals with joint issues, mobility

restrictions, or certain chronic conditions may find isometric exercises more accessible, offering various benefits, such as increased strength, greater muscle activity, and, as a result of training adaptation, a reduction in the root causes of MetS and CVDs. Edwards et al. highlighted that isometric exercise was more effective than aerobic exercise in improving vascular resistance and autonomic function measures (10).

The autonomic benefits of isometric exercise, such as enhanced parasympathetic activity and reduced sympathetic outflow, play a crucial role in blood pressure regulation (6, 26, 28). By modulating the autonomic nervous system, isometric exercise can lead to sustained blood pressure reductions and improved heart rate variability, both of which are important markers of cardiovascular health. Additionally, the muscle activation resulting from isometric exercise offers underlying benefits that may help decrease or even prevent metabolic syndrome and related health issues.

From a clinical and practical perspective, incorporating isometric exercise into hypertension management protocols offers a promising non-pharmacological strategy. Given the challenges associated with medication adherence and the potential side effects of antihypertensive drugs, isometric exercise provides a low-cost, low-risk intervention that patients can easily implement. The simplicity of isometric exercises, such as handgrip squeezes or wall sits, allows for flexibility in various settings, including at home or in clinical environments. However, exercise guideline developers must acknowledge several limitations. The heterogeneity of study designs, including variations in exercise protocols, duration, and intensity, makes it challenging to establish standardized guidelines, as each individual's circumstances may vary. Additionally, most studies had relatively short follow-up periods. As a result, the long-term sustainability of blood pressure reductions and improvements in endothelial function remains uncertain, which necessitates further cohort-based studies in diverse populations to track the ongoing adaptation to isometric training. Furthermore, there is a lack of large-scale, randomized controlled trials explicitly focusing on populations with metabolic syndrome, which limits the generalizability of the findings (36).

Future research should aim to address these limitations by conducting long-term, randomized controlled trials with larger and more diverse populations. Investigations into the optimal intensity, duration, and frequency of isometric exercise are necessary to develop evidence-based guidelines.

Additionally, exploring the molecular mechanisms underlying the benefits of isometric exercise could provide deeper insights into how these exercises influence cardiovascular health.

#### 4.1. Conclusions

This article presents a literature review of existing studies, which demonstrate that isometric training can effectively reduce and manage hypertension. Isometric exercise emerges as a potent, non-pharmacological intervention for lowering blood pressure and enhancing endothelial function in individuals with metabolic syndrome. Its ease of implementation and adaptability make it a practical addition to hypertension management strategies, particularly for those unable to participate in traditional exercise programs. By integrating isometric exercise into comprehensive lifestyle interventions, healthcare providers can offer patients a holistic approach to reducing cardiovascular risk and improving overall health outcomes.

#### Footnotes

**Authors' Contribution:** Study concept and design: M. B. and A. N.; Drafting of the manuscript: M. B. and A. N.

**Conflict of Interests Statement:** There is no conflict of interest to report.

**Data Reproducibility:** The dataset presented in the study is available on request from the corresponding author during submission or after publication.

**Funding/Support:** The review study did not receive any funding.

#### References

- Neutel JM, Smith DH. Hypertension control: multifactorial contributions. *Am J Hypertens*. 1999;**12**(12 Pt 1-2):164S-9S. [PubMed ID: 10619597]. [https://doi.org/10.1016/S0895-7061\(99\)00221-6](https://doi.org/10.1016/S0895-7061(99)00221-6).
- Bansal A, Hiwale K. Updates in the Management of Coronary Artery Disease: A Review Article. *Cureus*. 2023. <https://doi.org/10.7759/cureus.50644>.
- Huang PL. A comprehensive definition for metabolic syndrome. *Dis Model Mech*. 2009;**2**(5-6):231-7. [PubMed ID: 19407331]. [PubMed Central ID: PMC2675814]. <https://doi.org/10.1242/dmm.001180>.
- Hayden MR. Overview and New Insights into the Metabolic Syndrome: Risk Factors and Emerging Variables in the Development of Type 2 Diabetes and Cerebrocardiovascular Disease. *Med (Kaunas)*. 2023;**59**(3). [PubMed ID: 36984562]. [PubMed Central ID: PMC10059871]. <https://doi.org/10.3390/medicina59030561>.
- Myers J, Kokkinos P, Nyelin E. Physical Activity, Cardiorespiratory Fitness, and the Metabolic Syndrome. *Nutrients*. 2019;**11**(7). [PubMed ID: 31331009]. [PubMed Central ID: PMC6683051]. <https://doi.org/10.3390/nu11071652>.
- Carlson DJ, Dieberg G, Hess NC, Millar PJ, Smart NA. Isometric exercise training for blood pressure management: a systematic review and meta-analysis. *Mayo Clin Proc*. 2014;**89**(3):327-34. [PubMed ID: 24582191]. <https://doi.org/10.1016/j.mayocp.2013.10.030>.
- Collins R, Peto R, MacMahon S, Hebert P, Fiebach NH, Eberlein KA, et al. Blood pressure, stroke, and coronary heart disease. Part 2, Short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet*. 1990;**335**(8693):827-38. [PubMed ID: 1969567]. [https://doi.org/10.1016/0140-6736\(90\)90944-z](https://doi.org/10.1016/0140-6736(90)90944-z).
- Millar PJ, McGowan CL, Cornelissen VA, Araujo CG, Swaine IL. Evidence for the role of isometric exercise training in reducing blood pressure: potential mechanisms and future directions. *Sports Med*. 2014;**44**(3):345-56. [PubMed ID: 24174307]. <https://doi.org/10.1007/s40279-013-0118-x>.
- Dasgupta C, Zhang L. Angiotensin II receptors and drug discovery in cardiovascular disease. *Drug Discov Today*. 2011;**16**(1-2):22-34. [PubMed ID: 21147255]. [PubMed Central ID: PMC3022115]. <https://doi.org/10.1016/j.drudis.2010.11.016>.
- Edwards JJ, Coleman DA, Ritti-Dias RM, Farah BQ, Stensel DJ, Lucas SJE, et al. Isometric Exercise Training and Arterial Hypertension: An Updated Review. *Sports Med*. 2024;**54**(6):1459-97. [PubMed ID: 38762832]. [PubMed Central ID: PMC11239608]. <https://doi.org/10.1007/s40279-024-02036-x>.
- Seals DR, Jablonski KL, Donato AJ. Aging and vascular endothelial function in humans. *Clin Sci (Lond)*. 2011;**120**(9):357-75. [PubMed ID: 21244363]. [PubMed Central ID: PMC3482987]. <https://doi.org/10.1042/CS20100476>.
- Cornelissen VA, Smart NA. Exercise training for blood pressure: a systematic review and meta-analysis. *J Am Heart Assoc*. 2013;**2**(1). e004473. [PubMed ID: 23525435]. [PubMed Central ID: PMC3603230]. <https://doi.org/10.1161/JAHA.112.004473>.
- Nicoll R, Henein MY. Hypertension and lifestyle modification: how useful are the guidelines? *Br J Gen Pract*. 2010;**60**(581):879-80. [PubMed ID: 21144196]. [PubMed Central ID: PMC2991739]. <https://doi.org/10.3399/bjgp10X544014>.
- Krishnamoorthy Y, Rajaa S, Murali S, Sahoo J, Kar SS. Association Between Anthropometric Risk Factors and Metabolic Syndrome Among Adults in India: A Systematic Review and Meta-Analysis of Observational Studies. *Prev Chronic Dis*. 2022;**19**. E24. [PubMed ID: 35512304]. [PubMed Central ID: PMC9109643]. <https://doi.org/10.5888/pcd19.210231>.
- Smart NA, Way D, Carlson D, Millar P, McGowan C, Swaine I, et al. Effects of isometric resistance training on resting blood pressure: individual participant data meta-analysis. *J Hypertens*. 2019;**37**(10):1927-38. [PubMed ID: 30889048]. [PubMed Central ID: PMC6727950]. <https://doi.org/10.1097/HJH.0000000000002105>.
- Kawai T, Forrester SJ, O'Brien S, Baggett A, Rizzo V, Eguchi S. AT1 receptor signaling pathways in the cardiovascular system. *Pharmacol Res*. 2017;**125**(Pt A):4-13. [PubMed ID: 28527699]. [PubMed Central ID: PMC5607088]. <https://doi.org/10.1016/j.phrs.2017.05.008>.
- Oori MJ, Mohammadi F, Norozi K, Fallahi-Khoshknab M, Ebadi A, Gheshlagh RG. Prevalence of HTN in Iran: Meta-analysis of Published Studies in 2004-2018. *Curr Hypertens Rev*. 2019;**15**(2):113-22. [PubMed ID: 30657043]. [PubMed Central ID: PMC6635676]. <https://doi.org/10.2174/1573402115666190118142818>.
- Di Raimondo D, Buscemi S, Musiari G, Rizzo G, Pirera E, Corleo D, et al. Ketogenic Diet, Physical Activity, and Hypertension-A Narrative Review. *Nutrients*. 2021;**13**(8). [PubMed ID: 34444726]. [PubMed Central ID: PMC8398985]. <https://doi.org/10.3390/nu13082567>.
- Gollie JM, Cohen SD, Patel SS. Physical Activity and Exercise for Cardiorespiratory Health and Fitness in Chronic Kidney Disease. *Rev*

- Cardiovasc Med.* 2022;**23**(8):273. [PubMed ID: 36945353]. [PubMed Central ID: PMC10026709]. <https://doi.org/10.31083/j.rcm2308273>.
20. Williams MA, Haskell WL, Ades PA, Amsterdam EA, Bittner V, Franklin BA, et al. Resistance exercise in individuals with and without cardiovascular disease: 2007 update: a scientific statement from the American Heart Association Council on Clinical Cardiology and Council on Nutrition, Physical Activity, and Metabolism. *Circulation.* 2007;**116**(5):572-84. [PubMed ID: 17638929]. <https://doi.org/10.1161/CIRCULATIONAHA.107.185214>.
  21. Pedralli ML, Marschner RA, Kollet DP, Neto SG, Eibel B, Tanaka H, et al. Different exercise training modalities produce similar endothelial function improvements in individuals with prehypertension or hypertension: a randomized clinical trial Exercise, endothelium and blood pressure. *Sci Rep.* 2020;**10**(1):7628. [PubMed ID: 32376984]. [PubMed Central ID: PMC7203179]. <https://doi.org/10.1038/s41598-020-64365-x>.
  22. Kariuki JK, Imes CC, Engberg SJ, Scott PW, Klem ML, Cortes YI. Impact of lifestyle-based interventions on absolute cardiovascular disease risk: a systematic review and meta-analysis. *JBI Evid Synth.* 2024;**22**(1):4-65. [PubMed ID: 37551161]. [PubMed Central ID: PMC10841079]. <https://doi.org/10.1124/JBIES-22-00356>.
  23. Ozemek C, Tiwari S, Sabbahi A, Carbone S, Lavie CJ. Impact of therapeutic lifestyle changes in resistant hypertension. *Prog Cardiovasc Dis.* 2020;**63**(1):4-9. [PubMed ID: 31756356]. [PubMed Central ID: PMC7257910]. <https://doi.org/10.1016/j.pcad.2019.11.012>.
  24. Ferretti G, Fagoni N, Taboni A, Vinetti G, di Prampero PE. A century of exercise physiology: key concepts on coupling respiratory oxygen flow to muscle energy demand during exercise. *Eur J Appl Physiol.* 2022;**122**(6):1317-65. [PubMed ID: 35217911]. [PubMed Central ID: PMC9132876]. <https://doi.org/10.1007/s00421-022-04901-x>.
  25. Park W, Jung WS, Hong K, Kim YY, Kim SW, Park HY. Effects of Moderate Combined Resistance- and Aerobic-Exercise for 12 Weeks on Body Composition, Cardiometabolic Risk Factors, Blood Pressure, Arterial Stiffness, and Physical Functions, among Obese Older Men: A Pilot Study. *Int J Environ Res Public Health.* 2020;**17**(19). [PubMed ID: 33022918]. [PubMed Central ID: PMC7579509]. <https://doi.org/10.3390/ijerph17197233>.
  26. Rickson JJ, Maris SA, Headley SAE. Isometric Exercise Training: A Review of Hypothesized Mechanisms and Protocol Application in Persons with Hypertension. *Int J Exercise Sci.* 2021;**14**(2):1261-76. <https://doi.org/10.70252/kuhm5244>.
  27. Correia RR, Veras ASC, Tebar WR, Rufino JC, Batista VRG, Teixeira GR. Strength training for arterial hypertension treatment: a systematic review and meta-analysis of randomized clinical trials. *Sci Rep.* 2023;**13**(1):201. [PubMed ID: 36604479]. [PubMed Central ID: PMC9814600]. <https://doi.org/10.1038/s41598-022-26583-3>.
  28. Decaux A, Edwards JJ, Swift HT, Hurst P, Hopkins J, Wiles JD, et al. Blood pressure and cardiac autonomic adaptations to isometric exercise training: A randomized sham-controlled study. *Physiol Rep.* 2022;**10**(2). e15112. [PubMed ID: 35083878]. [PubMed Central ID: PMC8792514]. <https://doi.org/10.14814/phy2.15112>.
  29. Coronado F, Melvin SC, Bell RA, Zhao G. Global Responses to Prevent, Manage, and Control Cardiovascular Diseases. *Prev Chronic Dis.* 2022;**19**. E84. [PubMed ID: 36480801]. [PubMed Central ID: PMC9746707]. <https://doi.org/10.5888/pcd19.220347>.
  30. Higgins S, Pomeroy A, Bates LC, Paterson C, Barone Gibbs B, Pontzer H, et al. Sedentary behavior and cardiovascular disease risk: An evolutionary perspective. *Front Physiol.* 2022;**13**:962791. [PubMed ID: 35965885]. [PubMed Central ID: PMC9363656]. <https://doi.org/10.3389/fphys.2022.962791>.
  31. Kunutsor SK, Laukkanen JA. Physical activity, exercise and adverse cardiovascular outcomes in individuals with pre-existing cardiovascular disease: a narrative review. *Expert Rev Cardiovasc Ther.* 2024;**22**(1-3):91-101. [PubMed ID: 38488568]. [PubMed Central ID: PMC1057847]. <https://doi.org/10.1080/14779072.2024.2328644>.
  32. Lavie CJ, Ozemek C, Carbone S, Katzmarzyk PT, Blair SN. Sedentary Behavior, Exercise, and Cardiovascular Health. *Circ Res.* 2019;**124**(5):799-815. [PubMed ID: 30817262]. <https://doi.org/10.1161/CIRCRESAHA.118.312669>.
  33. Mancia G, Kreutz R, Brunstrom M, Burnier M, Grassi G, Januszewicz A, et al. 2023 ESH Guidelines for the management of arterial hypertension The Task Force for the management of arterial hypertension of the European Society of Hypertension: Endorsed by the International Society of Hypertension (ISH) and the European Renal Association (ERA). *J Hypertens.* 2023;**41**(12):1874-2071. [PubMed ID: 37345492]. <https://doi.org/10.1097/HJH.0000000000003480>.
  34. Onagbiye S, Guddemi A, Baruwa OJ, Alberti F, Odone A, Ricci H, et al. Association of sedentary time with risk of cardiovascular diseases and cardiovascular mortality: A systematic review and meta-analysis of prospective cohort studies. *Prev Med.* 2024;**179**:107812. [PubMed ID: 38081421]. <https://doi.org/10.1016/j.ypmed.2023.107812>.
  35. Oparil S, Acelajado MC, Bakris GL, Berlowitz DR, Cifkova R, Dominiczak AF, et al. Hypertension. *Nat Rev Dis Primers.* 2018;**4**:18014. [PubMed ID: 29565029]. [PubMed Central ID: PMC6477925]. <https://doi.org/10.1038/nrdp.2018.14>.
  36. Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, et al. 2018 ESC/ESH Guidelines for the management of arterial hypertension: The Task Force for the management of arterial hypertension of the European Society of Cardiology and the European Society of Hypertension: The Task Force for the management of arterial hypertension of the European Society of Cardiology and the European Society of Hypertension. *J Hypertens.* 2018;**36**(10):1953-2041. [PubMed ID: 30234752]. <https://doi.org/10.1097/HJH.0000000000001940>.
  37. Zhao M, Xiao M, Tan Q, Lyu J, Lu F. The effect of aerobic exercise on oxidative stress in patients with chronic kidney disease: a systematic review and meta-analysis with trial sequential analysis. *Ren Fail.* 2023;**45**(2):2252093. [PubMed ID: 37753870]. [PubMed Central ID: PMC10538458]. <https://doi.org/10.1080/0886022X.2023.2252093>.
  38. Cornelissen VA, Verheyden B, Aubert AE, Fagard RH. Effects of aerobic training intensity on resting, exercise and post-exercise blood pressure, heart rate and heart-rate variability. *J Hum Hypertens.* 2010;**24**(3):175-82. [PubMed ID: 19554028]. <https://doi.org/10.1038/jhh.2009.51>.
  39. Cornelissen VA, Fagard RH. Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *Hypertension.* 2005;**46**(4):667-75. [PubMed ID: 16157788]. <https://doi.org/10.1161/01.HYP.0000184225.05629.51>.
  40. Canu L, Parenti G, De Filipo G, Mannelli M. Pheochromocytomas and Paragangliomas as Causes of Endocrine Hypertension. *Front Endocrinol (Lausanne).* 2019;**10**:333. [PubMed ID: 31214117]. [PubMed Central ID: PMC6558199]. <https://doi.org/10.3389/fendo.2019.00333>.