Published Online: 2024 November 19



Comparison of Different Concentrations of Antibiotics with Nanoparticles Synthesized by Biological Method on *Pseudomonas aeruginosa* Isolated from The Surface

Fatemeh Mahmoodi¹, Tahereh Eslammanesh², Somayeh Sargolzaei³, Afsaneh Mirshekari⁴, Mehdi Jahantigh^{5,*}

¹ Plant Genetics and Breeding, Department of Plant Breeding and Biotechnology, University of Zabol, Zabol, Iran

² Assistant Professor, Department of Pathology, Faculty of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

³ Biochemistry, Payam Noor University of Mashhad, Mashhad, Iran

⁴ Department of Gastroenterology and Hepatology, Amir Al Momenin Hospital, School of Medicine, Zabol University of Medical Sciences, Zabol, Iran

⁵ Department of Clinical Sciences, Faculty of Veterinary Medicine, University of Zabol, Zabol, Iran

Corresponding Author: Department of Clinical Sciences, Faculty of Veterinary Medicine, University of Zabol, Zabol, Iran. Email: jahantighm@uoz.com

Received: 13 August, 2024; Revised: 18 September, 2024; Accepted: 22 September, 2024

Abstract

Background: The widespread and indiscriminate use of antibiotics, disinfectants, and antiseptics in medical centers has led to the emergence of resistance in *Pseudomonas aeruginosa*, a major cause of hospital infections.

Objectives: This study aims to compare the effects of antibiotics—cotrimoxazole, cephalexin, cefotaxime, cefazolin, gentamicin, imipenem, ampicillin, and ceftazidime—with silver nanoparticles synthesized in medicinal plant extract against *Pseudomonas aeruginosa* bacteria.

Methods: Ten antibiotic-resistant strains of *Pseudomonas aeruginosa* were isolated from surfaces at the maternity hospital of Amir al-Mominin Hospital in Zabul city. In this study, nanoparticles were synthesized using an aqueous extract of the medicinal plant *Capparis*. The minimum inhibitory concentration (MIC) and minimum lethal concentration (MLC) of the synthesized nanoparticles and antibiotics were determined by the microdilution method.

Results: The minimum inhibitory concentrations for the antibiotics ceftriaxone, cephalexin, cotrimoxazole, gentamicin, cefotaxime, imipenem, ceftazidime, ampicillin, and cefazolin were 128, 64, 32, 16, 4, 64, 64, 8, and 64 µg/mL, respectively. The minimum inhibitory concentration for the synthesized silver nanoparticles was 1024 µg/mL, with inhibition observed in four strains at this concentration.

Conclusions: This study demonstrated good antimicrobial effects of synthesized silver nanoparticles on *Pseudomonas aeruginosa*. Therefore, silver nanoparticles may be recommended as a potent antimicrobial agent for disinfecting hospital surfaces.

Keywords: Pseudomonas aeruginosa, Antibiotic, Biosynthesis of Silver, Surfaces

1. Background

Pseudomonas aeruginosa is a gram-negative bacillus, an aerobic, non-fermenting, spore-free, motile bacterium with a capsule-like cell enamel layer on its surface. In nature, it thrives in moist environments and can even survive in distilled water (1). In hospital settings, it is commonly found in incubators, sinks, urinary catheters, and various medical supplies, as well as on employees' clothing and hands.

The rise in antibiotic resistance, particularly multidrug resistance, in this bacterium presents significant challenges, complicating infection treatment (2). Resistance in *Pseudomonas aeruginosa* arises through several mechanisms, including altered permeability to drugs, efflux pumps, receptor modification, access to secondary metabolic pathways,

Copyright © 2024, Mahmoodi et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

and the production of drug-degrading enzymes. Among these, the production of drug-degrading enzymes is a primary mechanism, actively degrading antibiotics and conferring resistance (3).

Pseudomonas aeruginosa has a strong preference for moist environments and is widely found in soil, water, plants, manure, and animal skin. Additionally, it may sometimes reside on the skin, in the external ear, upper respiratory tract, or large intestine of healthy individuals and can cause a range of diseases in humans and livestock. These include urinary tract infections, chronic pneumonia in cystic fibrosis patients, eye and ear infections, wound infections, and mastitis, often with antibiotic resistance (4).

Herbal remedies have long been used to treat various ailments, with approximately 45,000 known plant species (5). Nanotechnology, a rapidly growing field, currently impacts many areas of human life, including industry, agriculture, and medicine. The synthesis of metal nanoparticles has garnered significant interest due to their applications in scientific and industrial fields (6).

Among various nanoparticles, silver nanoparticles have demonstrated wide-ranging benefits in bionanotechnology and medicine (7). In medical science, silver nanoparticles are used for treating burns, producing dental materials, metal coatings, improving water purification, and as disinfectants. They also exhibit antifungal, antiviral, antibacterial, antiinflammatory, anti-angiogenic, and anticancer activities (8).

Recently, biosynthesis using natural and biodegradable agents such as bacteria (9), fungi (10), and plants (11) has been established in various studies and is currently the primary focus of researchers in this field.

In the green synthesis method, metal ions are reduced to nanoparticles using plant compounds in a typically one-step reaction that does not require surfactants or other stabilizing agents. Bioactive substances and compounds in plant extracts, including flavonoids and other water-soluble metabolites, can reduce metal ions to nanoparticles at room temperature (12).

The antimicrobial activity of silver nanoparticles has been reported in numerous studies. The antibacterial mechanism of silver nanoparticles remains a topic of debate and is not yet fully understood. In studies on *Escherichia coli* and *Staphylococcus aureus*, it was observed that silver ions are released by nanoparticles, accumulate around the cell wall or within the cell, and impact DNA replication by interacting with protein thiol groups, causing protein inactivation (13). This interaction leads to the formation of reactive oxygen species (14).

Recent studies have demonstrated the antimicrobial, antioxidant, and anticancer activities of greensynthesized AgNPs (15-17). Biosynthesized silver nanoparticles have also been effectively used to degrade various chemicals (18).

Additionally, green-synthesized silver nanoparticles have a wide range of applications in biotechnology, including water filtration, disinfection, food preservation, cosmetics production, and the development of nano-insecticides and nano-pesticides (19-22).

The genus *Capparis* includes about 150 - 200 species found in tropical and subtropical regions worldwide (23), with four native species in Iran (24). In Iranian traditional medicine, *Capparis* species are used as antiinflammatory, analgesic (for gout and rheumatoid arthritis), diuretic, antibacterial, and anthelmintic agents (25).

Several compounds, including alkaloids, terpenes, flavonoids, and fatty acids, have been reported from different *Capparis* species (26). Scientific studies have shown that these plants possess biological activities, including antioxidant, antibacterial, anti-glycemic, blood fat-lowering, and pain-relieving effects.

The herbal medicine market has surpassed \$60 billion annually and continues to grow (27). Medicinal plants such as *Capparis*, which play a vital role in health systems, are therefore of considerable importance (27).

C. spinosa is one of the most economically significant species in the *Capparidaceae* family, with a vast diversity comprising approximately 40 - 50 genera and 700 - 900 species (28). *Capparis* is native to the Mediterranean basin and is widely distributed from Morocco to Crimea, Armenia, and Iran (29).

Several countries, including Greece, Italy, Spain, and Turkey, have extensive *C. spinosa* production (30). Spain and Turkey produce approximately 1,000 and 4,500 tons of *C. spinosa* (commonly known as snake grass) per year, respectively (31).

Afsharypuor et al. (32) reported that the leaves and ripe fruits of *C. spinosa* contain degradation products of glucosinolates, specifically methyl, isopropyl, and secbutyl isothiocyanates.

2. Objectives

Glucosinolates are plant secondary metabolites of medicinal importance, as they may contribute to disease prevention and reduce the risk of carcinogenesis. Similarly, Çaliş et al. (33) reported the presence of indole-3-acetonitrile glycosides in *C. spinosa* fruits.

3. Methods

3.1. Preparation of Extract

Five grams of powder prepared from the snake grass plant was mixed with 100 cc of deionized water and heated for 15 minutes at boiling temperature. After cooling, it was filtered using Whatman No. 1 paper, and the resulting aqueous extract was stored at 4°C for subsequent tests.

3.2. Synthesis of Nanoparticles

Ten cc of the prepared extract was mixed with 90 cc of a 1 mM silver nitrate solution, and the mixture was placed on a magnetic stirrer for 24 hours at room temperature. To observe color changes, the solution's absorbance was measured using a spectrophotometer in the range of 300 - 700 nm. The solution containing the synthesized nanoparticles was then centrifuged at 1200 rpm for 15 minutes, after which the supernatant was discarded.

3.3. X-ray Diffraction Analysis

The X-ray diffraction (XRD) method was used to examine the metallic nature of the produced nanoparticles. After biological reduction, the silver nanoparticle solution was centrifuged at 8000 rpm for 5 minutes. The resulting pellet was re-dissolved in 10 mL of sterilized deionized water and centrifuged three times to purify it. The final pellet was then dried in a vacuum oven for 24 hours. The structure and composition of the purified nanoparticles were subsequently analyzed using XRD.

3.4. Fourier Transform Infrared Spectroscopy

The dried powder of the produced nanoparticles, along with the leaf extract powder, was mixed with KBr, formed into pellets, and analyzed using the device.

3.5. Isolation of Bacteria

The strains of *Pseudomonas aeruginosa* used in this research were collected and isolated from various surfaces in the maternity department of Amirul Mominin Hospital in Zabol city. To identify *Pseudomonas aeruginosa*, gram staining, catalase, and oxidase tests were conducted, along with confirmatory sugar tests using triple sugar iron agar (TSI).

Subsequently, minimum inhibitory the concentration (MIC) of all strains was determined for different groups including of antibiotics, cephalexin, cefotaxime, cotrimoxazole, cefazolin, gentamicin, imipenem, ampicillin, and ceftazidime, using the agar dilution method according to the clinical and laboratory standards institute (CLSI) guidelines. To determine the MIC, the standard reference powder for each of the antibiotics was used. In the first step, the required amount of each antibiotic for the initial dilution was dissolved in specific solvents. Following this, serial dilutions of each antibiotic were incorporated into Mueller Hinton agar culture medium. After bacterial culturing and 18 hours of incubation, the lowest concentration of each antibiotic at which no colony growth was observed was recorded as the MIC. Data were collected and analyzed using MS Excel 2010 software.

4. Results

The results of this study showed that the lowest inhibitory concentrations of the antibiotics ceftriaxone, cephalexin, cotrimoxazole, gentamicin, cefotaxime, imipenem, ceftazidime, ampicillin, and cefazolin were 128, 64, 32, 16, 4, 64, 64, 8, and 64 μ g/mL, respectively. At these concentrations, 2, 3, 1, 2, 3, 2, 4, 2, and 3 strains of *Pseudomonas aeruginosa* were inhibited (Table 1).

This study showed that the lowest and highest inhibitory concentrations of the antibiotics ceftriaxone, cephalexin, cotrimoxazole, gentamicin, cefotaxime, imipenem, ceftazidime, ampicillin, and cefazolin were 256, 128, 64, 64, 8, 128, 16, and 128 μ g/mL, and 1024, 1024, 1024, 1024, 1024, 1024, 1024, 1024, 1024, 1024, 1024, and 1024 μ g/mL, respectively. At these concentrations, 10, 8, 6, 2, 4, 10, 7, 2, and 13 strains of *Pseudomonas aeruginosa* were destroyed (Table 2).

4.1. The Results of Fourier Transform Infrared Spectrometer Studies Related to Silver Nanoparticles

The results showed that the absorption peak observed in the range of 1680 - 1600 cm⁻¹ is related to the C=C stretching bond vibration, present in all studied samples, indicating the presence of alkene-containing compounds with covalent double bonds, such as ethene or ethylene. The absorption range of 2850 - 3000 cm⁻¹ corresponds to the presence of alkane compounds, such as methane and butane, in the sample. Additionally, the absorption observed in the range of 3600 - 3000 cm⁻¹ is associated with functional groups containing N-H and O-H bonds, suggesting the presence of phenolic compounds.

Bacterial	Ceftriaxone	Cephalexin	Cotrimoxazole	Gentamicin	Cefotaxime	Imi- Penin	Ceftazidime	Ampicillin	Cefazolin
1. P. aeruginosa	256	128	256	512	8	512	512	32	512
2. P. aeruginosa	1024	256	128	128	8	1024	256	125	512
3. P. aeruginosa	256	128	512	64	4	1024	64	64	512
4. P. aeruginosa	256	64	512	64	4	1024	64	512	1024
5. P. aeruginosa	1024	512	64	32	4	256	64	512	1024
6. P. aeruginosa	512	512	64	128	16	256	64	256	512
7. P. aeruginosa	512	1024	128	256	32	1024	256	128	512
8. P. aeruginosa	512	512	128	256	32	1024	512	64	512
9. P. aeruginosa	512	256	1024	32	64	1024	256	64	1024
10. P. aeruginosa	256	64	256	128	8	256	512	16	512
11. P. aeruginosa	256	64	256	256	16	256	512	16	512
12. P. aeruginosa	512	256	128	512	64	1024	1024	128	512
13. P. aeruginosa	512	128	128	32	32	128	256	256	256
14. P. aeruginosa	128	1024	512	16	64	64	128	256	64
15. P. aeruginosa	1024	1024	512	16	32	64	128	128	64
16. P. aeruginosa	128	512	1024	128	64	512	1024	8	512
17. P. aeruginosa	512	512	32	128	16	1024	512	8	64

Abbreviation: MIC, minimum inhibitory concentration.

	Table 2. Minimum Bad	ctericidal Concentration (of Different Antibiotics	Against Pseudomonas	aeruginosa
--	----------------------	----------------------------	--------------------------	---------------------	------------

Bacterial number	Ceftriaxone	Cephalexin	Cotrimoxazole	Gentamicin	Cefotaxime	Imipenem	Ceftazidime	Ampicillin	Cefazolin
1. P. aeruginosa	512	256	512	1024	16	1024	1024	64	1024
2. P. aeruginosa	1024	512	256	256	16	1024	512	256	1024
3. P. aeruginosa	512	256	1024	128	8	1024	128	128	1024
4. P. aeruginosa	512	128	1024	128	8	1024	128	1024	1024
5. P. aeruginosa	1024	1024	128	64	8	512	128	1024	1024
6. P. aeruginosa	1024	1024	128	256	32	512	128	512	1024
7. P. aeruginosa	1024	1024	256	512	64	1024	512	256	1024
8. P. aeruginosa	1024	1024	256	512	64	1024	1024	128	1024
9. P. aeruginosa	1024	512	1024	64	128	1024	512	128	1024
10. P. aeruginosa	512	128	512	256	16	512	1024	32	1024
11. P. aeruginosa	512	128	512	512	32	512	1024	32	1024
12. P. aeruginosa	1024	512	256	1024	128	1024	1024	256	1024
13. P. aeruginosa	1024	256	256	64	64	256	512	512	512
14. P. aeruginosa	256	1024	1024	32	128	128	256	512	128
15. P. aeruginosa	1024	1024	1024	32	64	128	256	256	128
16. P. aeruginosa	256	1024	1024	256	128	1024	1024	16	1024
17. P. aeruginosa	1024	1024	64	256	32	1024	1024	16	128

Abbreviations: MBC, minimum bactericidal concentration.

The vibration observed in the range of 1300 - 1400 cm⁻¹ corresponds to the N=O stretching bond, while the vibration in the range of 1400 - 1600 cm⁻¹ indicates the presence of aromatic compounds. The repeated observation of the absorption range of 2850 - 3000 cm⁻¹ further confirms the presence of alkane-containing

chemical bonds, such as methane and butane, in the sample (Figure 1).

To examine the shape, size, and distribution of the nanoparticles, a transmission electron microscope (TEM) was used. The analysis revealed that the silver



Figure 1. FTIR image related to synthetic silver nanoparticles in aqueous extract

Table 3. Minimum Inhibitory Concentrations and Minimum Bactericidal Concentration of Nano Silver Synthesis in Extract Plant						
Bacterial Number	MIC	MBC				
1	8	16				
2	4	8				
3	4	8				
4	8	16				
5	16	32				
6	16	32				
7	32	64				
8	8	16				
9	8	16				
10	32	64				
11	32	64				
12	64	128				
13	64	128				
14	32	64				
15	32	64				
16	16	32				
17	32	64				

Abbreviations: MIC, minimum inhibitory concentration, MBC, minimum bactericidal concentration.

nanoparticles are spherical, highly dense, and welldispersed, with a size of 150 nm in the stem extract. The MIC of silver nanoparticles synthesized in medicinal plant extract was found to be 4 μ g/mL, with 2 strains inhibited at this concentration. The maximum

inhibitory concentration was 32 μ g/mL, where 6 strains were inhibited (Table 3). The minimum and maximum lethal concentrations of the synthesized nanoparticles were 8 and 128 μ g/mL, respectively, with 2 strains killed at each of these concentrations (Table 3).

5. Discussion

In this study, the MIC of antibiotics for *Pseudomonas aeruginosa* were as follows: Ceftriaxone, 128 μ g/mL; cephalexin, 64 μ g/mL; cotrimoxazole, 32 μ g/mL; gentamicin, 16 μ g/mL; cefotaxime, 4 μ g/mL; imipenem, 64 μ g/mL; ceftazidime, 64 μ g/mL; ampicillin, 8 μ g/mL; and cefazolin, 64 μ g/mL.

Adabi et al. investigated the antibiotic resistance pattern of *Pseudomonas aeruginosa* in Zahedan. The results showed that beta-lactamase-producing strains were resistant to cefoxitin (96%), ceftazidime (15%), ceftriaxone (52%), cefotaxime (96%), cefotetan (30%), aztreonam (8%), and ciprofloxacin (8%) (34).

Habibi et al. studied the frequency and pattern of antibiotic resistance in *Pseudomonas aeruginosa* strains in Tehran hospitals. Their results showed that 88% of the strains were resistant to at least one antibiotic, with the highest resistance observed against cefotaxime (62.9%) and aztreonam (60.4%)(35).

In a study by Hemmati et al. examining *Pseudomonas aeruginosa* in Zanjan, the highest and lowest resistance rates were for cefotaxime (52 isolates, 43.3%) and amikacin (26 isolates, 21.6%), respectively. Resistance was also observed to imipenem in 35 isolates (29.2%), aztreonam in 45 isolates (37.5%), gentamicin in 45 isolates (37.5%), ceftazidime in 35 isolates (29.2%), and ciprofloxacin in 39 isolates (32.5%) (36).

Habibi et al. reported that, among 180 clinical samples, strains isolated from wound sources (44.1%) and urine (29.8%) had the highest frequency. Additionally, 97.5% of isolates exhibited beta hemolytic activity, while only 2.5% showed gamma hemolytic activity. Notably, 88% of isolates were resistant to one or more antibiotics (35).

In their study, Adabi et al. identified and confirmed strains using biochemical and genetic methods. Bacterial resistance to antibiotics was evaluated using the agar disk diffusion method, and MICs were determined for four representative antibiotics. Among 94 *Pseudomonas aeruginosa* strains, 83 (88.3%) exhibited multidrug resistance, with the highest resistance observed for cefepime (89.5%). For the antibiotics tested in MIC determinations, the highest resistance was to ciprofloxacin (89%) (37).

Jamshidi Gohar et al. found that *Pseudomonas aeruginosa* strains were most resistant to cephalothin, doxycycline, cefixime, amoxicillin, ampicillin, amikacin, and trimethoprim (38).

In the study by Rajabpour and Alikhani, the MIC of three different classes of antibiotics was investigated in Pseudomonas aeruginosa strains from hospitalized patients in Hamedan city. Among the eight selected antibiotics, the highest resistance was observed to ciprofloxacin, with 18 strains (58%) resistant, and levofloxacin, with 19 strains (61.2%) resistant. The lowest resistance was observed with imipenem, with only 3 strains (9.6%) resistant. Additionally, 2 strains (6.5%) showed complete resistance to all eight antibiotics. The MIC results showed that for gentamicin, 18 strains (58.06%) were resistant, 8 (25.08%) were intermediate, and 5 (16.12%) were sensitive. For ciprofloxacin, 28 strains (90.32%) were resistant and 3 (9.67%) intermediate. For imipenem, 12 strains (30.7%) were resistant, 13 (41.93%) intermediate, and 6 (19.35%) sensitive (39).

Golshani et al. investigated the resistance pattern of *Pseudomonas aeruginosa* strains in Isfahan hospitals. The resistance levels of the strains were as follows: Ciprofloxacin 56%, gentamicin 59%, tobramycin 61%, amikacin 65%, imipenem 55%, cefepime 55%, ceftazidime 57%, ceftriaxone 60%, cefotaxime 62%, oxacillin 100%, and piperacillin 48% (40).

In a study by Zarenia et al., the resistance pattern of *Pseudomonas aeruginosa* was investigated in samples from Kerman province. The results indicated the highest sensitivity was to cefizoxime, with 55 strains (91.7%), followed by imipenem (54 strains, 90%), meropenem (48 strains, 80%), and ciprofloxacin (40 strains, 66.7%). The highest resistance was observed with cefepime, with 36 strains (60%) resistant, and ciprofloxacin, with 19 strains (31.7%) resistant (41).

Roulova et al. studied the antibiotic resistance pattern of *Pseudomonas aeruginosa* isolated from six hospital wastewater sources. The resistance pattern, determined using the disc diffusion method for seven antibiotics, showed the highest resistance to ciprofloxacin (30.5%), followed by gentamicin (28.8%), meropenem (27.2%), ceftazidime (11.5%), amikacin (11.5%), piperacillin-tazobactam (11.5%), and aztreonam (8.5%) (42).

In the study by Zafer et al., the antibiotic resistance pattern of 122 *Pseudomonas aeruginosa* samples isolated from cancer patients was investigated. Resistance to beta-lactam antibiotics showed the following: Ceftazidime (60.6%), meropenem (45%), aztreonam (45.1%), imipenem (39.3%), and piperacillin/tazobactam (25.4%). Resistance to non-beta-lactam antibiotics included ciprofloxacin (43.4%), amikacin (32.18%), and polymyxin B(2.4%)(43).

In another study, 200 swab samples were obtained from the wound and burn department of an Iraqi teaching hospital. The results showed that out of 200 samples, only 31 were *Pseudomonas aeruginosa*. The antibiotic resistance pattern revealed resistance to gentamicin (83.87%), trimethoprim (67.74%), amikacin (54.83%), ceftazidime and tobramycin (25.80%), and bifloxacin and imipenem (22.58%) (44).

Jahantigh et al. investigated the antibiotic resistance pattern of *Pseudomonas aeruginosa* isolated from wounds at Zahedan Hospital. Their findings showed the highest resistance to trimethoprim/sulfamethoxazole (84.3%), cefipime (70.8%), piperacillin/tazobactam (20.8%), and colistin (8.3%) (45).

The antimicrobial properties of plant extracts are attributed to phenolic compounds and flavonoids, both of which are abundant in *Capparis* plant extracts (46, 47). *Capparis* extract is rich in phenolic, flavonoid, rutin, tocopherol, carotenoid, and vitamin C compounds (31, 48). Other studies have also demonstrated that *Capparis* extract is high in phenolic and flavonoid content (49).

Krishna Raj reported the minimum inhibitory concentration of nanosilver against *Escherichia coli* and *Pseudomonas aeruginosa* as 10 and 20 μ g/mL, respectively, whereas in the present study, these values were 5 and 10 μ g/mL, respectively (50).

In the study by Azizian Shermeh et al., silver nanoparticles were biosynthesized using the aqueous extract of mace plant leaves, and their antimicrobial activity was evaluated. Upon adding the extract to a silver nitrate solution, the solution turned brown. The silver nanoparticles showed maximum absorption at 405 nm, had a spherical shape, and ranged in size from 8 - 12 nm. These nanoparticles exhibited significant antimicrobial activity against tested samples, inhibiting bacterial and fungal growth at very low concentrations (51).

In 2014, Shams et al. investigated the antimicrobial properties of silver nanoparticles derived from lentil seed extract against several gram-positive and gramnegative bacteria. The study confirmed that silver nanoparticles are effective bactericidal agents against a wide range of gram-positive and gram-negative bacteria, including antibiotic-resistant strains (52).

5.1. Conclusions

The results of this study showed that synthesized silver nanoparticles demonstrated a strong inhibitory effect on antibiotic-resistant *Pseudomonas aeruginosa*.

Given that this bacterium has contributed to increased hospital infections, these nanoparticles could potentially serve as a solution to reduce and eliminate bacterial strains.

Acknowledgements

The authors of this article would like to thank all the professors who helped collect and write this article.

Footnotes

Authors' Contribution: All authors read and approved the final manuscript.

Conflict of Interests Statement: The authors declared no conflict of interests.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Funding/Support: The authors declared no financial support to write this manuscript.

References

- Amos D, Joklik WK, Wilfert CM, Willett HP. Zinsser microbiology. 20. ed. Norwalk, Conn: Appleton & Lange; 1992. XI p. 1294 Seiten Illustrationen 29 cm. EN.
- 2. Ahani-Azari A, Danesh A. [Prevalence and antibiotic susceptibility of Pseudomonas aeruginosa isolated from Taleghani hospital, Gorgan-Iran]. J Gorgan Univ Med Sci. 2007;9(3):69-73. FA.
- 3. Peymani A, Farivar TN, Rahimi H, Ranjbar M, Najafipour R. Frequency of class I integron among multidrug resistant Pseudomonas aeruginosa isolates from the selected hospitals in Qazvin and Tehran, Iran. *Qom Univ Med Sci J.* 2014;**8**(3):69-1. FA.
- Qingyu J, Yufeng Y, Kerui X, Xiudong B, Mengchao L, Bei Z, et al. Microbial diversity analysis of subclinical mastitis in dairy cattle in Northeast China. *African Journal of Microbiology Research*. 2015;9(10):687-94. https://doi.org/10.5897/ajmr2014.7172.
- Grover JK, Yadav S, Vats V. Medicinal plants of India with anti-diabetic potential. J Ethnopharmacol. 2002;81(1):81-100. [PubMed ID: 12020931]. https://doi.org/10.1016/s0378-8741(02)00059-4.
- Nadagouda MN, Hoag G, Collins J, Varma RS. Green Synthesis of Au Nanostructures at Room Temperature Using Biodegradable Plant Surfactants. Crystal Growth & Design. 2009;9(11):4979-83. https://doi.org/10.1021/cg9007685.
- Jegadeeswaran P, Shivaraj R, Venckatesh R. Green synthesis of silver nanoparticles from extract of Padina tetrastromatica leaf. *Digest Journal of Nanomaterials and Biostructures*. 2012;7(3):991-8.
- Kalishwaralal K, Deepak V, Ram Kumar Pandian S, Kottaisamy M, BarathmaniKanth S, Kartikeyan B, et al. Biosynthesis of silver and gold nanoparticles using Brevibacterium casei. *Colloids Surf B Biointerfaces*. 2010;77(2):257-62. [PubMed ID: 20197229]. https://doi.org/10.1016/j.colsurfb.2010.02.007.

- Shahverdi AR, Minaeian S, Shahverdi HR, Jamalifar H, Nohi A. Rapid synthesis of silver nanoparticles using culture supernatants of Enterobacteria: A novel biological approach. *Process Biochemistry*. 2007;42(5):919-23. https://doi.org/10.1016/j.procbio.2007.02.005.
- Gajbhiye M, Kesharwani J, Ingle A, Gade A, Rai M. Fungus-mediated synthesis of silver nanoparticles and their activity against pathogenic fungi in combination with fluconazole. *Nanomedicine*. 2009;5(4):382-6. [PubMed ID: 19616127]. https://doi.org/10.1016/j.nano.2009.06.005.
- Shankar SS, Rai A, Ahmad A, Sastry M. Rapid synthesis of Au, Ag, and bimetallic Au core-Ag shell nanoparticles using Neem (Azadirachta indica) leaf broth. J Colloid Interface Sci. 2004;275(2):496-502. [PubMed ID: 15178278]. https://doi.org/10.1016/j.jcis.2004.03.003.
- Mittal AK, Chisti Y, Banerjee UC. Synthesis of metallic nanoparticles using plant extracts. *Biotechnol Adv.* 2013;31(2):346-56. [PubMed ID: 23318667]. https://doi.org/10.1016/j.biotechadv.2013.01.003.
- Feng QL, Wu J, Chen GQ, Cui FZ, Kim TN, Kim JO. A mechanistic study of the antibacterial effect of silver ions on Escherichia coli and Staphylococcus aureus. J Biomed Mater Res. 2000;52(4):662-8.
 [PubMed ID: 11033548]. https://doi.org/10.1002/1097-4636(20001215)52:4<662::aid-jbm10>3.0.co;2-3.
- Matsumura Y, Yoshikata K, Kunisaki S, Tsuchido T. Mode of bactericidal action of silver zeolite and its comparison with that of silver nitrate. *Appl Environ Microbiol*. 2003;**69**(7):4278-81. [PubMed ID: 12839814]. [PubMed Central ID: PMC165194]. https://doi.org/10.1128/AEM.69.7.4278-4281.2003.
- El-Naggar NE, Hussein MH, El-Sawah AA. Bio-fabrication of silver nanoparticles by phycocyanin, characterization, in vitro anticancer activity against breast cancer cell line and in vivo cytotxicity. *Sci Rep.* 2017;7(1):10844. [PubMed ID: 28883419]. [PubMed Central ID: PMC5589729]. https://doi.org/10.1038/s41598-017-11121-3.
- Kedi PBE, Meva FE, Kotsedi L, Nguemfo EL, Zangueu CB, Ntoumba AA, et al. Eco-friendly synthesis, characterization, in vitro and in vivo anti-inflammatory activity of silver nanoparticle-mediated Selaginella myosurus aqueous extract. *Int J Nanomedicine*. 2018;**13**:8537-48. [PubMed ID: 30587976]. [PubMed Central ID: PMC6296690]. https://doi.org/10.2147/IJN.S174530.
- Huq MA. Green Synthesis of Silver Nanoparticles Using Pseudoduganella eburnea MAHUQ-39 and Their Antimicrobial Mechanisms Investigation against Drug Resistant Human Pathogens. Int J Mol Sci. 2020;21(4). [PubMed ID: 32098417]. [PubMed Central ID: PMC7073201]. https://doi.org/10.3390/ijms21041510.
- Fouda A, Abdel-Maksoud G, Abdel-Rahman MA, Eid AM, Barghoth MG, El-Sadany MA. Monitoring the effect of biosynthesized nanoparticles against biodeterioration of cellulose-based materials by Aspergillus niger. *Cellulose*. 2019;26(11):6583-97. https://doi.org/10.1007/s10570-019-02574-y.
- Rafique M, Sadaf I, Rafique MS, Tahir MB. A review on green synthesis of silver nanoparticles and their applications. *Artif Cells Nanomed Biotechnol.* 2017;45(7):1272-91. [PubMed ID: 27825269]. https://doi.org/10.1080/21691401.2016.1241792.
- Burdusel AC, Gherasim O, Grumezescu AM, Mogoanta L, Ficai A, Andronescu E. Biomedical Applications of Silver Nanoparticles: An Up-to-Date Overview. Nanomaterials (Basel). 2018;8(9). [PubMed ID: 30200373]. [PubMed Central ID: PMC6163202]. https://doi.org/10.3390/nano8090681.
- Akter S, Huq MA. Biologically rapid synthesis of silver nanoparticles by Sphingobium sp. MAH-11(T) and their antibacterial activity and mechanisms investigation against drug-resistant pathogenic microbes. Artif Cells Nanomed Biotechnol. 2020;48(1):672-82. [PubMed ID: 32075448]. https://doi.org/10.1080/21691401.2020.1730390.
- 22. Hamouda RA, Hussein MH, Abo-Elmagd RA, Bawazir SS. Synthesis and biological characterization of silver nanoparticles derived from the cyanobacterium Oscillatoria limnetica. *Sci Rep.* 2019;**9**(1):13071.

[PubMed ID: 31506473]. [PubMed Central ID: PMC6736842]. https://doi.org/10.1038/s41598-019-49444-y.

- 23. Trombetta D, Occhiuto F, Perri D, Puglia C, Santagati NA, De Pasquale A, et al. Antiallergic and antihistaminic effect of two extracts of Capparis spinosa L. flowering buds. *Phytother Res.* 2005;**19**(1):29-33. [PubMed ID: 15799005]. https://doi.org/10.1002/ptr.1591.
- 24. Mozafarian V. Dictionary of Iranian plants. Netherlands: Moaser; 1998.
- 25. Aghili MH. [Makhzan al-Advie]. Iran Univ Med Sci. 2008:549-8. FA.
- Yang T, Wang C, Chou G, Wu T, Cheng X, Wang Z. New alkaloids from Capparis spinosa: Structure and X-ray crystallographic analysis. *Food Chemistry*. 2010;**123**(3):705-10. https://doi.org/10.1016/j.foodchem.2010.05.039.
- 27. Gunjan M, Naing TW, Saini RS, Ahmad A, Naidu JR, Kumar I. Marketing trends & future prospects of herbal medicine in the treatment of various disease. *World Journal of Pharmaceutical Research*. 2015;**4**(9):132-55.
- Hall JC, Sytsma KJ, Iltis HH. Phylogeny of Capparaceae and Brassicaceae based on chloroplast sequence data. Am J Bot. 2002;89(11):1826-42. [PubMed ID: 21665611]. https://doi.org/10.3732/ajb.89.11.1826.
- 29. Ramezani-Gask M, Bahrani MJ, Shekafandeh A, Salehi H, Taghvaei M, Al-Ahmadi MJ. A comparison of different propagation methods of common Caper-bush (Capparis spinosa L.) as a new horticultural crop. *Int J Plant Dev Biol*. 2008;**2**(2):106-10.
- Inocencio C, Rivera D, Alcaraz F, Tomás-Barberán FA. Flavonoid content of commercial capers (Capparis spinosa, C. sicula and C. orientalis) produced in mediterranean countries. European Food Research and Technology. 2000;212(1):70-4. https://doi.org/10.1007/s002170000220.
- Tilii N, Elfalleh W, Saadaoui E, Khaldi A, Triki S, Nasri N. The caper (Capparis L.): ethnopharmacology, phytochemical and pharmacological properties. *Fitoterapia*. 2011;82(2):93-101. [PubMed ID: 20851750]. https://doi.org/10.1016/j.fitote.2010.09.006.
- Afsharypuor S, Jeiran K, Jazy AA. First investigation of the flavour profiles of the leaf, ripe fruit and root of Capparis spinosa var. mucronifolia from Iran. *Pharmaceutica Acta Helvetiae*. 1998;**72**(5):307-9. https://doi.org/10.1016/s0031-6865(97)00023-x.
- Çaliş İ, Kuruüzüm A, Rüedi P. 1H-Indole-3 acetonitrile glycosides from Capparis spinosa fruits. *Phytochemistry*. 1999;**50**(7):1205-8. https://doi.org/10.1016/s0031-9422(98)00669-4.
- Adabi J, Shahraki Zahedan S, Bokaeian M, Tahmasebi H. The phenotype of antibiotic resistance in Pseudomonas aeruginosa isolates producing ESBL in Zahedan 2014. Pars of Jahrom University of Medical Sciences. 2017;15(1):7-15. https://doi.org/10.29252/jmj.15.1.7.
- 35. Habibi A, Mozafari A, Fallah Mehrabadi J, Kazemi Darsanky R. [The frequency and antibiotic resistance modeling in clinical strains of Pseudomonas aeruginosa isolated from different parts of Tehran hospitals]. *Razi J Med Sci.* 2016;**23**(146):10-6.
- 36. Hemmati F, Sourouri Zanjani R, Haghi F, Zeighami H. [Determination of antibiotic resistance profile and frequency of metallo-betalactamases in Pseudomonas aeruginosa isolates]. *J Zanjan Univ Med Sci Health Services*. 2014;**22**(93):77-85. FA.
- Adabi M, Talebi Taher M, Arbabi L, Afshar M, Fathizadeh S, Minaeian S, et al. [Determination of antibiotic resistance pattern of Pseudomonas aeruginosa strains isolated from patients with burn wounds]. J Ardabil Univ Med Sci. 2015;15(1):66-74. FA.
- Jamshidi Gohar M, Rahimi Fard N, Hosseini Doust SR. The pattern of antibiotic resistance within clinical isolates of Pseudomonas aeruginosa and detection of AmpC. *Medical Sciences Journal*. 2018;28(3):212-9. https://doi.org/10.29252/iau.28.3.212.
- 39. Rajabpour M, Alikhani MY. [MIC determination of Pseudomonas aeruginosa strains were isolated from clinical specimens of patients

admitted to educational hospitals in Hamedan (90-91)]. *Iran J Med Microbiol*. 2013;7(3):18-25. FA.

- 40. Golshani Z, Ahadi AM, Sharifzadeh A. [Identification and determination of the antibiotic resistance pattern of Pseudomonas aeruginosa strains isolated from patients hospitalized in Isfahan hospitals with multiple resistance]. *J Veterinary Laboratory Res.* 2013;1(4):119. FA.
- 41. Zarenia M, Salehi M, Kheirkhah B. Identification of Drug-Repellent Encoded Genes (Efflux pump) in Pseudomonas aeruginosa Isolated from Stool Samples, Kerman Province. *journal of ilam university of medical sciences*. 2017;**25**(4):108-16. https://doi.org/10.29252/sjimu.25.4.108.
- Roulova N, Mot'kova P, Brozkova I, Pejchalova M. Antibiotic resistance of Pseudomonas aeruginosa isolated from hospital wastewater in the Czech Republic. J Water Health. 2022;20(4):692-701. [PubMed ID: 35482385]. https://doi.org/10.2166/wh.2022.101.
- Zafer MM, Al-Agamy MH, El-Mahallawy HA, Amin MA, Ashour MS. Antimicrobial resistance pattern and their beta-lactamase encoding genes among Pseudomonas aeruginosa strains isolated from cancer patients. *Biomed Res Int.* 2014;2014:101635. [PubMed ID: 24707471]. [PubMed Central ID: PMC3953503]. https://doi.org/10.1155/2014/101635.
- 44. Rashid Mahmood A, Mansour Hussein N. Study of antibiotic resistant genes in Pseudomonas aeroginosa isolated from burns and wounds. *Archives of Razi Institute.* 2022;77(1):403-11. https://doi.org/10.22092/ARI.2021.356681.1893.
- Jahantigh M, Tahmasbi H, Bakaeiyan M. [Antibiotic Resistance Pattern of Pseudomonas aeruginosa Strains Isolated from Burn Patients]. Medical J Mashhad Univ Med Sci. 2018;60(6):1121-31. FA. https://doi.org/10.22038/mjms.2019.14778.

- Zia-Ul-Haq M, Cavar S, Qayum M, Imran I, de Feo V. Compositional studies: antioxidant and antidiabetic activities of Capparis decidua (Forsk.) Edgew. Int J Mol Sci. 2011;12(12):8846-61. [PubMed ID: 22272107]. [PubMed Central ID: PMC3257104]. https://doi.org/10.3390/ijms12128846.
- 47. Imran Imran II, Muhammad Zia-ul-Haq MZ, Calani L, Mazzeo T, Pellegrini N. Phenolic profile and antioxidant potential of selected plants of Pakistan. J Applied Botany Food Quality. 2014;87:35-0. https://doi.org/10.5073/JABFQ.2014.087.005.
- Tlili N, Nasri N, Khaldi A, Triki S, MunnÉ-Bosch S. Phenolic Compounds, Tocopherols, Carotenoids and Vitamin C of Commercial Caper. *Journal of Food Biochemistry*. 2011;35(2):472-83. https://doi.org/10.1111/j.1745-4514.2010.00396.x.
- 49. Sivropoulou A, Kokkini S, Lanaras T, Arsenakis M. Antimicrobial activity of mint essential oils. *Journal of Agricultural and Food Chemistry*. 2002;**43**(9):2384-8. https://doi.org/10.1021/jf00057a013.
- 50. Krishnaraj C, Jagan EG, Rajasekar S, Selvakumar P, Kalaichelvan PT, Mohan N. Synthesis of silver nanoparticles using Acalypha indica leaf extracts and its antibacterial activity against water borne pathogens. *Colloids Surf B Biointerfaces*. 2010;**76**(1):50-6. [PubMed ID: 19896347]. https://doi.org/10.1016/j.colsurfb.2009.10.008.
- Azizian Shermeh O, Valizadeh J, Noroozifar M, Qasemi A. Investigating the Antimicrobial Activities of Silver Nanoparticles Biosynthesized by Aqueous Extract of Sambucus ebulus L. *journal of ilam university of medical sciences*. 2016;24(5):92-108. https://doi.org/10.18869/acadpub.sjimu.24.5.92.
- 52. Shams S, Pourseyedi S, Hashemipour Rafsanjani H. Green synthesis of silver nanoparticles: eco-friendly and antibacterial. *International Journal of nanoscience and nanotechnology*. 2014;**10**(2):127-32.