



Characteristics of College Students Attending the Smoking Cessation Clinic at Sultan Qaboos University Hospital: A Cross-Sectional Study

AbdulMalik AL Attabi¹, Majid Al Rawahi¹, Hana AlSumri^{2,*}

¹ College of Medicine and Health Sciences, Sultan Qaboos University, Muscat, Oman

² Department of Family Medicine and Public Health, College of Medicine and Health Sciences, Sultan Qaboos University, Muscat, Oman

*Corresponding Author: Department of Family Medicine and Public Health, College of Medicine and Health Sciences, Sultan Qaboos University, Muscat, Oman. Email: alsumry@squ.edu.om

Received: 28 December, 2024; Revised: 25 February, 2025; Accepted: 5 March, 2025

Abstract

Background: Smoking cessation clinics provide essential support and resources for individuals seeking to quit smoking.

Objectives: This study aimed to describe the sociodemographic and clinical characteristics of college students attending a university smoking cessation clinic in Oman, focusing on factors such as age and the motivations behind both starting and quitting smoking.

Materials and Methods: This retrospective cross-sectional study involved 63 college students who attended the smoking cessation clinic between the years of 2013 to 2020. Data regarding sociodemographic and smoking characteristics were collected from the electronic medical records. Descriptive and inferential statistics were carried out using a statistical software program.

Results: The vast majority of college students were male (98.4%), with a median age of smoking initiation of 16 years (range: 8-22 years). Most students exhibited high (30.2%) or moderate (28.6%) levels of nicotine dependence, with the most common utilized type of tobacco product being the midwakh (pipe) (79.4%) and cigarettes (60.3%). The primary reason for smoking initiation was friends (66.7%), while health-related concerns were the most common reason for quitting (65.1%). Students from scientific colleges more frequently quit for health-related reasons than non-scientific college students (82.4% versus 46.2%; $P = 0.040$). Students with low nicotine dependency levels more frequently quit due to social image compared to those with moderate or high dependency levels (47.1% versus 34.8% and 10.5%, respectively; $P = 0.050$). Moreover, students with low nicotine dependency levels more frequently started smoking out of curiosity, compared to those with moderate and high dependency levels (58.8% versus 9.5% and 27.8%, respectively; $P = 0.004$).

Conclusions: This study provides valuable insights into college students' smoking behaviors at a university smoking cessation clinic. Understanding the reasons behind decisions to start and stop smoking can help inform tailored smoking cessation interventions and campaigns.

Keywords: Smoking, Health Risk Behaviors, Smoking Cessation, Nicotine Dependence, Students, University, Oman

1. Background

Smoking is a major public health issue worldwide, causing significant preventable morbidity and mortality (1). According to the World Health Organization (WHO), Smoking is responsible for more than 8 million deaths each year, predominantly in low- and middle-income countries (2). Smoking not only affects the smoker but also poses a significant risk to non-smokers through exposure to secondhand smoke (3). Smoking has been linked to several health problems,

including cardiovascular and respiratory diseases, impaired wound healing, and reduced fertility (2, 4). Furthermore, Smoking has been shown to increase rates of depression and stress among smokers and has been linked to decreased productivity and increased healthcare costs (5, 6).

Despite the well-documented risks associated with this behavior, Smoking remains prevalent in many countries (2). In Oman, the WHO estimated that approximately 11% of the local population were smokers

in 2010, affecting approximately 18% of men and 1% of women. If current tobacco control measures remain unchanged, the WHO predicts that Smoking prevalence could increase to 23% by 2025, with approximately 33% of men smoking, while the rate among women is expected to remain at around 1% (7).

Several studies have investigated the characteristics of college student smokers and associated predictors for smoking behaviors. Findings suggest that sociodemographic factors such as gender and socioeconomic status play a significant role in college students' smoking behaviors in the Eastern Mediterranean region (8, 9). In Oman, previous research from 2008 involving 1,962 school students indicated that 26.6% reported having smoked tobacco at some point in their lives, with 9.6% currently using tobacco products (10). Among males, 15.5% were current users, whereas only 2.6% of females reported currently smoking. Another study from Oman showed that 4.6% of secondary school students were current smokers (11). The probability of participants smoking increased if they had a friend or parent who smoked. However, adolescents were less inclined to smoke if they believed that Smoking had adverse health effects (10).

Evidence suggests that smoking cessation interventions can effectively help college students quit smoking (12). In Oman, the Sultan Qaboos University Hospital (SQUH) in Muscat Governorate offers a dedicated smoking cessation clinic integrated into the university's primary student health services since 2013. This clinic offers psychological counseling and nicotine replacement therapy to college students registered at the affiliated Sultan Qaboos University. However, despite the relatively high prevalence of Smoking among college students in Oman, there is limited information on the number and characteristics of those who utilize this service. Such knowledge is essential for healthcare providers to assess clinic outcomes and design tailored interventions that address their specific needs and motivations to quit.

2. Objectives

This study, therefore, aimed to investigate the sociodemographic and clinical characteristics of college students attending the SQUH Smoking Cessation Clinic from the year 2013 to 2020 using the information available in the electronic health records. Baseline nicotine dependency levels were also assessed using the Fagerström test for nicotine dependence (13).

Furthermore, the study sought to explore the reasons behind the students' smoking behaviors and their motivations for quitting.

3. Materials and Methods

3.1. Study Design and Participants

A retrospective cross-sectional study was conducted among 63 college students attending the SQUH Smoking Cessation Clinic from 2013 to 2020. The inclusion criteria encompassed all undergraduate and graduate college students, regardless of gender and nationality, who smoked and attended the clinic during this period. Non-university students and SQUH employees attending the cessation clinic were excluded. Data collection continued until the end of 2020, when the clinic temporarily ceased services due to the global Coronavirus Disease 2019 pandemic.

3.2. Data Collection

Relevant data were collected from the students' electronic medical records. Sociodemographic information such as gender, age, place of residence, and specific college was gathered. In addition, clinical and smoking-related characteristics were assessed, including the presence of comorbidities, age at first clinic visit, type of tobacco used, nicotine dependence level, age of smoking initiation, reasons for starting to smoke, and motivations for quitting. Nicotine dependence was assessed using the Fagerström test for nicotine dependence by the treating physician. Dependency levels were classified as low (scores of 1 - 2), low-to-moderate (scores of 3 - 4), moderate (scores of 5 - 7), or high (scores of ≥ 8). The Fagerström test for nicotine dependence has demonstrated reliability in assessing nicotine dependence across various settings and populations (14).

3.3. Statistical Analysis

Collected data were analyzed using the statistical package for the social sciences (SPSS) software, version 26 (IBM Corp., Armonk, New York, USA). The analysis included both descriptive statistics and inferential statistics. Categorical variables were reported as frequencies and percentages, while continuous categorical variables were presented as medians with ranges. Associations between dependent and independent variables were analyzed using appropriate significance tests, including chi-squared, Mann-Whitney

U, and Kruskal-Wallis tests. A P-value of ≤ 0.05 was considered statistically significant. The Kolmogorov-Smirnov test was employed to explore the normality of distribution between the independent continuous variables and the dependent categorical variables. A P-value of < 0.05 indicated a non-normal distribution. Place of residence was categorized as Muscat versus elsewhere, while type of college attended was divided into scientific (agricultural and marine sciences, engineering, science, medicine and health science, and nursing) and non-scientific (art and social sciences, economics and political science, education, and law) categories.

3.4. Ethical Considerations

This study was approved by the Medical Research and Ethics Committee of the College of Medicine and Health Sciences, Sultan Qaboos University (MREC #3033).

4. Results

4.1. Baseline Characteristics

Most college students who attended the SQUH Smoking Cessation Clinic were male (98.4%), with only one female participant (1.6%). The median age at smoking initiation was 16 years (range: 8 - 22 years old), while the median age upon first clinic visit was 21 years (range: 17 - 26 years old). The most frequently represented colleges were the College of Economics & Political Science and the College of Engineering (15.9% each), followed by the College of Education (11.1%). Most students originated from Al-Sharqiyah South (15.9%), followed by Muscat and Al-Batinah North (14.3%). The majority smoked midwakh pipes (79.4%) and cigarettes (60.3%). Additionally, 30.2% smoked Shisha Waterpipes, and 17.5% used Afdhal Chewing Tobacco. Only three students (4.8%) used electronic cigarettes or vape pens (Table 1).

Upon their first clinic visit, the majority of students exhibited either high (30.2%) or moderate (28.6%) levels of nicotine dependence (Figure 1). In terms of the reasons for smoking initiation, 42 students (66.7%) started smoking due to peer influence, 17 (27.0%) out of curiosity, 14 (22.2%) due to stress and emotional reasons, 10 (15.9%) due to the influence of a family member who smoked, and 10 (15.9%) for social reasons. Health-related concerns were the most common reason to quit smoking (65.1%), followed by family reasons (34.9%), social image (28.6%), and religious beliefs and feelings of

guilt (23.8%). Additionally, 14 students (22.2%) indicated a desire to quit smoking due to peer influence, while only three (4.8%) attempted to quit either because they wanted to marry or were already married (Table 2).

4.2. Associations Between Smoking Behaviors and Sociodemographic Characteristics

Age at smoking initiation was not associated with the level of nicotine dependence, place of residence, or type of college attended ($P > 0.05$) (Table 3). However, significantly more students from scientific colleges wanted to quit smoking for health-related reasons compared to students attending non-scientific colleges (82.4% versus 46.2%; $P = 0.040$). In addition, students with low nicotine dependence levels more frequently began smoking out of curiosity compared to those with moderate or high nicotine dependence levels (58.8% versus 9.5% and 27.8%, respectively; $P = 0.050$). Similarly, students with low nicotine dependence levels more frequently identified social image as a reason to quit smoking compared to those with moderate or high dependence levels (47.1% versus 34.8% and 10.5%, respectively; $P = 0.004$) (Table 4).

5. Discussion

This study is one of the first in Oman to explore students' sociodemographic differences, tobacco consumption types, and levels of nicotine addiction at the time of their first visit to a smoking cessation clinic. It also sheds light on key factors influencing smoking initiation, such as peer pressure and curiosity, as well as common motivations for quitting, including health-related concerns. The findings offer valuable insights for healthcare providers and public health organizations to design tailored smoking cessation campaigns and educational programs targeting high school and college students.

This study investigated the characteristics of college students attending the SQUH Smoking Cessation Clinic, revealing a predominance of male gender students (98.4%). This likely reflects the low smoking rates among Omani female students. These findings are consistent with a systematic review that identified a similar pattern among Saudi college students, where male gender was strongly associated with smoking behaviors (15). Additionally, research across five countries in the Eastern Mediterranean region, including Egypt, Jordan, Occupied Palestinian Territories, Oman, and the United Arab Emirates, has also indicated a higher prevalence of

Table 1. Characteristics of College Students Attending a University Hospital Smoking Cessation Clinic in Muscat, Oman (N = 63)

Characteristic	Values ^a
Gender	
Male	62 (98.4)
Female	1 (1.6)
Median age in years (range)	
At smoking initiation (8 - 22)	16
Upon first clinic visit (17 - 26)	21
College	
Agricultural and Marine Sciences	4 (6.3)
Arts and Social Sciences	6 (9.5)
Economics and Political Science	10 (15.9)
Education	7 (11.1)
Engineering	10 (15.9)
Law	3 (4.8)
Nursing	0 (0)
Medicine and Health Sciences	0 (0)
Science	3 (4.8)
Unknown	20 (31.7)
Place of residence	
Ad Dakhiliyah	8 (12.7)
Ad Dhahirah	5 (7.9)
Al Batinah North	9 (14.3)
Al Batinah South	4 (6.3)
Al Buraymi	2 (3.2)
Al Wusta	1 (1.6)
Ash Sharqiyah North	7 (11.1)
Ash Sharqiyah South	10 (15.9)
Dhofar	8 (12.7)
Muscat	9 (14.3)
Musandam	0 (0)
Type of tobacco product used ^b	
Pipe (midwakh)	50 (79.4)
Cigarettes	38 (60.3)
Hookah/waterpipe (shisha)	19 (30.2)
Chewing tobacco (Afdhal)	11 (17.5)
E-cigarette/vape pen	3 (4.8)
Unknown	3 (4.8)

^a Values are expressed as No. (%) unless otherwise indicated.

^b Percentages may not add up to 100% as some individuals may have used more than one type of tobacco product.

waterpipe smoking among male university students, as well as a lower perceived risk of harm from engaging in this behavior (8). This underscores the need for targeted interventions addressing the unique challenges and influences driving smoking behaviors in this group.

Similarly, the present study also identified an early age of smoking initiation, with a median age of 16 years, which aligns with patterns reported in Oman, where the mean age of starting to smoke was 18.7 years old (16).

Two studies conducted in Saudi Arabia also reported a mean age of 16 years for the initiation of tobacco use, highlighting the early onset of smoking behaviors during adolescence (17, 18). Regarding geographical distribution, most smokers in the current study originated from Al-Sharqiyah South, followed by Muscat and Al-Batinah North. This information is significant as it can help guide the development of targeted smoking awareness and outreach programs in these regions.




Figure 1. Baseline nicotine dependency levels among college students attending a university hospital Smoking Cessation Clinic in Muscat, Oman (N = 63). Dependency was assessed using the Fagerström test for nicotine dependence, with scores of 1 - 2, 3 - 4, 5 - 7, and ≥ 8 classified as low, low-to-moderate, moderate, and high nicotine dependence, respectively.

Furthermore, further investigation into cultural or environmental factors could help explain the higher prevalence of smoking in these regions.

Our research revealed that college students utilized various forms of tobacco products, with many participating in culturally ingrained practices such as smoking midwakh pipes (79.4%) and Shisha Waterpipes (30.2%). Cigarettes were also commonly used (60.3%), while a minority opted for Afdhal chewing tobacco (17.5%) and electronic cigarettes or vape pens (4.8%). A previous study of adult Omani smokers documented that the majority (82.9%) smoked manufactured cigarettes, with smaller percentages smoking Shisha (6.4%), Gadou (traditional waterpipes; 7.9%), pipes (7.7%),

and other forms of tobacco products, like chewable tobacco (4.5%). Acknowledging and respecting these cultural preferences while designing effective smoking cessation programs is imperative.

The study also underscores the significant challenge of nicotine addiction, given that the majority of the cohort demonstrated either high or moderate levels of dependence at their first visit to the smoking cessation clinic. Similarly, a study conducted among students at King Faisal University in Saudi Arabia found that 47.0% of current smokers exhibited high nicotine dependence levels according to the Fagerström test for nicotine dependence (19). This highlights the necessity of developing tailored, intensive cessation strategies,

Table 2. Reasons for Smoking Initiation and Cessation Among College Students Attending a University Hospital Smoking Cessation Clinic in Muscat, Oman (N = 63)

Reason	No. (%)
For starting to smoke	
Friends	42 (66.7)
Curiosity	17 (27)
Stress/emotional reasons	14 (22.2)
Family	10 (15.9)
Social reasons	10 (15.9)
Unknown	5 (7.9)
For deciding to quit ^a	
Health	41 (65.1)
Family	22 (34.9)
Social image	18 (28.6)
Religious reasons/guilt	15 (23.8)
Friends	14 (22.2)
Marriage	3 (4.8)
Unknown	2 (3.2)

^a Percentages may not add up to 100% as some individuals may have had more than one reason for quitting.

Table 3. Associations Between Age of Smoking Initiation and Various Characteristics Among College Students Attending a University Hospital Smoking Cessation Clinic in Muscat, Oman (N = 63)

Factors	Mean Rank	P-Value
Nicotine dependency level		0.082 ^a
Low	37.12	
Moderate	25.75	
High	27.03	
Place of residence		0.920 ^b
Muscat	30.56	
Elsewhere	29.91	
College type		0.877 ^b
Scientific	21.15	
Non-scientific	21.74	

^a Using a Kruskal-Wallis test.

^b Using a Mann-Whitney U test.

including educational campaigns targeting adolescents early to prevent the uptake of tobacco products or to assist those who are still in the mild stages of addiction, as quitting is more manageable at lower levels of dependence.

Understanding the reasons for starting and quitting provides valuable insights into the motivations behind college students' smoking behaviors. In this study, peer influence was the most prevalent reason for smoking initiation (66.7%), followed by curiosity (27.0%) and stress or emotional reasons (22.2%). Similarly, a study from neighboring Saudi Arabia also indicated that

curiosity was the primary reason cited for smoking, cited by 44.3% of respondents, with tension relief following closely behind at 26.1%. These findings suggest that social dynamics and the exploratory nature play a significant role in initiating this behavior. Additionally, numerous studies have identified the behaviors and attitudes of friends as significant influencers of individual behavior (20). Regarding cessation, we found health-related reasons to constitute the most common motivation for quitting (65.1%), underscoring the importance of health awareness in driving individual cessation efforts. This finding aligns with a previous

Table 4. Associations Between Reasons for Smoking Initiation/Cessation and Various Characteristics Among College Students Attending a University Hospital Smoking Cessation Clinic in Muscat, Oman (N = 63)^a

Factors	College Type		Nicotine Dependency Level			Nicotine Dependency Level			P-Value
	Scientific	Non-scientific	Low	Moderate	High	Low	Moderate	High	
Health as reason for quitting									0.040 ^b
Yes	14 (82.4)	12 (46.2)							
No	3 (17.6)	14 (53.8)							
Social image as reason for quitting									0.050
Yes			8 (47.1)	8 (34.8)	2 (10.5)				
No			9 (52.2)	15 (65.2)	17 (89.5)				
Curiosity as reason for starting to smoke									0.004
Yes						10 (58.8)	2 (9.5)	5 (27.8)	
No						7 (41.2)	19 (90.5)	13 (72.2)	

^a Values are presented as No. (%).^b Using Pearson's chi-squared test.

study, which showed that approximately 70% of smokers quit due to increased awareness of the hazards associated with smoking, highlighting the potential effectiveness of health-focused interventions in promoting smoking cessation (17).

Interestingly, a later age of smoking initiation was linked with lower nicotine dependence, although this trend was not statistically significant ($P = 0.082$). The higher mean age of smoking initiation in the low nicotine dependence group could indicate that individuals with lower nicotine dependence may start smoking later compared to those with moderate or high dependence. This finding suggests a complex relationship that warrants further investigation, possibly with a larger sample size to enhance the study's power to detect genuine associations between these variables. Moreover, no significant relationship was found between the age of smoking initiation and place of residence or type of college attended.

Nonetheless, the associations between reasons for quitting smoking and sociodemographic factors provide insights into the motivations behind smoking cessation among college students. In particular, a significantly higher percentage of students from scientific colleges cited health-related concerns as their primary motivation to quit smoking compared to non-scientific colleges ($P = 0.040$), reaffirming the influence of educational background in shaping attitudes towards smoking cessation. Additionally, the link between social image and nicotine dependence ($P = 0.050$), as well as curiosity and nicotine dependence ($P = 0.004$), highlights key psychosocial factors affecting

smoking behaviors. Specifically, curiosity as a reason to start smoking was significantly associated with low nicotine dependence, while social image represented a significant motivator for quitting in the same group. These findings underscore the necessity for tailored interventions addressing both the physiological and psychosocial aspects of nicotine dependence.

The findings of this study should be interpreted in light of its limitations. Results were based on data collected from a single center, which provides smoking-quitting services only to college students attending one affiliated university. This impacted the generalizability of the study findings about students attending other public or private colleges. Furthermore, the sample size was small, consisting of only 63 college students. This limited sample size could have affected the study's statistical power, potentially influencing the results of the association analyses. Additionally, the cross-sectional nature of the study design and the retrospective data collection method could have affected the availability of certain information, leading to missing data.

5.1. Conclusions

The findings of this study provide essential baseline information that may help healthcare providers and public health organizations develop tailored smoking cessation campaigns and educational programs targeting college students. These initiatives should emphasize the adverse health consequences of smoking while taking into account common motives for starting and the types of tobacco that lead to dependence.

Moreover, the low uptake of the freely available Smoking Cessation Clinic on campus, despite the likely high prevalence of student smokers, highlights the need for better promotion of these services and the benefits of quitting smoking among this population. Future studies could benefit from a prospective longitudinal design to address the limitations of missing data. Expanded smoking cessation services across Oman would also provide a more comprehensive understanding of the health profile of affected patients, offering a better representation of the broader population.

Acknowledgements

The researchers would like to thank the relevant authorities at SQUH for granting access to the electronic health records of participating students, which facilitated data collection for this study.

Footnotes

Authors' Contribution: Study Concept and design: Hana AlSumri; Acquisition of data: Hana AlSumri, AbdulMalik AL Attabi, and Majid Al Rawahi; Analysis and interpretation of data: Hana AlSumri, AbdulMalik AL Attabi, and Majid Al Rawahi; Drafting of the manuscript: AbdulMalik AL Attabi, Hana AlSumri, and Majid Al Rawahi; Critical revision of the manuscript for important intellectual content: Hana AlSumri; Statistical analysis: Hana AlSumri, AbdulMalik AL Attabi, and Majid Al Rawahi; Administrative, technical, and material support: Hana AlSumri, AbdulMalik AL Attabi, and Majid Al Rawahi; Study supervision: Hana AlSumri.

Conflict of Interests Statement: One author is an Assistant Professor at SQU, and one author holds an unpaid membership in a government or non-governmental organization.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication. The data are not publicly available due to the sensitivity and confidentiality of patient information.

Ethical Approval: This study was approved by the Medical Research and Ethics Committee of the College of Medicine and Health Sciences, Sultan Qaboos University (MREC #3033).

Funding/Support: This research was not supported financially with any grant.

References

- Centers for Disease Control and Prevention. *Smoking and tobacco use: Fast facts*. Centers for Disease Control and Prevention; 2024, [cited 2024]. Available from: https://www.cdc.gov/tobacco/data_statistics/fact_sheets/fast_facts/index.htm.
- World Health Organization. *Fact sheets: Tobacco*. Geneva, Switzerland: World Health Organization; 2024, [cited 2024]. Available from: <https://www.who.int/news-room/fact-sheets/detail/tobacco>.
- Al-Lawati JA, Al-Thuhli Y, Qureshi F. Measuring Secondhand Smoke in Muscat, Oman. *Sultan Qaboos Univ Med J*. 2015;**15**(2):e288-91. [PubMed ID: 26052464]. [PubMed Central ID: PMC4450794].
- National Institute on Drug Abuse. *Cigarettes and other tobacco products*. 2024, [cited 2024]. Available from: https://www.myflfamilies.com/sites/default/files/2023-05/drugfacts_cigarettes2018.pdf.
- Mental Health Foundation. *Smoking and mental health*. 2024, [cited 2024]. Available from: <https://www.mentalhealth.org.uk/explore-mental-health/a-z-topics/smoking-and-mental-health>.
- Centers for Disease Control and Prevention. *Smoking and tobacco use: Health effects of cigarette smoking*. Centers for Disease Control and Prevention; 2024, [cited 2024]. Available from: https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm.
- World Health Organization. *WHO global report on trends in prevalence of tobacco smoking*. Geneva, Switzerland: World Health Organization; 2024, [cited 2024]. Available from: https://iris.who.int/bitstream/handle/10665/156262/9789241564922_eng.pdf?sequence=1.
- Abu-Rmeileh NME, Alkhuffash O, Kheirallah K, Mostafa A, Darawad M, Al-Farsi Y, et al. Harm perceptions of waterpipe tobacco smoking among university students in five Eastern Mediterranean Region countries: A cross-sectional study. *Tob Induc Dis*. 2018;**16**:20. [PubMed ID: 31516420]. [PubMed Central ID: PMC6659482]. <https://doi.org/10.18332/tid/89966>.
- Hitchman SC, Fong GT, Zanna MP, Thrasher JF, Chung-Hall J, Siahpush M. Socioeconomic status and smokers' number of smoking friends: findings from the International Tobacco Control (ITC) Four Country Survey. *Drug Alcohol Depend*. 2014;**143**:158-66. [PubMed ID: 25156228]. [PubMed Central ID: PMC4209373]. <https://doi.org/10.1016/j.drugalcdep.2014.07.019>.
- Al-Lawati JA, Muula AS, Hilmi SA, Rudatsikira E. Prevalence and Determinants of Waterpipe Tobacco Use among Adolescents in Oman. *Sultan Qaboos Univ Med J*. 2008;**8**(1):37-43. [PubMed ID: 21654955]. [PubMed Central ID: PMC3087736].
- Jaffer YA, Afifi M, Al Ajmi F, Alouhaishi K. Knowledge, attitudes and practices of secondary-school pupils in Oman: I. health-compromising behaviours. *East Mediterr Health J*. 2006;**12**(1-2):35-49. [PubMed ID: 17037220].
- Cobb NK, Graham AL, Abrams DB. Social network structure of a large online community for smoking cessation. *Am J Public Health*. 2010;**100**(7):1282-9. [PubMed ID: 20466971]. [PubMed Central ID: PMC2882421]. <https://doi.org/10.2105/AJPH.2009.165449>.
- Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO. The Fagerstrom Test for Nicotine Dependence: a revision of the

- Fagerstrom Tolerance Questionnaire. *Br J Addict.* 1991;**86**(9):1119-27. [PubMed ID: 1932883]. <https://doi.org/10.1111/j.1360-0443.1991.tb01879.x>.
14. Meneses-Gaya IC, Zuardi AW, Loureiro SR, Crippa JA. Psychometric properties of the Fagerstrom Test for Nicotine Dependence. *J Bras Pneumol.* 2009;**35**(1):73-82. [PubMed ID: 19219334]. <https://doi.org/10.1590/s1806-37132009000100011>.
15. Alotaibi SA, Alsuliman MA, Durgampudi PK. Smoking tobacco prevalence among college students in the Kingdom of Saudi Arabia: Systematic review and meta-analysis. *Tob Induc Dis.* 2019;**17**:35. [PubMed ID: 31516478]. [PubMed Central ID: PMC6662783]. <https://doi.org/10.18332/tid/105843>.
16. Al Riyami AA, Afifi M. Smoking in Oman: prevalence and characteristics of smokers. *East Mediterr Health J.* 2004;**10**(4-5):600-9. [PubMed ID: 16335652].
17. Ansari K, Farooqi FA. Comparison and prevalence of smoking among Saudi females from different Departments of the College of Applied Medical Sciences in Dammam. *Int J Health Sci (Qassim).* 2017;**11**(5):56-62. [PubMed ID: 29114195]. [PubMed Central ID: PMC5669512].
18. Koura MR, Al-Dossary AF, Bahnassy AA. Smoking pattern among female college students in Dammam, Saudi Arabia. *J Family Community Med.* 2011;**18**(2):63-8. [PubMed ID: 21897913]. [PubMed Central ID: PMC3159230]. <https://doi.org/10.4103/2230-8229.83370>.
19. Al-Mohamed HI, Amin TT. Pattern and prevalence of smoking among students at King Faisal University, Al Hassa, Saudi Arabia. *East Mediterr Health J.* 2010;**16**(1):56-64. [PubMed ID: 20214159].
20. Urberg KA, Luo Q, Pilgrim C, Degirmencioglu SM. A two-stage model of peer influence in adolescent substance use: individual and relationship-specific differences in susceptibility to influence. *Addict Behav.* 2003;**28**(7):1243-56. [PubMed ID: 12915166]. [https://doi.org/10.1016/s0306-4603\(02\)00256-3](https://doi.org/10.1016/s0306-4603(02)00256-3).