Published Online: 2025 January 5

Research Article



Exploring the Relationship Between Obsessive-Compulsive Disorder and Chronic Prostatitis/Chronic Pelvic Pain Syndrome: A Case-Control Study

Ayoub Salehi Mishani (1)¹, Seyed Shahab Banihashem (1)², Amir Alinejad Khorram (1)³, Shadi Kordbacheh (1)², Farzad Allameh (1)^{4,*}

¹ Department of Urology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

² Department of Psychosomatic Medicine, Taleghani Hospital Research Development Committee, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³ Clinical Research Development Unit, Shohada-e Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴ Men's Health and Reproductive Health Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

* Corresponding Author: Department of Urology, Shohada-e Tajrish Hospital, Tehran, Iran. Email: farzadallame@gmail.com

Received: 29 July, 2024; Revised: 15 November, 2024; Accepted: 16 December, 2024

Abstract

Background: Chronic prostatitis/chronic pelvic pain syndrome (CP/CPPS) is recognized as a challenging condition with unclear underlying factors. Despite the difficulties in treatment, psychological factors are thought to contribute to its symptoms.

Objectives: This investigation sought to examine the correlation between obsessive-compulsive disorder (OCD) and CP/CPPS.

Methods: We assembled a case group comprising 30 patients diagnosed with CP/CPPS and a control group consisting of 30 healthy individuals without the condition. The Yale-Brown Obsessive-Compulsive Scale (Y-BOCS) was used to assess the prevalence of OCD in both groups. In contrast, the National Institutes of Health (NIH) Chronic Prostatitis Symptom Index (NIH-CPSI) was used to evaluate the severity of CP/CPPS symptoms.

Results: The findings revealed no statistically significant difference in the prevalence of OCD between the group of individuals with CP/CPPS and the healthy control group. However, patients with CP/CPPS had significantly higher Y-BOCS scores compared to the control group. Furthermore, we observed a significant positive correlation between Y-BOCS scores and NIH-CPSI scores, suggesting that increasing OCD symptoms were associated with increased disease severity, pain, and decreased quality of life. Multivariate analysis revealed that increasing Y-BOCS scores independently predicted CP/CPPS symptom severity.

Conclusions: Our findings highlight the importance of considering psychological factors, including OCD, in the management of CP/CPPS. The results suggest that addressing OCD symptoms may be an important adjunctive treatment approach for improving symptom outcomes in patients with CP/CPPS. Future studies should investigate the effectiveness of psychotherapeutic interventions targeting OCD in this population.

Keywords: Obsessive-Compulsive Disorder, Prostatitis, Psychosomatic Disorders

1. Background

Chronic prostatitis/chronic pelvic pain syndrome (CP/CPPS) is a complex and debilitating condition characterized by a range of symptoms. According to the National Institutes of Health (NIH) guidelines, CP/CPPS is classified as a distinct subgroup of prostatitis, with two categories: IIIA and IIIB. This syndrome is defined by the presence of pain in the perineum, pelvis, suprapubic

region, or external genitalia, accompanied by varying degrees of urination and/or ejaculation disorders, without evidence of bacterial infection (1).

Chronic prostatitis/chronic pelvic pain syndrome is a significant and complex health issue, affecting approximately 2 - 10% of men worldwide. This condition significantly impairs patients' quality of life due to the chronic nature of its symptoms. Despite significant progress in basic and clinical research, the underlying

Copyright © 2025, Salehi Mishani et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

causes and mechanisms of CP/CPPS remain poorly understood, underscoring the need for further investigation (2, 3). Some studies have suggested that underlying psychiatric factors may contribute to the development of CP/CPPS, with some research even

Despite the availability of symptom definitions, a good classification system, and a standardized measurement instrument [National Institutes of Health (NIH) Chronic Prostatitis Symptom Index (NIH-CPSI)] (7), research on the role of psychosocial factors in CP/CPPS lags behind that of other chronic pain conditions. Due to the diverse and complex nature of symptoms in CP/CPPS, it is essential to gain a more detailed understanding of the psychological and social factors underlying the condition to develop targeted and effective treatment strategies for patients with CP/CPPS (8).

classifying it as a psychosomatic disorder (4-6).

Obsessive-compulsive disorder (OCD) is a psychiatric disorder that affects around 1% to 3% of the population, causing significant impairment and distress (9). This disorder is characterized by obsessions and compulsions that consume considerable time and lead to marked discomfort and functional impairment in patients (10). Some previous studies have suggested a relationship between OCD and chronic urological diseases, such as overactive bladder syndrome and CP/CPPS (4, 11).

Recent investigations into the psychiatric factors associated with CP/CPPS have emerged; however, there remains a critical knowledge gap regarding the underlying causes of this condition and the relationship between psychiatric factors and the development and exacerbation of its symptoms (8, 12, 13).

2. Objectives

The primary objective of the present study is to explore the potential correlation between OCD and CP/CPPS. Specifically, we will examine the co-occurrence of OCD and CP/CPPS, as well as the influence of OCD on symptom severity.

3. Methods

3.1. Study Population

This study recruited patients from the Urology Clinic of Shohada-e-Tajrish Referral Hospital between April 2023 and April 2024. The case group consisted of patients diagnosed with CP/CPPS, while the control group consisted of healthy individuals with no history of urological disease or pelvic pain. The inclusion criteria were as follows.

3.1.1. Case Group

- Confirmation of CP/CPPS diagnosis by a urologist not affiliated with the research team.

- Presence of prostatitis symptoms for at least 6 months.

- Absence of genitourinary infection, as confirmed by urine analysis (U/A), urinalysis (U/C), and the 2-glass test.

3.1.2. Control Group

- Healthy individuals with no history of urinary irritation symptoms or vague pelvic pain

The exclusion criteria are as follows:

- Previous history of genitourinary infection
- Malignancies
- Recurrent urinary stones
- Inflammatory bowel disease
- Benign prostatic hyperplasia
- Psychotic disorder
- Neurological diseases

Patients were excluded from the study if a relevant specialist confirmed any of the above conditions. Ultimately, 60 participants were recruited, with 30 individuals in the case group and 30 in the control group.

3.2. Study Design

This case-control study aimed to explore the association between OCD and CP/CPPS. The study comprised a case group of 30 individuals with CP/CPPS and a control group of 30 healthy individuals. We investigated the prevalence of OCD in both groups and compared the results. Additionally, we examined the relationship between OCD and symptom severity in patients with CP/CPPS.

3.3. Assessment Tools

The prevalence of OCD in both groups was assessed by a psychiatrist using the Yale-Brown Obsessive-Compulsive Scale (Y-BOCS) (14). The Y-BOCS scores range from 0 to 40, with a cut-off point of 17 or higher indicating a diagnosis of OCD.

The severity of CP/CPPS symptoms in the case group was evaluated by a urologist using the NIH-CPSI questionnaire (15). This scale assesses three domains: Pain (0 - 21), urinary symptoms (0 - 10), and quality of life (0 - 12), with a total score range of 0 to 43.

Both questionnaires have been validated for use in Iranian populations (16, 17).

3.4. Outcome Assessment

The primary outcome is to investigate the prevalence of OCD in individuals with CP/CPPS compared to healthy individuals. The secondary outcome is to examine the association between OCD and the severity of CP/CPPS symptoms.

3.5. Statistical Methods

Data analysis was conducted using SPSS software version 22. The results are presented as means and standard deviations for quantitative variables, and frequencies and percentages for qualitative variables.

For the analysis of quantitative data, Pearson's correlation test was employed to examine the relationships between variables. To compare the groups in terms of quantitative variables, the normality of the data distribution was assessed using the Shapiro-Wilk test. Then, either the Mann-Whitney U test or the *t*-test was used to compare the two groups. For comparing qualitative variables, Fisher's exact test or the chi-square test was used. Multivariate linear regression analysis was performed to calculate odds ratios (OR) and identify potential predictors.

In all statistical analyses, a P-value of 0.05 or lower was considered statistically significant.

3.6. Ethical Statement

This study did not incur any costs for participants, and the researchers did not influence the patients' decision-making or treatment processes in any way. Each participant provided informed consent before participating in the study.

This research was approved by the Ethics Committee of Shahid Beheshti University of Medical Sciences, with approval number IR.SBMU.MSP.REC.1401.481.

4. Results

The study included 60 participants: Thirty controls and 30 individuals with CP/CPPS. The mean age of the participants was 42.83 ± 14.38 years (range: 18 - 80 years). A comparison of demographic variables between the case and control groups showed no statistically significant differences, as presented in Table 1.

In this study, 12 (20%) of the 60 individuals examined met the criteria for OCD based on a Y-BOCS cut-off score of 17. Of these, 9 were from the CP/CPPS group and 3 were from the healthy individuals group. Although there was no statistically significant difference in the prevalence of OCD between the case and control groups (P = 0.10), the average Y-BOCS scores were significantly higher in the CP/CPPS group (12.83 \pm 7.02) compared to the control group (6.1 \pm 8.4) (P = 0.001), as shown in Figure 1.

To examine the relationship between OCD symptoms and the severity of CP/CPPS symptoms, we analyzed the correlation between Y-BOCS scores and scores from various components of the NIH-CPSI questionnaire, including pain, urinary symptoms, and quality of life. A significant correlation was found between Y-BOCS scores and pain scores, quality of life scores, and total NIH-CPSI scores in patients with CP/CPPS (Table 2). Furthermore, the multivariate regression test results revealed that Y-BOCS scores were an independent predictor of symptom severity, even after controlling for other potential risk factors (Table 3).

5. Discussion

Our findings revealed a statistically significant difference between the patients with CP/CPPS and the healthy individuals in terms of Y-BOCS scores, with the patient group exhibiting higher scores. However, this difference did not translate into a significant difference in the prevalence of OCD itself.

A striking finding of this study was the direct and significant correlation between Y-BOCS scores, pain scores, quality of life scores, and total scores of NIH-CPSI. Specifically, patients with CP/CPPS who had higher Y-BOCS scores experienced more pain and lower quality of life. Furthermore, our analysis revealed that OCD, as evaluated by Y-BOCS scores, was an independent predictor of the severity of CP/CPPS symptoms. These findings suggest that addressing OCD may be an important component of comprehensive treatment plans for individuals with CP/CPPS.

The CP/CPPS is a chronic condition marked by debilitating pelvic pain and discomfort, which can be notoriously challenging to treat (18). Despite its prevalence, the underlying causes of CP/CPPS remain debated among healthcare professionals. Given the significant impact this disease has on patients' quality of life and the substantial costs it incurs for healthcare systems, it is essential to investigate the underlying factors that contribute to this complex condition (3, 19).

The heterogeneity of CP/CPPS underscores the need for a more comprehensive understanding of the psychological mechanisms underlying this condition. Studies indicate that psychological factors have a substantial impact on the development and worsening of CP/CPPS symptoms, a notion that has been supported by studies for decades (8). For instance, Piontek et al.

Variables	Case (N = 30)	Control(N=30)	P-Value
Age (y)	43.43±14.36	42.25 ± 14.61	0.75
BMI (kg/m ²)	24.76 ± 4.26	26.56 ± 4.97	0.13
Duration of chronic prostatitis (mon)	37.00 ± 37.29	-	-
Marital status (married)	21(70)	21 (70)	0.99
Smoking history	10 (33.3)	9 (30)	0.78
Diabetes mellitus	2 (6.7)	1(3.3)	0.99

Abbreviation: BMI, Body Mass Index.

^a Values are expressed as mean \pm SD or No. (%).



Figure 1. Comparison of Yale-Brown Obsessive-Compulsive Scale (Y-BOCS) scores between case and control groups

Table 2. Correlation Between Yale-Brown Obsessive-Compulsive Scale Scores and National Institutes of Health Chronic Prostatitis Symptom Index				
Variables	r Coefficient	P-Value		
Y-BOCS scores/pain scores of NIH-CPSI	0.4	0.02		
Y-BOCS scores/urinary symptom scores of NIH-CPSI	-0.27	0.15		
Y-BOCS scores/quality of life scores of NIH-CPSI	0.37	0.04		
Y-BOCS scores/total scores of NIH-CPSI	0.35	0.05		

Abbreviations: Y-BOCS, Yale-Brown Obsessive-Compulsive Scale; NIH-CPSI, National Institutes of Health Chronic Prostatitis Symptom Index.

identified behavioral patterns associated with depression in patients with CP/CPPS, particularly during symptom flare-ups (20). Furthermore, Naliboff et al. reported higher levels of lifetime stress in this patient population, along with increased self-reported cognitive deficits compared to individuals in the general population (21). In light of the accumulating evidence regarding the relationship between psychological factors and CP/CPPS symptoms, the application of psychiatric approaches in treating these patients has been increasingly explored and developed (22, 23). A randomized controlled trial by Wang et al. found that

Variables		Multivariate Regression			
	SE	P-Value	Odds Ratio		
Age	0.11	0.58	0.54		
BMI	0.38	0.34	0.96		
Smoking history	3.38	0.98	0.02		
Diabetes mellitus	6.71	0.66	0.44		
Y-BOCS scores	0.24	0.04	2.11		

Table 3. Association Between Study Variables and National Institutes of Health Chronic Prostatitis Symptom Index Scores in Patients with Chronic Prostatitis/Chronic Pelvic Pain Syndrome

Abbreviations: BMI, Body Mass Index; Y-BOCS, Yale-Brown Obsessive-Compulsive Scale.

psychological interventions significantly improved treatment outcomes for patients with CP/CPPS (22). Despite these advancements, there remains a critical need for deeper insights into the psychological factors contributing to this condition to further enhance the efficacy of treatment strategies.

Obsessive-compulsive disorder is a common psychiatric condition that can manifest with various urological symptoms, including sexual dysfunction or vague urinary-genital infections. Accurate diagnosis of OCD in these cases is crucial to avoid misdiagnosis, which can lead to associated risks to patients, unnecessary physical examinations, and treatment of the wrong disease. Therefore, urologists should consider the possibility of OCD in patients presenting with these symptoms, particularly when the underlying cause is not immediately apparent (24).

A study by Tournikioti et al. evaluated 57 patients with chronic prostatitis, assessing symptoms and OCD using the Leyton Obsessional Inventory (LOI). The results revealed that 58% of patients exhibited abnormal LOI scores, indicating the presence of obsessivecompulsive characteristics, a remarkably high percentage (4).

Some research suggests that cellular and molecular changes related to psychological problems may be primarily responsible for the development of CP/CPPS in some patients (25). For example, studies have shown that individuals with depression tend to have lower levels of interleukin-10 in their peripheral mononuclear cells compared to healthy controls. Additionally, in patients with stage III prostate disease, the level of concern is associated with lower levels of interleukin-6 and 10 in the seminal plasma (26).

A growing body of evidence suggests that urological chronic pain diseases frequently co-occur with psychiatric disorders (27, 28). For instance, Weissman et al.'s study found that patients with interstitial cystitis are at a higher risk of developing panic disorder over their lifetime (29). Similarly, Allameh et al.'s study revealed that Y-BOCS scores were significantly higher in patients with Overactive Bladder Syndrome compared to a control group (11). Our study's results also indicate that, although OCD in patients with prostatitis was not significantly higher than in the control group, these patients had a significantly higher Y-BOCS score than the control group. Furthermore, a significant change in pain intensity and quality of life was observed as this score increased.

The interpretation of these studies' results suggests that an underlying psychiatric issue may be a key contributor to the onset and progression of chronic urological diseases. Moreover, these findings confirm previous studies' results, emphasizing the importance of psychiatric treatments in achieving more effective treatment outcomes for this patient population.

5.1. Limitations of the Study

The main limitation of the present study is its small sample size. As a result, the findings may not be generalizable to all patients with CP/CPPS. Additionally, the limited sample size may have reduced the power to detect statistically significant differences between groups. Future studies with larger, multi-center cohorts covering diverse racial and age groups can help address these limitations and provide more accurate and reliable insights into the relationship between psychiatric treatments and chronic urological diseases. Ultimately, such studies could inform more effective clinical practices and treatment decisions.

5.2. Conclusions

The present study underscores the significance of psychological factors in CP/CPPS, emphasizing the need for a more comprehensive understanding of this complex condition. Our findings suggest that brief screening assessments can facilitate timely diagnosis and referral, which are critical for effective management. Furthermore, we highlight the importance of future research to thoroughly explore psychosocial variables and evaluate the efficacy of psychotherapeutic interventions in patients with CP/CPPS.

Footnotes

Authors' Contribution: Study concept and design: F. A.; Acquisition of data: A. S. and S. K.; Analysis and interpretation of data: A. A.; Drafting of the manuscript: S. B. and A. S.; Critical revision of the manuscript for important intellectual content: F. A.; Study supervision: S. B. and F. A.

Conflict of Interests Statement: The authors declared no conflict of interests.

Data Availability: After requesting from corresponding author the raw data was available.

Ethical Approval: IR.SBMU.MSP.REC.1401.481.

Funding/Support: This study was not supported in part or complete from the any national health agency or any research center.

Informed Consent: The informed consent was obtained from each participant.

References

- Maeda K, Shigemura K, Fujisawa M. A review of current treatments for chronic prostatitis/chronic pelvic pain syndrome under the UPOINTS system. Int J Urol. 2023;30(5):431-6. [PubMed ID: 36788717]. https://doi.org/10.1111/jju.15149.
- Lee JH, Yoo TK, Kang JY, Cho JM, Park YW, Lee SW, et al. Relationship between erectile dysfunction and moderate to severe prostatitis-like symptoms in middle-aged men: a propensity score-matched analysis. *Int Urol Nephrol.* 2021;**53**(11):2261-6. [PubMed ID: 34546557]. https://doi.org/10.1007/s11255-021-02991-8.
- Sugimoto M, Hijikata Y, Tohi Y, Kuroda H, Takei M, Matsuki T, et al. Low quality of life in men with chronic prostatitis-like symptoms. *Prostate Cancer Prostatic Dis.* 2022;25(4):785-90. [PubMed ID: 35752656]. [PubMed Central ID: PMC9705241]. https://doi.org/10.1038/s41391-022-00559-w.
- Tournikioti K, Moussas G, Korkoliakou P, Christodoulou C, Gourgoulis GM, Tsiodras S, et al. P01-311 - Alexithymia, Hypochondriasis and Obsessive-compulsive Symptoms in Patients with Chronic Prostatitis. *European Psychiatry*. 2020;25(S1). https://doi.org/10.1016/s0924-9338(10)70519-8.
- Keltikangas-Jarvinen L, Ruokolainen J, Lehtonen T. Personality pathology underlying chronic prostatitis. *Psychother Psychosom*. 1982;37(2):87-95. [PubMed ID: 7122791]. https://doi.org/10.1159/000287558.
- 6. Stamatiou K, Trinchieri M, Trinchieri M, Perletti G, Magri V. Chronic prostatitis and related psychological problems. Which came first:

The chicken or the egg? A systematic review. Arch Ital Urol Androl. 2023;95(1). https://doi.org/10.4081/aiua.2023.11300.

- Dun RL, Tsai J, Hu XH, Mao JM, Zhu WJ, Qi GC, et al. A systematic review of cross-cultural adaptation of the National Institutes of Health Chronic Prostatitis Symptom Index. *Health Qual Life Outcomes*. 2021;**19**(1):159. [PubMed ID: 34059073]. [PubMed Central ID: PMC8166010]. https://doi.org/10.1186/s12955-021-01796-8.
- Riegel B, Bruenahl CA, Ahyai S, Bingel U, Fisch M, Lowe B. Assessing psychological factors, social aspects and psychiatric co-morbidity associated with Chronic Prostatitis/Chronic Pelvic Pain Syndrome (CP/CPPS) in men – a systematic review. J Psychosom Res. 2014;77(5):333-50. [PubMed ID: 25300538]. https://doi.org/10.1016/j.jpsychores.2014.09.012.
- Zai G, Barta C, Cath D, Eapen V, Geller D, Grunblatt E. New insights and perspectives on the genetics of obsessive-compulsive disorder. *Psychiatr Genet.* 2019;29(5):142-51. [PubMed ID: 31464995]. https://doi.org/10.1097/YPG.00000000000230.
- Cervin M. Obsessive-Compulsive Disorder: Diagnosis, Clinical Features, Nosology, and Epidemiology. Psychiatr Clin North Am. 2023;46(1):1-16. [PubMed ID: 36740346]. https://doi.org/10.1016/j.psc.2022.10.006.
- Allameh F, Motamed M, Poury M, Tayyebiazar A, Ghiasy S, Fallah-Karkan M, et al. Relationship between Obsessive-compulsive Disorder and Overactive Bladder Syndrome. *Iran J Psychiatry Behav Sci.* 2021;**15**(1). https://doi.org/10.5812/ijpbs.101858.
- Li AS, Van Niekerk L, Wong ALY, Matthewson M, Garry M. Psychological management of patients with chronic prostatitis/chronic pelvic pain syndrome (CP/CPPS): a systematic review. Scand J Pain. 2023;23(1):25-39. [PubMed ID: 35938980]. https://doi.org/10.1515/sjpain-2022-0049.
- Huang X, Qin Z, Cui H, Chen J, Liu T, Zhu Y, et al. Psychological factors and pain catastrophizing in men with chronic prostatitis/chronic pelvic pain syndrome (CP/CPPS): a meta-analysis. *Transl Androl Urol.* 2020;9(2):485-93. [PubMed ID: 32420154]. [PubMed Central ID: PMC7214995]. https://doi.org/10.21037/tau.2020.01.25.
- Goodman WK, Price LH, Rasmussen SA, Mazure C, Fleischmann RL, Hill CL, et al. The Yale-Brown Obsessive Compulsive Scale. I. Development, use, and reliability. Arch Gen Psychiatry. 1989;46(11):1006-11. [PubMed ID: 2684084]. https://doi.org/10.1001/archpsyc.1989.01810110048007.
- Litwin MS, McNaughton-Collins M, Fowler FJ, Nickel JC, Calhoun EA, Pontari MA, et al. The National Institutes of Health chronic prostatitis symptom index: development and validation of a new outcome measure. Chronic Prostatitis Collaborative Research Network. J Urol. 1999;162(2):369-75. [PubMed ID: 10411041]. https://doi.org/10.1016/s0022-5347(05)68562-x.
- Allameh F, Mansouri Tehrani MM, Tasharrofi MA, Ganji Jameshouran MA. Validation of the Persian Version of the National Institute of Health Chronic Prostatitis Symptom Index. Urol J. 2020;18(1):117-21. [PubMed ID: 32149374]. https://doi.org/10.22037/uj.v0i0.5444.
- 17. Esfahani SR, Motaghipour Y, Kamkari K, Zahiredin A, Janbozorgi M. Reliability and validity of the Persian version of the Yale-Brown Obsessive-Compulsive Scale (Y-BOCS). *Iran J Psychiatry Behav Sci.* 2012;**17**(4):297-303.
- Pena VN, Engel N, Gabrielson AT, Rabinowitz MJ, Herati AS. Diagnostic and Management Strategies for Patients with Chronic Prostatitis and Chronic Pelvic Pain Syndrome. *Drugs Aging*. 2021;38(10):845-86. [PubMed ID: 34586623]. https://doi.org/10.1007/s40266-021-00890-2.
- Clemens JQ, Markossian T, Calhoun EA. Comparison of economic impact of chronic prostatitis/chronic pelvic pain syndrome and interstitial cystitis/painful bladder syndrome. *Urology*. 2009;**73**(4):743-6. [PubMed ID: 19193408]. [PubMed Central ID: PMC2684702]. https://doi.org/10.1016/j.urology.2008.11.007.

- 20. Piontek K, Ketels G, Albrecht R, Schnurr U, Dybowski C, Brunahl CA, et al. Somatic and psychosocial determinants of symptom severity and quality of life in male and female patients with chronic pelvic pain syndrome. *J Psychosom Res.* 2019;**120**:1-7. [PubMed ID: 30929698]. https://doi.org/10.1016/j.jpsychores.2019.02.010.
- Naliboff BD, Stephens AJ, Afari N, Lai H, Krieger JN, Hong B, et al. Widespread Psychosocial Difficulties in Men and Women With Urologic Chronic Pelvic Pain Syndromes: Case-control Findings From the Multidisciplinary Approach to the Study of Chronic Pelvic Pain Research Network. Urology. 2015;85(6):1319-27. [PubMed ID: 26099876]. [PubMed Central ID: PMC4479402]. https://doi.org/10.1016/j.urology.2015.02.047.
- 22. Wang J, Liang K, Sun H, Li L, Wang H, Cao J. Psychotherapy combined with drug therapy in patients with category III chronic prostatitis/chronic pelvic pain syndrome: A randomized controlled trial. *Int J Urol.* 2018;**25**(8):710-5. [PubMed ID: 29862568]. https://doi.org/10.1111/iju.13706.
- Lackner JM, Clemens JQ, Radziwon C, Danforth TL, Ablove TS, Krasner SS, et al. Cognitive Behavioral Therapy for Chronic Pelvic Pain: What Is It and Does It Work? J Urol. 2024;211(4):539-50. [PubMed ID: 38228093]. [PubMed Central ID: PMC10939861]. https://doi.org/10.1097/JU.000000000003847.
- 24. Conti D, Pellegrini L, Riaz M, Mpavaenda D, Sim A, Fineberg N. Obsessive-compulsive disorder and men's health. Part 1: Recognition

and diagnosis. *Trend Urol Mens Heal*. 2023;**14**(1):11-6. https://doi.org/10.1002/tre.895.

- Pontari MA, Ruggieri MR. Mechanisms in prostatitis/chronic pelvic pain syndrome. J Urol. 2008;179(5 Suppl):S61-7. [PubMed ID: 18405756]. https://doi.org/10.1016/j.juro.2008.03.139.
- Miller LJ, Fischer KA, Goralnick SJ, Litt M, Burleson JA, Albertsen P, et al. Nerve growth factor and chronic prostatitis/chronic pelvic pain syndrome. Urology. 2002;59(4):603-8. [PubMed ID: 11927336]. https://doi.org/10.1016/s0090-4295(01)01597-7.
- Mazza M, Margoni S, Mandracchia G, Donofrio G, Fischetti A, Kotzalidis GD, et al. This pain drives me crazy: Psychiatric symptoms in women with interstitial cystitis/bladder pain syndrome. World J Psychiatry. 2024;14(6):954-84. [PubMed ID: 38984334]. [PubMed Central ID: PMC11230088]. https://doi.org/10.5498/wjp.v14.i6.954.
- Mahjani B, Koskela LR, Batuure A, Gustavsson Mahjani C, Janecka M, Hultman CM, et al. Systematic review and meta-analysis identify significant relationships between clinical anxiety and lower urinary tract symptoms. *Brain Behav.* 2021;11(9). e2268. [PubMed ID: 34402598]. [PubMed Central ID: PMC8442597]. https://doi.org/10.1002/brb3.2268.
- Weissman MM, Gross R, Fyer A, Heiman GA, Gameroff MJ, Hodge SE, et al. Interstitial cystitis and panic disorder: a potential genetic syndrome. Arch Gen Psychiatry. 2004;61(3):273-9. [PubMed ID: 14993115]. https://doi.org/10.1001/archpsyc.61.3.273.