



Key Factors for Predicting Appendicitis from Contrast-Enhanced Computed Tomography Images Obtained Through Multiplanar Reconstruction

Jiun-Ying Chern¹, Yon Cheong Wong^{1,2,*}, Li Jen Wang^{1,2}, Cheng Hsien Wu¹, Huan Wu Chen¹, Being Chuan Lin³

¹ Division of Emergency and Critical Care Radiology, Department of Medical Imaging & Intervention, Chang Gung Memorial Hospital, Taoyuan, Taiwan

² College of Medicine, Chang Gung University, Taoyuan 330, Taiwan

³ Division of Trauma & Emergency Surgery, Department of Surgery, Chang Gung Memorial Hospital, Taoyuan 333, Taiwan

*Corresponding Author: Division of Emergency and Critical Care Radiology, Department of Medical Imaging & Intervention, Chang Gung Memorial Hospital, Taoyuan, Taiwan. Email: ycwong@cgmh.org.tw

Received: 3 March, 2024; Revised: 17 September, 2024; Accepted: 23 September, 2024

Abstract

Background: The diameter of the appendix is a key parameter in diagnosing appendicitis. The diagnostic threshold for this parameter is 6 mm, originally established through graded compression sonography of the right lower quadrant (RLQ) of the abdomen. However, without corroborative findings from computed tomography (CT), this threshold may not be a reliable indicator of appendicitis. To ensure accurate diagnosis, clinicians should perform a comprehensive, multiparameter imaging assessment of the appendix, rather than relying solely on appendix diameter.

Objectives: This study aimed to identify key factors for predicting appendicitis using contrast-enhanced coronal and sagittal CT images obtained through multiplanar reconstruction.

Patients and Methods: This single-center, retrospective, cross-sectional study included patients who presented to our emergency department (ED) with RLQ abdominal pain and subsequently underwent contrast-enhanced CT between July 2019 and September 2020. The primary study outcome was pathologically confirmed appendicitis. Two experienced radiologists assessed parameters such as appendix diameter, wall thickness, abnormal appendix enhancement, abnormal appendix content, appendix erection, and periappendiceal fat stranding. Multivariate logistic regression was performed to identify significant predictive factors for appendicitis.

Results: The study included 173 patients (median age: 37 years; women: 86). They were divided into appendicitis ($n = 102$) and alternative diagnosis ($n = 71$) groups. Significant differences were observed between the groups in terms of appendix diameter, wall thickness, wall enhancement, luminal content, appendix erection, and periappendiceal fat stranding ($P < 0.001$). The diagnostic sensitivity and specificity values for an appendix diameter threshold of 7.7 mm were 91% and 82%, respectively. An appendix diameter of > 7.7 mm (OR: 15.3; $P < 0.001$), abnormal appendix enhancement (OR: 12.5; $P < 0.001$), and appendix erection (OR: 6.1; $P = 0.004$) emerged as significant independent predictors of appendicitis.

Conclusion: An appendix diameter of 7.7 mm appears to be the optimal threshold for diagnosing appendicitis. Additionally, the detection of abnormal appendix enhancement and appendix erection on contrast-enhanced CT images holds considerable diagnostic value.

Keywords: Appendicitis, Appendix, Abdominal Pain, Multidetector Computed Tomography, Diagnostic Imaging

1. Background

Acute appendicitis, with an overall global incidence of approximately 100 cases per 100,000 adults, is the most common reason for emergency surgery (1, 2).

Although the clinical diagnosis of appendicitis is typically straightforward in patients presenting with classic signs and symptoms, various conditions (e.g., gynecologic, renal, and colonic diseases) may complicate the diagnosis, leading to treatment delays or

unnecessary procedures (3-5). Over the last decade, computed tomography (CT) has been increasingly used to diagnose appendicitis due to its widespread availability, standardized technique, rapid data acquisition, and potential for differential diagnosis (6). Clinicians and radiologists primarily use the appendix diameter as a key parameter for assessing appendicitis, with a threshold of 6 mm. This value was originally derived from graded compression sonography of the right lower quadrant (RLQ) of the abdomen; however, CT images are acquired without compression (7-9). Evidence suggests that over 20% of patients without appendicitis have an appendix diameter greater than 7 mm, making the 6 mm threshold unreliable in the absence of corroborative CT findings (9, 10). Therefore, a comprehensive, multiparameter imaging evaluation of the appendix is essential for accurate diagnosis. This study aimed to identify key factors for predicting appendicitis from contrast-enhanced CT images obtained through multiplanar reconstruction.

2. Objectives

The main objective of this study was to identify key factors for predicting appendicitis from contrast-enhanced coronal and sagittal CT images obtained through multiplanar reconstruction.

3. Patients and Methods

3.1. Study Cohort

This retrospective cross-sectional study was approved by the institutional review board of our hospital. The board waived the requirement for informed consent due to the retrospective nature of the study and the use of deidentified patient data. Between July 2019 and September 2020, 185 consecutive patients presented to our emergency department (ED) with RLQ pain and underwent contrast-enhanced CT. For patients diagnosed with appendicitis, pathology confirmation after appendectomy was required. Patients who presented with typical imaging features of appendicitis but were treated with antibiotic therapy were excluded. Those who had a negative appendectomy or RLQ pain but were diagnosed with a different condition were included in the alternative diagnosis group. Medical records were monitored for 1 month to confirm the absence of any ED or outpatient department revisits, ensuring there was no misdiagnosis in the alternative diagnosis group. Patients who were discharged against medical advice or had a history of appendectomy were excluded from the study.

3.2. Imaging Modality

Computed tomography imaging was performed using the Aquilion 64 scanner (Toshiba Medical Systems, Otawara, Japan). No enteric contrast medium was administered; however, 100 mL of an intravenous contrast medium (iohexol; 350 mg iodine/mL; Omnipaque 350; GE Healthcare, Princeton, NJ, USA) was administered at a rate of 2 mL/s. Portal venous phase imaging was conducted 70 seconds after the contrast medium injection. The CT parameters were as follows: 120 kVp with automatic tube current modulation. For analysis, all images were reconstructed with a 5 mm thickness at 5 mm intervals, and coronal and sagittal images were obtained through multiplanar reconstruction.

3.3. Data Collection and Analysis

All data were reviewed by two abdominal radiologists (JYC and YCW, with 5 and 25 years of experience, respectively) who were blinded to the diagnosis. Patient identifiers, including numbers and names, were removed and replaced with random numbers serving as virtual identifiers. Only members of the research team had access to patient data and images. A standardized questionnaire was used to assess imaging parameters such as appendix diameter, wall thickness, abnormal appendix enhancement, abnormal appendix content, appendix erection, and periappendiceal fat stranding, without revealing the CT scan diagnosis. The final values for appendix diameter and wall thickness were the averages of the two radiologists' measurements. For the other parameters, the final values were determined based on consensus between the radiologists (in cases of discrepancies).

Abnormal appendix enhancement was defined as a difference in enhancement between the appendix and the cecal or ileal wall. Abnormal appendix content was defined as the presence of intraluminal fluid (Figure 1) or appendicoliths, rather than intraluminal air, within the appendix. Appendix erection was defined as the presence of an I-shaped or C-shaped appendix rather than a V-shaped, O-shaped, or S-shaped flaccid appendix on axial, sagittal, and coronal CT images. An I-shaped appendix was characterized by a straight or nearly straight appearance, while a C-shaped appendix was characterized by a curved, bow-like appearance (Figure 1). A V-shaped appendix had an acute angle at its turn, an O-shaped appendix appeared with the tip close to the opening, resembling a "kissing" posture, and an S-shaped appendix had a biconcave appearance (Figure 2).




Figure 1. Image appearance of an erected appendix: A and B, a 21-year-old male patient with appendicitis. Post contrast computed tomography (CT) on axial view showed an erected “I shape” appendix (red arrows). The appendix diameter was 6.6 mm. Note also the appendix had abnormal wall enhancement and filled with fluid; C and D, a 62-year-old male patient with appendicitis. Post contrast CT on coronal view showed an erected “C shape” appendix (red arrows) with abnormal wall enhancement and intraluminal fluid. Appendix diameter was 8 mm.

3.4. Statistical Analysis

All data were analyzed using SPSS (IBM Corp. Released 2013. IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp.). Intraclass correlation coefficients were used to assess interobserver agreement for appendix diameter and wall thickness. Cohen’s kappa coefficient was used to evaluate interobserver agreement regarding abnormal appendix

enhancement, abnormal appendix content, appendix erection, and periappendiceal fat stranding. Categorical variables were presented as frequencies and percentages, while continuous variables with a normal distribution were presented as mean and standard deviation (SD) values. Continuous variables with a non-normal distribution were presented as median and interquartile range (IQR) values.

Categorical variables were analyzed using the chi-square or Fisher exact test, and continuous variables




Figure 2. Image appearance of a flaccid appendix: A, B, and C, a 53-year-old male patient with right lower quadrant (RLQ) pain. Post contrast computed tomography (CT) on sagittal view showed a "V shape" appendix (red arrows) distinguished from a "C shape" appendix by an acute angle turn (yellow arrows in A and B). The appendix diameter was 7.8 mm. The patient was diagnosed with right ureterovesicle junction (UVJ) stone. D, E, and F, an "O shape" appendix (red arrows) with mesentery fat stranding in RLQ of abdomen on post contrast CT coronal view in a patient with RLQ pain. The appendix tip "kissing" the opening of the appendix (yellow arrow in E) with a diameter of 7 mm. The patient was diagnosed with ascending colon diverticulitis instead of appendicitis. G and H, a 42-years-old male patient visited our ED due to RLQ pain with final diagnosis of enteritis. Post contrast CT on coronal view showed a bi-curved "S shape" appendix (red arrows). The appendix diameter was 7.2 mm.

were analyzed using the independent-sample *t*-test or Mann-Whitney U test, as appropriate. A P-value of < 0.05 was considered statistically significant. The optimal thresholds for appendix diameter and wall thickness were determined using the maximum Youden index values, which were derived from the corresponding

receiver operating characteristic (ROC) curves. Variables with a P-value < 0.1 in the univariate analysis were included in a backward stepwise multivariate logistic regression model to identify significant independent factors for predicting appendicitis.

4. Results

4.1. Patient Characteristics

Between July 2019 and September 2020, 185 patients visited our hospital. Of these, 12 patients were excluded for the following reasons: Five patients presented with typical imaging features of appendicitis but received antibiotic treatment, 5 had a history of appendectomy, and 2 were discharged against medical advice. Ultimately, this study included 173 patients [median age: 37 years (IQR: 27 to 54)]. Of these, 102 received a pathologically confirmed diagnosis of appendicitis, and 71 were diagnosed with an alternative condition or had a negative appendectomy ($n = 6$). None of the patients with alternative diagnoses revisited our hospital's ED or outpatient department within 1 month of their initial presentation (Figure 3).




Figure 3. Flowchart of patients presented to emergency department (ED) with right lower quadrant (RLQ) pain between July 2019 to September 2020.

The median age of the appendicitis group was 43.4 years, which included 54 male patients (53%). The mean age of the alternative diagnosis group was 38.8 years, which included 33 male patients (46.5%). No significant differences were observed between the two groups in terms of age or sex. However, significant differences were noted in leukocytosis ($> 10,000/\text{mL}$; $P = 0.001$) and leukocyte left shift ($> 75\%$; $P < 0.001$), but not in body temperature ($P = 0.279$) or C-reactive protein level ($P = .196$).

4.2. Image Characteristics

The interobserver intraclass correlation coefficients for appendix diameter and wall thickness were 0.915 and 0.756, respectively. Cohen's kappa coefficients for abnormal appendix enhancement, abnormal appendix content, appendix erection, and periappendiceal fat

stranding were 0.766, 0.657, 0.833, and 0.835, respectively. The overall incidence of appendicitis among patients presenting to our ED with right RLQ pain was 59% (102/173).

The mean diameter of the appendix was significantly greater in the appendicitis group compared to the alternative diagnosis group (11.9 ± 3.5 mm vs 6.6 ± 1.4 mm, respectively; $P < 0.001$). Similarly, the median appendix wall thickness was significantly higher in the appendicitis group than in the alternative diagnosis group (3.1 [IQR: 2.3 to 3.8] mm vs 1.7 [IQR: 1.5 to 2.3] mm, respectively; $P < 0.001$). Univariate analysis identified the following significant ($P < 0.001$) predictors of appendicitis: Abnormal appendix enhancement, abnormal appendix content, appendix erection, and periappendiceal fat stranding (Table 1).

Table 1. Demographic, Laboratory and Imaging Findings of Patients with and without Appendicitis^a

Characteristics	Appendicitis (n = 102)	Alternative diagnosis (n = 71)	P-value
Age, median (IQR)	34 (27 - 49)	39 (29 - 56)	0.09
Gender, male	54/102 (52.9)	33/71 (46.5)	0.403
Body temperature ($^{\circ}\text{C}$)	36.7 (36.2 - 37.3)	36 (36.2 - 37.5)	0.279
Leukocytosis ($> 10000/\text{uL}$)	72/102 (70)	33/71 (46)	0.001
Leukocyte left shift ($> 75\%$)	80/102 (78)	32/71 (45)	< 0.001
C-reactive protein level (mg/L)	19.6 (3.0 - 67.3)	27.4 (6.1 - 122.8)	0.196
Surgical time (min)	142 (77 - 300)	90 (70 - 120)	0.03
Appendix diameter (mm)	11.9 ± 3.5	6.6 ± 1.4	< 0.001
Appendix wall thickness (mm)	1.7 (1.5 - 2.3)	3.1 (2.3 - 3.8)	< 0.001
Abnormal wall enhancement	89/102 (87)	10/71 (11)	< 0.001
Abnormal intraluminal content	96/102 (93)	29/71 (33)	< 0.001
Erection of appendix	89/102 (87)	13/71 (18)	< 0.001
Peri-appendiceal fat stranding	92/102 (90)	25/71 (35)	< 0.001
Appendix diameter ≥ 7.7 mm	93/102 (91.1)	10/71 (14)	< 0.001
Appendix wall thickness ≥ 2 mm	90/102 (88.2)	30/71 (42.2)	< 0.001

Abbreviations: SD, standard deviation; IQR, interquartile range.

^a Values are expressed as No. (%), mean \pm SD or median (IQR).

The area under the ROC curve values for appendix diameter and wall thickness were 0.940 and 0.855, respectively. The Youden index indicated that an appendix diameter threshold of 7.7 mm achieved sensitivity and specificity values of 91% and 82%, respectively. In contrast, an appendix diameter

threshold of 6 mm achieved sensitivity and specificity values of 99% and 36%, respectively. For appendix wall thickness, a threshold of 2 mm yielded sensitivity and specificity values of 86% and 65%, respectively (Table 2).

Table 2. Diagnostic Accuracy of Diameter and Thickness of The Appendix for diagnosis of Acute Appendicitis

Variables	AUC (95 % CI)	Sensitivity (%)	Specificity (%)	Cut off value (mm)
Appendix diameter	0.940	99.0	36.2	6
		91.2	85.9	7.7
Appendix thickness	0.855	86.3	64.8	2

Abbreviations: AUC, area under curve; CI, confidence interval.

4.3. Significant Predictive Factors for Appendicitis

After adjusting for covariates, the multivariate logistic regression identified the following independent predictors of appendicitis (Table 3): Appendix diameter greater than 7.7 mm [odds ratio (OR): 15.3; 95% confidence interval (CI): 4.8 to 47.6; $P < 0.001$], abnormal appendix enhancement (OR: 12.5; 95% CI: 3.7 to 41.7; $P < 0.001$), and appendix erection (OR: 6.1; 95% CI: 1.9 to 20.8; $P = 0.004$).

Table 3. Predictors of Appendicitis on Multivariable Logistic Regression Analysis

Variables	Odds ratio (95% CI)	P-value
Appendix diameter ≥ 7.5 mm	15.3 (4.8 - 47.6)	< 0.001
Abnormal appendix wall enhancement	12.5 (3.7 - 41.7)	< 0.001
Erection of appendix	6.1 (1.9 - 20.8)	0.004

Abbreviation: CI, confidence interval.

5. Discussion

Acute appendicitis is a common cause of abdominal pain, a leading reason for emergency abdominal surgery, and a frequent condition associated with lawsuits against emergency physicians or radiologists (8). Despite extensive discussion over the years and the use of clinical scoring systems, diagnosing acute appendicitis remains challenging. Approximately 60% of patients with uncomplicated appendicitis can be treated successfully with antibiotics (11, 12). However, appendectomy remains the definitive treatment for acute appendicitis as it provides a single-session resolution and allows for pathological confirmation when the diagnosis is uncertain (13). This underscores the importance of ensuring an accurate diagnosis to avoid unnecessary surgery.

With advancements in imaging technology, significant improvements have been made in CT image acquisition time, image resolution, and multiplanar reconstruction, enabling CT imaging to offer crucial insights for patients suspected of having appendicitis (14). Currently, intravenous administration of contrast media is recommended to enhance diagnostic accuracy, except in specific clinical scenarios (15, 16). As a result, imaging studies continue to play a vital role in preventing misdiagnosis (17, 18). Although several imaging parameters are used to diagnose appendicitis, the disease presentation on imaging is not always straightforward (19, 20).

Appendicitis is believed to result from outlet obstruction caused by appendicoliths, lymphoid hyperplasia, or other factors, leading to inflammation of the appendix. As the disease progresses, increased luminal pressure causes small vessel thrombosis, resulting in bacterial proliferation and tissue ischemia (13). These pathological changes are reflected in CT images. Although the appearance of appendicitis on CT may vary among patients, several key imaging parameters can aid in diagnosis, including appendix diameter, wall thickness, and periappendiceal fat stranding (21-23). Other important parameters include the presence of fluid or air within the appendix and abnormal enhancement of the appendix wall (24-26).

Various thresholds have been proposed for appendix diameter and wall thickness (9, 27). The current appendix diameter threshold of 6 mm is based on measurements obtained through graded compression sonography of the RLQ of the abdomen; however, CT images are acquired without compression (7). Although measuring appendix diameter is straightforward, it can be influenced by factors such as body mass index and sex. One study reported that over 20% of patients without appendicitis had an appendix diameter greater than 7 mm (10). Another study indicated that the appendix diameter exceeds 6 mm in more than 42% of healthy individuals (9). These findings are consistent with those of our study, where the mean appendix diameter in the alternative diagnosis group was 6.6 mm, and 35% of patients in this group had an appendix diameter greater than 7 mm. Setting the threshold at 6 mm resulted in a sensitivity of 99%, but a specificity of only 36%. Consequently, the 6-mm threshold may not be suitable for diagnosis, and relying solely on appendix diameter could increase the rate of negative appendectomies. Therefore, multiple imaging parameters should be assessed to avoid unnecessary treatments.

In our study, an appendix diameter threshold of 7.7 mm achieved sensitivity and specificity values of 91% and 86%, respectively. We identified the following significant predictors of appendicitis: Appendix diameter greater than 7.7 mm, abnormal appendix enhancement, and appendix erection. Abnormal appendix enhancement and changes in appendix diameter have been previously discussed in the literature, as these parameters reflect different clinical stages of appendicitis. Early inflammation may manifest as hyperemia, whereas late-stage disease may present as ischemic changes and gradual fluid accumulation.

Our evaluation of appendix shapes using CT images obtained through multiplanar reconstruction revealed that, in addition to abnormal appendix enhancement and diameter, appendix erection was a significant predictor of appendicitis ($P < 0.004$). By analyzing sonographic data, Rettenbacher et al. identified an association between appendix shape and appendicitis, indicating that appendix shape assessment can aid in ruling out the condition (28). However, previous studies have mostly relied on transverse section examinations of the appendix rather than its complete shape. The approach proposed in the current study is more comprehensive, as the accumulation of fluids within the appendix may cause gradual distension, leading to alterations in both appendix diameter and shape on imaging. Through sequential image analysis and multiplanar reconstruction, we could accurately delineate the exact shape of the appendix. Few studies have investigated the correlation between appendix shape and appendicitis using CT images obtained through multiplanar reconstruction, making our findings particularly valuable.

This study has several limitations. First, the single-center, retrospective cross-sectional design might have introduced selection bias, particularly given the low rate of negative appendectomies at our institution. Second, although abdominal pain, particularly RLQ pain, is the primary complaint in patients with acute appendicitis, we excluded patients with atypical symptoms, potentially missing cases where appendicitis presented with uncommon symptoms. Third, although measurements were conducted by experienced radiologists, the possibility of interobserver variability affecting diagnostic consistency cannot be overlooked. Finally, we excluded patients with a history of appendectomy, meaning conditions such as stump appendicitis were not considered in this study.

Although appendix diameter is often the initial parameter considered in the diagnosis of appendicitis (29), our findings suggest that the traditional threshold

of 6 mm may not be adequate for CT imaging-based evaluations. Evidence shows that patients without appendicitis can present with a wide range of appendix diameters (9, 10, 30). Therefore, additional imaging parameters should be considered to avoid unnecessary surgeries and improper treatments.

In conclusion, an appendix diameter greater than 7.7 mm, abnormal appendix enhancement, and appendix erection appear to be significant independent predictors of appendicitis in patients presenting with RLQ pain.

Footnotes

Authors' Contribution: Study concept and design: Y. C. W. and L. J. W.; acquisition of data: J. Y. H.; analysis and interpretation of data: J. Y. H. and Y. C. W.; drafting of the manuscript: J. Y. H., C. H. W., and H. W. C.; critical revision of the manuscript for important intellectual content: Y. C. W. and B. H. L.; statistical analysis: J. Y. H., L. J. W., and Y. C. W.; administrative, technical, and material support: C. H. W., H. W. C., and B. C. L.; study supervision: Y. C. W.

Conflict of Interests Statement: The authors declare no conflicts of interest that are relevant to the content of this article.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after its publication. The data are not publicly available due to their containing information that could compromise the privacy of research participants.

Ethical Approval: This study was approved by the institution reviewed board of our hospital (Reference number: 202300938B0).

Funding/Support: This article did not receive any financial support from any governmental or private organizations.

Informed Consent: Written informed consents were waived.

References

1. Ferris M, Quan S, Kaplan BS, Molodecky N, Ball CG, Chernoff GW, et al. The Global Incidence of Appendicitis. *Ann Surg*. 2017;**266**(2):237-41. <https://doi.org/10.1097/sla.0000000000002188>.
2. Coward S, Kareemi H, Clement F, Zimmer S, Dixon E, Ball CG, et al. Incidence of Appendicitis over Time: A Comparative Analysis of an Administrative Healthcare Database and a Pathology-Proven Appendicitis Registry. *PLoS One*. 2016;**11**(11). e0165161. [PubMed ID:

- 27820826]. [PubMed Central ID: PMC5098829]. <https://doi.org/10.1371/journal.pone.0165161>.
3. Petroianu A. Diagnosis of acute appendicitis. *Int J Surg*. 2012;**10**(3):115-9. [PubMed ID: 22349155]. <https://doi.org/10.1016/j.ijsu.2012.02.006>.
 4. Cartwright SL, Knudson MP. Evaluation of acute abdominal pain in adults. *Am Fam Physician*. 2008;**77**(7):971-8. [PubMed ID: 18441863].
 5. Dahabreh IJ, Adam GP, Halladay CW, Steele DW, Daiello LA, Wieland LS, et al. *Diagnosis of Right Lower Quadrant Pain and Suspected Acute Appendicitis*. Rockville: Agency for Healthcare Research and Quality (US); 2015.
 6. Heller MT, Hattoum A. Imaging of acute right lower quadrant abdominal pain: differential diagnoses beyond appendicitis. *Emerg Radiol*. 2012;**19**(1):61-73. [PubMed ID: 22072087]. <https://doi.org/10.1007/s10140-011-0997-9>.
 7. Kessler N, Cyteval C, Gallix B, Lesnik A, Blayac PM, Pujol J, et al. Appendicitis: evaluation of sensitivity, specificity, and predictive values of US, Doppler US, and laboratory findings. *Radiol*. 2004;**230**(2):472-8. [PubMed ID: 14688403]. <https://doi.org/10.1148/radiol.2302021520>.
 8. Pinto Leite N, Pereira JM, Cunha R, Pinto P, Sirlin C. CT evaluation of appendicitis and its complications: imaging techniques and key diagnostic findings. *AJR Am J Roentgenol*. 2005;**185**(2):406-17. [PubMed ID: 16037513]. <https://doi.org/10.2214/ajr.185.2.01850406>.
 9. Willekens I, Peeters E, De Maeseneer M, de Mey J. The normal appendix on CT: does size matter? *PLoS One*. 2014;**9**(5). e96476. [PubMed ID: 24802879]. [PubMed Central ID: PMC4011757]. <https://doi.org/10.1371/journal.pone.0096476>.
 10. Moskowitz E, Khan AD, Cribari C, Schroepel TJ. Size matters: Computed tomographic measurements of the appendix in emergency department scans. *Am J Surg*. 2019;**218**(2):271-4. [PubMed ID: 30558802]. <https://doi.org/10.1016/j.amjsurg.2018.12.010>.
 11. Podda M, Poillucci G, Pacella D, Mortola L, Canfora A, Aresu S, et al. Appendectomy versus conservative treatment with antibiotics for patients with uncomplicated acute appendicitis: a propensity score-matched analysis of patient-centered outcomes (the ACTUAA prospective multicenter trial). *Int J Colorectal Dis*. 2021;**36**(3):589-98. [PubMed ID: 33454817]. <https://doi.org/10.1007/s00384-021-03843-8>.
 12. Salminen P, Tuominen R, Pajajenen H, Rautio T, Nordstrom P, Aarnio M, et al. Five-Year Follow-up of Antibiotic Therapy for Uncomplicated Acute Appendicitis in the APPAC Randomized Clinical Trial. *JAMA*. 2018;**320**(12):1259-65. [PubMed ID: 30264120]. [PubMed Central ID: PMC6233612]. <https://doi.org/10.1001/jama.2018.13201>.
 13. Harnoss JC, Zelenka I, Probst P, Grummich K, Muller-Lantzsch C, Harnoss JM, et al. Antibiotics Versus Surgical Therapy for Uncomplicated Appendicitis: Systematic Review and Meta-analysis of Controlled Trials (PROSPERO 2015: CRD42015016882). *Ann Surg*. 2017;**265**(5):889-900. [PubMed ID: 27759621]. <https://doi.org/10.1097/SLA.0000000000002039>.
 14. Foley WD. CT Features for Complicated versus Uncomplicated Appendicitis: What Is the Evidence? *Radiol*. 2018;**287**(1):116-8. [PubMed ID: 29558303]. <https://doi.org/10.1148/radiol.2018180022>.
 15. Moris D, Paulson EK, Pappas TN. Diagnosis and Management of Acute Appendicitis in Adults: A Review. *JAMA*. 2021;**326**(22):2299-311. [PubMed ID: 34905026]. <https://doi.org/10.1001/jama.2021.20502>.
 16. Rud B, Vejborg TS, Rapoport ED, Reitsma JB, Wille-Jørgensen P. Computed tomography for diagnosis of acute appendicitis in adults. *Cochrane Database Syst Rev*. 2019. [PubMed ID: 31743429]. [PubMed Central ID: PMC6953397]. <https://doi.org/10.1002/14651858.CD009977.pub2>.
 17. Di Saverio S, Podda M, De Simone B, Ceresoli M, Augustin G, Gori A, et al. Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines. *World J Emerg Surg*. 2020;**15**(1):27. [PubMed ID: 32295644]. [PubMed Central ID: PMC7386163]. <https://doi.org/10.1186/s13017-020-00306-3>.
 18. van Randen A, Bipat S, Zwinderman AH, Ubbink DT, Stoker J, Boermeester MA. Acute appendicitis: meta-analysis of diagnostic performance of CT and graded compression US related to prevalence of disease. *Radiol*. 2008;**249**(1):97-106. [PubMed ID: 18682583]. <https://doi.org/10.1148/radiol.2483071652>.
 19. Birnbaum BA, Wilson SR. Appendicitis at the Millennium. *Radiol*. 2000;**215**(2):337-48. [PubMed ID: 10796905]. <https://doi.org/10.1148/radiology.215.2.r00ma24337>.
 20. Chin CM, Lim KL. Appendicitis: atypical and challenging CT appearances. *Radiograph*. 2015;**35**(1):123-4. [PubMed ID: 25590392]. <https://doi.org/10.1148/rg.351140122>.
 21. Monson B, Mandoul C, Millet I, Taourel P. Imaging of appendicitis: Tips and tricks. *Eur J Radiol*. 2020;**130**:109165. [PubMed ID: 32663765]. <https://doi.org/10.1016/j.ejrad.2020.109165>.
 22. Garcia EM, Camacho MA, Karolyi DR, Kim DH, Cash BD, Chang KJ, et al. ACR Appropriateness Criteria® Right Lower Quadrant Pain-Suspected Appendicitis. *J American College Radiol*. 2018;**15**(11):S373-87. [PubMed ID: 30392606]. <https://doi.org/10.1016/j.jacr.2018.09.033>.
 23. Song X, Shi M, Liu W, Ge Y, Wang P. Relative CT number of periappendiceal fat stranding may be an applicable index for estimating the severity of acute appendicitis. *Br J Radiol*. 2021;**94**(1120):20200437. [PubMed ID: 33684313]. [PubMed Central ID: PMC8010530]. <https://doi.org/10.1259/bjrr.20200437>.
 24. Moteki T, Horikoshi H. New CT criterion for acute appendicitis: maximum depth of intraluminal appendiceal fluid. *AJR Am J Roentgenol*. 2007;**188**(5):1313-9. [PubMed ID: 17449776]. <https://doi.org/10.2214/AJR.06.1180>.
 25. Hong HS, Cho HS, Woo JY, Lee Y, Yang I, Hwang JY, et al. Intra-Appendiceal Air at CT: Is It a Useful or a Confusing Sign for the Diagnosis of Acute Appendicitis? *Korean J Radiol*. 2016;**17**(1):39-46. [PubMed ID: 26798214]. [PubMed Central ID: PMC4720809]. <https://doi.org/10.3348/kjr.2016.17.1.39>.
 26. Park G, Lee SC, Choi BJ, Kim SJ. Stratified computed tomography findings improve diagnostic accuracy for appendicitis. *World J Gastroenterol*. 2014;**20**(38):13942-9. [PubMed ID: 25320531]. [PubMed Central ID: PMC4194577]. <https://doi.org/10.3748/wjg.v20.i38.13942>.
 27. Kim MY, Kim Y, Ryu JA, Kim TY. How to evaluate appendices with borderline diameters on CT: proposal of a quick solution to overcome the limitations of the established CT criteria. *Acad Radiol*. 2014;**21**(12):1573-8. [PubMed ID: 25127845]. <https://doi.org/10.1016/j.acra.2014.07.005>.
 28. Rettenbacher T, Hollerweger A, Macheiner P, Gritzmann N, Daniaux M, Schwamberger K, et al. Ovoid shape of the vermiform appendix: a criterion to exclude acute appendicitis-evaluation with US. *Radiol*. 2003;**226**(1):95-100. [PubMed ID: 12511674]. <https://doi.org/10.1148/radiol.226101496>.
 29. Lai V, Chan WC, Lau HY, Yeung TW, Wong YC, Yuen MK. Diagnostic power of various computed tomography signs in diagnosing acute appendicitis. *Clin Imaging*. 2012;**36**(1):29-34. [PubMed ID: 22226440]. <https://doi.org/10.1016/j.clinimag.2011.04.003>.
 30. Rettenbacher T, Hollerweger A, Macheiner P, Rettenbacher L, Tomaselli F, Schneider B, et al. Outer diameter of the vermiform appendix as a sign of acute appendicitis: evaluation at US. *Radiol*. 2001;**218**(3):757-62. [PubMed ID: 11230651]. <https://doi.org/10.1148/radiology.218.3.r01fe20757>.