Published Online: 2025 March 30

## Systematic Review



# Prevalence of Problematic Internet Use (PIU) After the COVID-19 Pandemic in Iranian Adolescent: A Systematic Review

Fatemeh Hamedi<sup>1</sup>, Nasrin Shaabani<sup>2</sup>, Fatemeh Mahdavi<sup>3</sup>, Somaye Nezamdoust<sup>4</sup>, Shifteh Arshian<sup>5</sup>, Arezoo Zare Mohammadi<sup>6,\*</sup>

<sup>1</sup> Faculty of Medical Sciences, Tabriz Branch, Islamic Azad University, Tabriz, Iran

<sup>2</sup> Shiraz Branch, Islamic Azad University, Shiraz, Iran

<sup>3</sup> Sohanek Branch, Islamic Azad University, Tehran, Iran

<sup>4</sup> Hakim Nezami Institute of Higher Education, Quchan, Iran

<sup>5</sup> Department of Psychology, Iranian University, Tehran, Iran

 $^{6}$  Tehran Central Unit, Tehran Central Branch, Islamic Azad University, Tehran, Iran

\* Corresponding Author: Tehran Central Unit, Tehran Central Branch, Islamic Azad University, Tehran, Iran. Email: arezoo.zare.mohammadi@gmail.com

Received: 11 February, 2025; Accepted: 16 March, 2025

## Abstract

**Introduction:** The widespread use of the internet has resulted in both beneficial and detrimental effects on adolescents and young adults globally. This systematic review and meta-analysis aim to examine the prevalence of problematic internet use (PIU) among Iranian adolescents following the COVID-19 pandemic.

**Methods:** We conducted a systematic search using electronic databases, including PubMed, Scopus, and Google Scholar, covering the period from January 1, 2020, to January 1, 2025. The search focused on studies examining the prevalence of internet addiction among students aged 13 - 18 years in Iran following the coronavirus epidemic. Data from these studies were analyzed using meta-analysis methods, employing the meta-prop command in STATA-17, and results were reported with 95% confidence intervals (CIs). The I-squared test was used to identify heterogeneity between studies.

Results: The meta-analysis results indicate that the prevalence of PIU among Iranian adolescents post-COVID-19 pandemic is

31% (95% CI: 17% to 45%). This finding represents a significant increase compared to pre-pandemic studies. High heterogeneity (I<sup>2</sup> = 99.43%) was observed among the studies, potentially due to variations in study populations, measurement tools, and temporal changes throughout the post-pandemic period.

**Conclusions:** The findings underscore the urgent need for culturally tailored interventions to address PIU among Iranian adolescents. Recommendations include developing effective prevention strategies, enhancing parental guidance, and implementing educational programs to promote healthy internet use. Future research should focus on understanding the long-term impacts of increased internet use and developing targeted interventions to mitigate associated risks.

Keywords: COVID-19 Pandemic, Problematic Internet Use, Iran

## 1. Introduction

Adolescents' problematic internet use (PIU) has garnered international attention (1). Recent research indicates that this phenomenon is associated with several detrimental effects. For instance, studies have shown that excessive internet use can lead to social relationship problems, increased anxiety and depression, and poorer academic performance (2). Moreover, a 2020 study found that reducing social media use to 30 minutes daily significantly enhances mental health and reduces anxiety (3). Following the COVID-19 pandemic, internet usage among Iranian adolescents has increased markedly (4). Recent studies report that the prevalence of PIU among Iranian adolescents has reached 10.3% post-pandemic. This alarming rise underscores the need for a comprehensive examination of preventive strategies. Excessive internet

Copyright @ 2025, Journal of Archives in Military Medicine. This open-access article is available under the Creative Commons Attribution-NonCommercial 4.0 (CC BY-NC 4.0) International License (https://creativecommons.org/licenses/by-nc/4.0/), which allows for the copying and redistribution of the material only for noncommercial purposes, provided that the original work is properly cited.

How to Cite: Hamedi F, Shaabani N, Mahdavi F, Nezamdoust S, Arshian S, et al. Prevalence of Problematic Internet Use (PIU) After the COVID-19 Pandemic in Iranian Adolescent: A Systematic Review. J Arch Mil Med. 2025; 13 (1): e160299. https://doi.org/10.5812/jamm-160299.

use can result in psychosocial issues, academic performance disruption, and a decline in physical health (5). Given that adolescents are at higher risk for developing PIU (6), it is crucial to develop effective preventive strategies tailored to Iranian culture. This systematic review aims to evaluate and synthesize existing evidence on PIU prevention methods for Iranian adolescents following the COVID-19 pandemic, providing practical guidance for policymakers, educators, and health professionals.

Examining PIU among adolescent students is of paramount importance. The ICD-11 approach appears to offer a realistic and reliable prevalence rate. In a study conducted using ICD-11 identification criteria, the prevalence rate for PIU was 2.98%, and for problematic gaming, it was 1.8%. Female gender predicted PIU, while male gender predicted problematic gaming (7). These concerning statistics highlight the urgent need for serious attention to this issue. Excessive internet use among adolescents is linked to numerous negative consequences, including academic decline, reduced interpersonal relationships, anxiety, depression, and decreased physical activity (8). Given that the internet has become an integral part of adolescents' lives playing a significant role in their academic, recreational, and social activities - it is vital to develop effective strategies tailored to Iranian culture for preventing and managing PIU in this age group. Doing so will not only improve adolescents' mental health and academic performance but also help prevent more serious issues in the future.

## 2. Methods

## 2.1. Search Strategy

We conducted a systematic search using electronic databases, including PubMed, Scopus, and Google Scholar, covering the period from January 1, 2020, to January 1, 2025. The search focused on studies examining the prevalence of internet addiction among students aged 13-18 years in Iran following the coronavirus epidemic. Additionally, a manual search of references in relevant review articles was performed to ensure the inclusion of all related published articles. The search strategy was developed by combining Medical Subject Headings (MeSH) terms in both Persian and English from the PubMed database, as outlined below: ("Iran"[ti] AND "Internet addiction"[ti] OR "student\*"[ti] OR "13 to 18 years of age"[ti] OR

"Adolescent"[ti]) AND (COVID-19[mesh] OR COVID-19[ti] OR "covid 19"[ti] OR covid19[ti] OR Coronavirus[ti] OR SARS-CoV-2[ti] OR "SARS CoV 2"[ti] OR 2019-nCoV[ti] OR "2019 nCoV Disease"[ti] AND 2020/01/01:2025/01/01[dp]).

Our search yielded a number of potentially relevant articles from the aforementioned electronic databases. After removing duplicates and excluding studies based on title and abstract reviews, several studies were selected for full-text screening. Ultimately, five studies were included in the final analysis due to their relevance to the subject matter. The flow diagram of the screening process and study selection is presented in Figure 1.

For each electronic database, a specific search strategy was applied. The POLIS criteria were employed to develop observational studies related to the combination of evidence for study selection: (1) Patients: Adolescent students (13 - 18 years of age), (2) outcome: Internet addiction (IA), (3) location: Iran, (4) indicator: Prevalence of internet addiction, (5) study design: Cross-sectional studies.

#### 2.2. Exclusion and Inclusion Criteria for the Studies

The inclusion criteria encompassed any studies that examined the prevalence of internet addiction among Iranian adolescent students. No restrictions were applied based on participant age, gender, or search time frame for study inclusion. This review included crosssectional studies. Case-report studies, studies lacking full text, and those focusing on other target groups such as Iranian university students were excluded from the review process.

## 2.3. Study Selection and Data Extraction

Relevant studies were selected independently by two individuals at every stage of the study selection process, including screening, full-text review, and quality assessment. A pre-designed Excel form for data extraction and quality assessment of articles was provided to the two evaluators. For data extraction, variables included the publication year, name of the study's first author, sample size, study design type, age range or mean age, target population, internet addiction prevalence rate, key findings, and quality assessment score.

#### 2.4. Qualitative Assessment of Studies

To execute this section, two independent assessors employed a modified version of the quality assessment



Figure 1. Identification of studies through databases and registries based on (9) PRISMA follow diagram (2020)

checklist for prevalence studies (adapted from Hoy et al.), specifically designed for cross-sectional prevalence studies (10). This tool comprises ten questions aimed at evaluating the risk of bias. For this study, the questions in this tool were modified to reflect the conditions of the included studies, and subsequently, a qualitative assessment of the articles was performed based on these modifications.

#### 2.5. Data Synthesis

Data from studies examining the prevalence of internet addiction among adolescent students in Iran were analyzed using meta-analysis methods, employing the meta-prop command in STATA-17, and reported with 95% confidence intervals (CIs). To identify heterogeneity between studies, the I-squared test was used. The interpretation of I-squared values was as follows:  $I^2 < 25\%$ : No heterogeneity,  $I^2 = 25 - 50\%$ : Moderate heterogeneity,  $I^2 > 50\%$ : Substantial heterogeneity. To investigate the effects of potential factors contributing

to the heterogeneity of internet addiction prevalence, Cochrane's proposed approach of meta-analysis was used for sample size. The Egger test was employed to evaluate publication bias, with a P-value of less than 0.05 deemed statistically significant.

# 3. Results

## 3.1. Features of the Included Studies

Of the total number of relevant articles that were systematically reviewed, the characteristics of a total of 5 articles are reported in Table 1.

#### 3.2. Qualitative Assessment

All studies had a low risk of bias, and this is the greatest strength of our study, as articles with high assessment quality entered the work process (Table 2).

3.3. Prevalence of Internet Addiction Among Iranian Adolescent Students During the COVID-19 Pandemic

| Table 1. Features of                       | able 1. Features of the Included Studies |                |                   |                        |  |                                     |  |  |
|--|--|----------------|-------------------|------------------------|--|-------------------------------------|--|--|
| Authors, Year                              | Men<br>Age                               | Sample<br>Size | Number<br>of Male | Number<br>of<br>Female | Type of Addiction  | Percent of<br>Internet<br>Addiction | Key Finding  |  |
| Mokhtarinia<br>et al., 2022 (11)           | 14.49                                    | 589            | 385               | 385                    | Smartphone   | 53.3                                | The increased reliance on smartphones by students during the Covid-19<br>pandemic has been linked primarily to eye strain and neck discomfort. It is<br>crucial for parents to be aware of the potential long-term musculoskeletal<br>and postural issues that may arise from prolonged smartphone use. They<br>should closely monitor their children's smartphone habits, focusing on<br>promoting proper posture and usage techniques that minimize eye strain<br>and musculoskeletal stress, with particular attention to neck health. By<br>doing so, parents can help mitigate the risk of future complications<br>associated with excessive smartphone use.    |  |
| Firoozi et al.,<br>2022 (4)                | 13 - 18                                  | 1385           | 859               | 859                    | Internet addiction   | 10.3                                | Research indicates that the onset of Internet use and duration of online<br>activity are key predictors of Internet addiction, while gender does not<br>appear to be a determining factor. The study reveals a strong association<br>between Internet addiction and leisure activities, particularly gaming, as<br>well as inadequate parental guidance. Interestingly, educational use of the<br>Internet was not found to contribute significantly to addiction. These<br>findings highlight the importance of early intervention and parental<br>involvement in shaping healthy Internet usage habits among young people,<br>especially in non-academic contexts. |  |
| Salehi et al.,<br>2022 (12)                | 13 - 18                                  | 943            | 440               | 440                    | Online addiction   | 3.6                                 | The study findings revealed an inverse relationship between secure<br>attachment styles and online addictions. Conversely, insecure attachment<br>styles demonstrated a positive correlation with all three categories of online<br>addiction examined. The research also indicated that young adults<br>exhibited higher engagement in online activities, with online addictive<br>behaviors showing an increase with age. Furthermore, a gender disparity<br>was observed in online gaming addiction, with males displaying a higher<br>prevalence compared to females.  |  |
| Shokri et al.,<br>2024 ( <mark>13</mark> ) | 13 - 19                                  | 407            | 222               | 222                    | Internet addiction   | 25                                  | The study revealed significant differences in lifestyle habits between students with and without internet addiction. Non-addicted students demonstrated markedly higher overall healthy lifestyle scores ( $P < 0.05$ ). Notably, scores in spiritual growth, health responsibility, and nutrition were significantly better among non-addicted students compared to their addicted counterparts.  |  |
| Akbari et al.,<br>2023 (14)                | 13 - 19                                  | 2390           | 1555              | 1555                   | Problematic social<br>media use with<br>gaming disorder<br>and gambling with<br>social media | 18.5                                | Adolescents experiencing concurrent problems with social media, gaming,<br>and gambling demonstrate heightened psychological vulnerabilities,<br>including increased internalizing symptoms, sensation-seeking behaviors,<br>and social anxiety.   |  |

#### Table 2. Qualitative Assessment of All Studies

| Authors and<br>Year                        | Was an Appropriate Case<br>Definition Utilized in<br>the Study? | Was the Study Instrument<br>That Measured the Parameter<br>of Internet Addiction? | Was the Same Method of<br>Data Collection Employed<br>for All Target Groups? | Was the Length of the Shortest<br>Prevalence Period for the<br>Parameter of Interest Appropriate? | Score                         |
|--|---|---|--|---|-------------------------------|
| Mokhtarinia et<br>al., 2022 (11)           | Yes   | Yes   | Yes  | Yes   | High quality of<br>assessment |
| Firoozi et al.,<br>2022 (4)                | Yes   | Yes   | Yes  | Yes   | High quality of<br>assessment |
| Salehi et al.,<br>2022 (1 <mark>2</mark> ) | Yes   | Yes   | Yes  | Yes   | High quality of<br>assessment |
| Shokri et al.,<br>2024 ( <mark>13</mark> ) | Yes   | Yes   | Yes  | Yes   | High quality of<br>assessment |
| Akbari et al.,<br>2023 (14)                | Yes   | Yes   | Yes  | Yes   | High quality of<br>assessment |

The primary outcome of interest for this study was to examine the prevalence of internet addiction among Iranian adolescent students. A total of five studies were included in the meta-analysis. Using a random-effects model, the prevalence of this behavior was calculated to be ES: 0.31 [95% CI = 0.17 - 0.45]. The heterogeneity (I<sup>2</sup>) was reported as I<sup>2</sup> = 99.43%, which is statistically significant (P = 0.000), indicating high heterogeneity among these studies (Figure 2). The eligible studies were not suitable for subgroup analysis to reduce heterogeneity (Figure 2).

# 3.4. Meta-regression

To examine the effects of potential factors influencing the heterogeneity in the prevalence of internet addiction, the Cochrane-recommended



Figure 2. Prevalence of internet addiction among Iranian adolescent students after the COVID-19 pandemic (4, 11-14)



#### Figure 3. Investigating the effects of heterogeneity through meta-regression

approach called meta-regression was used for sample size. Based on Figure 3, the results indicate that as the sample size increases in studies, the mean score of internet addiction also increases. However, this relationship is not statistically significant (P = 0.41).

## 3.5. Publication Bias

To assess publication bias and examine the smallstudy effect for this outcome, the Egger's plot method was utilized. As shown in Figure 3, there is no regular distribution of studies, indicating the presence of publication bias. In the Egger's plot, which examines the small-study effect, a P-value of 0.09 was reported. Although this value suggests a trend towards significance, it indicates the presence of publication bias (Figure 4).

## 4. Discussion

The widespread use of the internet has resulted in both beneficial and detrimental effects on adolescents and young adults worldwide. The results of this metaanalysis indicate that the prevalence of PIU among



#### Figure 4. Egger's test for small-study effects

Iranian adolescents after the COVID-19 pandemic is 31% (95% CI: 17% to 45%). This finding demonstrates a significant increase compared to pre-pandemic studies. In comparison with similar findings among Iranian university students in the post-COVID era, this result shows relatively similar outcomes. For example, a study conducted in 2022 reported the prevalence of internet addiction among Iranian university students to be 39.8% (15).

Moreover, compared to similar target groups of students before the COVID-19 pandemic, this result shows a substantial increase. In another study conducted in 2020, the prevalence of severe internet addiction among Iranian adolescents and young adults was reported to be only 2.4% (16). This significant increase could be attributed to various factors, including changes resulting from the COVID-19 pandemic. During the period of quarantine and social restrictions, adolescents spent more time online, which may have led to an increase in PIU. This finding is consistent with the results of a study conducted in 2022, which reported an increase in the prevalence of internet addiction among Tehran adolescents to 10.3% following the COVID-19 pandemic (4).

Understanding the factors contributing to the effects of increased internet use can aid in developing culturally appropriate strategies. Furthermore, the implications of these findings underscore the need for policymakers, educators, and healthcare providers to devise effective strategies to combat this emerging public health concern. The high heterogeneity ( $I^2 = 99.43\%$ ) in this meta-analysis indicates significant differences between studies. These variations may stem from factors such as differences in study populations, diverse measurement tools, and temporal changes throughout the post-pandemic period. These findings highlight the importance of addressing PIU among Iranian adolescents.

Results from a similar study suggest that adolescents' internet behavior patterns during the COVID-19 pandemic have been diverse. Researchers found that various factors such as education level, academic score rankings, family functioning, and emotional and behavioral symptoms differed across different groups. These differences could explain the high heterogeneity in studies related to PIU among adolescents (17). Given the negative consequences of internet addiction on mental health, academic performance, and social relationships, there is a need for effective preventive and therapeutic interventions.

Ultimately, this study demonstrates that PIU among Iranian adolescents has increased following the COVID-19 pandemic. These findings highlight the necessity for greater attention to this issue and the design of appropriate intervention programs. Furthermore, conducting additional studies to examine the factors influencing this increase and strategies to address it appears essential.

## 4.1. Conclusions

This meta-analysis shows that the prevalence of PIU among Iranian adolescents has significantly increased after the COVID-19 pandemic. Considering the negative impacts of this phenomenon on adolescents' mental health, academic performance, and social relationships, there is an urgent need to design and implement effective preventive and therapeutic interventions. Moreover, further studies are necessary to gain a deeper understanding of the factors contributing to this increase and to find suitable strategies to address it.

## Acknowledgements

We sincerely thank all the researchers whose published studies made this meta-analysis possible. We also express our gratitude to all colleagues and experts who assisted us at various stages of this research. The scientific and ethical support of these individuals has played a significant role in advancing this study. Finally, we extend our deepest appreciation to the families and adolescents who participated in the primary studies and provided valuable data. We hope that the results of this research will contribute effectively to improving the mental and social health of adolescents.

## Footnotes

**Authors' Contribution:** A. Z. M.: Study concept and design; F. H.: Acquisition of data; A. Z. M. and Sh. A.: Analysis and interpretation of data; F. M., A. Z. M., and S. N.: Drafting of the manuscript; F. M. and N. Sh.: Critical revision of the manuscript for important intellectual content; A. Z. M.: Statistical analysis; F. H.: Administrative, technical, and material support; F. H. and A. Z. M.: Study supervision.

**Conflict of Interests Statement:** The authors declare no conflict of interest.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication. The data are not publicly available due to confidentiality.

**Funding/Support:** The present study received no funding/support.

#### References

- Kalantari E, Zarei M, Abdolahi F. Negative Consequences Affecting the Lifestyle of Students During Covid-19 Pandemic. Social Welfare. 2023;22(87):321-55. https://doi.org/10.32598/refahj.22.87.4082.1.
- Shalchi PB, Kolahi Hamed S, Shafiee Kandjani PA. [Confirmatory Factor Analysis of Problematic Internet Entertainment Use Scale for Adolescents (PIEUSA)]. *Quarterly J Family Res.* 2021;**18**(1):105-20. FA.
- Bernard CE. Lonely zs: examining the relationships among time spent on social networking sites, the fear of missing out (fomo), and loneliness among undergraduate college students [dessertation]. Lynchburg, VA: Liberty University; 2020.
- Firoozi M, Firoozi AA, Karimi Riabi E. The Global Prevalence of Internet Addiction in Adolescents Post-COVID-19 Period: Examining Iran's Situation. J World Sociopolitical Studies. 2022;6(4):673-703. https://doi.org/10.22059/wsps.2023.358588.1346.
- Paulus FW, Joas J, Gerstner I, Kuhn A, Wenning M, Gehrke T, et al. Problematic Internet Use among Adolescents 18 Months after the Onset of the COVID-19 Pandemic. *Children (Basel)*. 2022;9(11). [PubMed ID: 36360452]. [PubMed Central ID: PMC9689314]. https://doi.org/10.3390/children9111724.
- Romero Saletti SM, Van den Broucke S, Chau C. The effectiveness of prevention programs for problematic Internet use in adolescents and youths: A systematic review and meta-analysis. *Cyberpsychology: Journal of Psychosocial Research on Cyberspace*. 2021;15(2). https://doi.org/10.5817/cp2021-2-10.
- Nogueira-Lopez A, Rial-Boubeta A, Guadix-Garcia I, Villanueva-Blasco VJ, Billieux J. Prevalence of problematic Internet use and problematic gaming in Spanish adolescents. *Psychiatry Res.* 2023;**326**:115317. [PubMed ID: 37352749]. https://doi.org/10.1016/j.psychres.2023.115317.
- Jelleli H, Ben Aissa M, Kaddech N, Saidane M, Guelmami N, Bragazzi NL, et al. Examining the interplay between physical activity, problematic internet use and the negative emotional state of depression, anxiety and stress: insights from a moderated mediation path model in university students. *BMC Psychol.* 2024;**12**(1):406. [PubMed ID: 39054507]. [PubMed Central ID: PMC11274773]. https://doi.org/10.1186/s40359-024-01736-3.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;**372**:n71. [PubMed ID: 33782057]. [PubMed Central ID: PMC8005924]. https://doi.org/10.1136/bmj.n71.
- Hoy D, Brooks P, Woolf A, Blyth F, March L, Bain C, et al. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. J Clin Epidemiol. 2012;65(9):934-9. [PubMed ID: 22742910]. https://doi.org/10.1016/j.jclinepi.2011.11.014.
- Mokhtarinia HR, Torkamani MH, Farmani O, Biglarian A, Gabel CP. Smartphone addiction in children: patterns of use and musculoskeletal discomfort during the COVID-19 pandemic in Iran. *BMC Pediatr.* 2022;22(1):681. [PubMed ID: 36435748]. [PubMed Central ID: PMC9701359]. https://doi.org/10.1186/s12887-022-03748-7.
- Salehi E, Fallahchai R, Griffiths M. Online Addictions Among Adolescents and Young Adults in Iran: The Role of Attachment Styles and Gender. Social Science Computer Review. 2022;41(2):554-72. https://doi.org/10.1177/08944393221111242.
- Shokri A, Mohamadi A, Mohammadi D, Moradi M, Sadeghi S, Mahmoodi H, et al. The relationship between internet addiction and lifestyle among high school students: A cross sectional in the west of Iran. *PLoS One*. 2024;**19**(9). e0308333. [PubMed ID: 39240897].

| [PubMed          | Central              | ID:       | PMC11379132 |
|------------------|----------------------|-----------|-------------|
| https://doi.org/ | 10.1371/journal.pone | .0308333. |             |

- Akbari M, Bahadori MH, Khanbabaei S, Milan BB, Horvath Z, Griffiths MD, et al. Psychological predictors of the co-occurrence of problematic gaming, gambling, and social media use among adolescents. *Computers in Human Behavior*. 2023;**140**. https://doi.org/10.1016/j.chb.2022.107589.
- 15. Toumari R, Ramezani-Badr F, Bulfone G, Amini K. Prevalence of social media addiction and its related factors among students of nursing and midwifery in Iranian context: A cross-sectional study. *Journal of*

 Multidisciplinary
 Care.
 2023;12(1):3-11.

 https://doi.org/10.34172/jmdc.1132.

- 16. Mousavi SV. [Prevalence of Internet addiction and the Status of the use of virtual social networks in Iranian Teenagers and Youths in 2018]. *Mil Med*. 2020;**22**(3):281-8. FA.
- Wu Q, Ren Q, Zhong N, Bao J, Zhao Y, Du J, et al. Internet behavior patterns of adolescents before, during, and after COVID-19 pandemic. *Front Psychiatry*. 2022;13:947360. [PubMed ID: 36440398]. [PubMed Central ID: PMC9685307]. https://doi.org/10.3389/fpsyt.2022.947360.