Published Online: 2025 March 8

Research Article



Assessing the Impact of Demographics Factors on Healthcare Professionals' Performance: A Cross-Sectional Study

Saeed Biroudian¹, Mehdi Mohit², Shahriar Mousavinejad³, Hamid Mohammadi Kojidi⁴, Zohreh Kazempour Keleshteri², Mirsaeed Attarchi⁵, Kourosh Delpasand (b^{6,*})

³ Assistant Professor of Medical Ethics, Department of Medical Ethics, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴ Associate Professor of Forensic Medicine, Department of Forensic Medicine, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran

⁵ Professor of Occupational Medicine, Department of Forensic Medicine, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran ⁶ Associate Professor of Medical Ethics, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran

* Corresponding Author: Associate Professor of Medical Ethics, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran. Email: kd388@yahoo.com

Received: 5 November, 2024; Revised: 25 February, 2025; Accepted: 2 March, 2025

Abstract

Background: The safeguarding of patients' rights is a fundamental aspect of healthcare that ensures ethical and respectful treatment. Healthcare professionals play a critical role in upholding these rights, which include informed consent, confidentiality, and the right to receive appropriate care.

Objectives: This study evaluated the performance of healthcare professionals in safeguarding patients' rights at Guilan University of Medical Sciences, Rasht, Iran.

Methods: A cross-sectional design was employed, involving 130 healthcare professionals. Data on demographics (age, gender, marital status, occupation, and education) were collected. Performance was assessed using a self-designed questionnaire. All data were analyzed using SPSS version 21, with a significance level set at less than 0.05.

Results: Participants had a mean age of 29.42 years, were predominantly female (59.2%), with most being single (56.2%) and holding bachelor's degrees (61.5%). The average performance score was 8.38 out of a maximum score, indicating moderate performance. Statistical analysis revealed no significant differences in scores based on demographic variables (P > 0.05), despite trends suggesting higher scores among females and medical doctors.

Conclusions: The findings indicated that demographic factors do not significantly impact healthcare professionals' performance in safeguarding patients' rights. This underscores the necessity for targeted training and policy interventions to enhance the understanding and implementation of patient rights within healthcare settings.

Keywords: Healthcare Professionals, Patient Rights, Informed Consent

1. Background

The safeguarding of patients' rights is a cornerstone of modern healthcare systems, reflecting the fundamental ethical principles of beneficence, nonmaleficence, autonomy, and justice. As healthcare delivery becomes increasingly complex, the role of healthcare professionals in protecting and promoting these interests has gained paramount importance (1-3). The concept of patients' rights has evolved significantly over the past century, encompassing a broad spectrum of entitlements, including the right to informed consent, confidentiality, dignity, and autonomy in medical decision-making (4, 5). It requires healthcare professionals to navigate complex ethical dilemmas, balance competing priorities, and make decisions that align with both professional standards and individual patient needs. The multifaceted nature of this

¹ Assistant Professor of Medical Ethics, Department of Medical Ethics, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

² Razi Clinical Research Development Unit, Razi Hospital, Guilan University of Medical Sciences, Rasht, Iran

Copyright © 2025, Biroudian et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

responsibility demands a nuanced approach to performance assessment (6, 7).

Conflict of interest occurs when a professional's judgment regarding а primary interest is inappropriately influenced by a secondary interest. It arises when an individual in a position of trust has personal or group interests that conflict with their professional duties (8). This situation can lead to compromised decision-making, even unintentionally, where the primary interest may be neglected in favor of the secondary one. Importantly, an actual compromise of the primary interest isn't necessary for a conflict to exist; the mere potential for such compromise due to existing conditions or relationships is sufficient to constitute a conflict of interest (9, 10).

Recent years have witnessed a growing emphasis on patient-centered care, shared decision-making, and the recognition of patients as active participants in their healthcare journey. This paradigm shift has further underscored the importance of healthcare professionals' ability to effectively safeguard patients' interests (11, 12). It has also highlighted the need for robust assessment methodologies that can capture the subtleties and complexities of this aspect of healthcare delivery.

2. Objectives

However, developing effective assessment strategies in this area presents significant challenges. The subjective nature of many aspects of patient safeguarding, the diversity of healthcare settings and specialties, and the potential influence of systemic factors on individual performance all contribute to the complexity of this task (13-15). The current study investigated healthcare professionals' performance regarding safeguarding patients' rights.

3. Methods

This research employed a cross-sectional design to assess the performance of healthcare professionals in safeguarding patients' rights at Guilan University of Medical Sciences in Rasht, Iran. The study aimed to evaluate how well healthcare professionals understand and implement patient rights, which are essential for ethical healthcare delivery.

3.1. Inclusion Criteria

Participants included a diverse range of healthcare professionals, specifically nurses, residents, general physicians, and medical students. This variety allowed us to capture a comprehensive perspective on performance related to patient rights across different levels of experience and specialization. All individuals who consented to participate were included in the study. Informed consent was obtained from each participant prior to data collection, ensuring that they understood the purpose of the study and their right to withdraw at any time without any consequences.

3.2. Exclusion Criteria

Healthcare professionals with less than six months of experience in their respective roles were excluded to ensure that participants had sufficient exposure to patient interactions and relevant ethical considerations.

Data were collected through a self-designed questionnaire that included demographic information (age, gender, marital status, occupation, and education level) as well as specific questions related to knowledge and practices concerning patients' rights. The questionnaire was developed based on existing literature and guidelines on patient rights to ensure relevance and comprehensiveness. Informed consent was obtained from all participants prior to data collection, ensuring that they understood the purpose of the study and their right to withdraw at any time.

Data were analyzed using IBM SPSS Statistics version 26. Descriptive statistics were calculated to summarize demographic characteristics and performance scores. The Kolmogorov-Smirnov test was used to assess the normality of the data distribution. For comparisons between groups, t-tests and ANOVA were employed as Additionally, Pearson appropriate. correlation coefficients were calculated to explore relationships between demographic variables and performance scores. Multiple linear regression analysis was conducted to determine associations between demographic characteristics and informed consent quality scores.

4. Results

A total of 130 healthcare professionals were enrolled in the study. The mean age of participants was $29.42 \pm$ 5.54 years (range, 22 - 51 years), and the majority were female (59.2%). Most healthcare professionals were

Fable 1. Demographic Characteristics of Participants (N = 130)					
Variables	No. (%)				
Gender					
Male	53 (40.8)				
Female	77 (59.2)				
Marital status					
Single	73 (56.2)				
Married	57 (43.8)				
Educational status					
Bachelor degree	80 (61.5)				
Medical student	6(4.6)				
Medical doctor	44 (33.8)				
Occupation					
Nurse	74 (56.9)				
General physician	13 (10.0)				
Resident	13 (10.0)				
Student	30 (23.1)				

ariables	Score	t	df ^b	P-Value ^c
ender		0.68	128	0.498 ^d
Male	8.24 ± 2.06			
Female	8.47 ± 1.66			
larital status		1.015	128	0.312 ^d
Single	8.52 ± 1.98			
Married	8.19 ± 1.61			
ducational status		1.64	127	0.197 ^e
Bachelor degree	8.35 ± 1.77			
Medical student	7.17 ± 1.72			
Medical doctor	8.59 ± 1.91			
Occupation		0.713	126	0.546 ^d
Nurse	8.46 ± 1.74			
General physician	7.79 ± 1.81			
Resident	8.69 ± 1.84			
Student	8.54 ± 2.33			

 a Values are expressed as mean \pm SD.

^b Degree of freedom within a group.

^c Significant level < 0.05.

^dt-test.

^e ANOVA.

single (56.2%), nurses (56.9%), and held bachelor's degrees (61.5%). The mean performance score among healthcare professionals was 8.38 ± 1.83 , with scores ranging from 4 to 14 (Table 1).

Analysis of professional performance scores showed no statistically significant differences based on gender,

marital status, educational status, or occupation (P > 0.05). While females, single individuals, medical doctors, and residents tended to have higher performance scores, these differences were not statistically significant (P > 0.05). For instance, the mean score for females was 8.52 ± 1.79 compared to 8.23 ± 1.87

for males, but this difference did not reach statistical significance (Table 2).

Regression analysis revealed that none of the variables (gender, marital status, occupation, and educational status) significantly predicted performance scores (P > 0.05). Although gender had a positive unstandardized coefficient (B = 0.060), indicating a potential positive association, this effect was not statistically significant (P > 0.05) (Table 3).

5. Discussion

Protecting patients' rights is a foundational principle in contemporary healthcare, encompassing the ethical obligation to respect individuals' autonomy, dignity, and privacy in medical settings. These rights, which include informed consent, the right to refuse treatment, confidentiality, and access to care, are crucial for maintaining the trust that underpins patient-provider relationships (2, 16). In an era where healthcare systems are rapidly evolving due to technological advancements, policy shifts, and the increasing complexity of care delivery, ensuring that these rights are upheld is more important than ever. Ethical medical practices require that healthcare professionals remain vigilant in their efforts to balance clinical objectives with respect for patients' choices and needs (17, 18). Moreover, as healthcare becomes more interdisciplinary and datadriven, there is a growing need to continually evaluate how well healthcare systems support patient rights, adapting policies and practices to meet emerging challenges (19, 20). Protecting these rights not only safeguards individuals from harm or neglect but also fosters a healthcare environment rooted in respect, transparency, and accountability.

In the current study, the average performance score suggested a moderate level of performance in safeguarding patients' rights. This result aligns with previous studies that have highlighted varying levels of awareness and implementation of patient rights among providers. Despite the healthcare moderate performance score, our analysis showed no statistically significant differences in performance based on demographic factors such as gender, marital status, educational status, or occupation (P > 0.05). This lack of significant findings contrasts with some existing literature that suggests demographic factors can influence healthcare professionals' knowledge and

practice regarding patient rights. For instance, studies by Alraimi and Shelke (21) and Czajowska et al. (22) have indicated that specific demographic characteristics may correlate with higher awareness and better communication of patients' rights. However, our results suggest that the factors influencing performance may be more nuanced than previously understood.

Interestingly, although females, single individuals, medical doctors, and residents exhibited higher average performance scores, these differences were not statistically significant. This trend raises important questions about the underlying reasons for these observations. For example, while female healthcare professionals often report higher levels of empathy and communication skills – traits essential for safeguarding patient rights – our study did not find a corresponding statistical significance in their performance scores. This discrepancy may indicate that other factors, such as institutional support or training opportunities, play a more critical role in determining performance than demographic characteristics alone.

The regression analysis further confirmed that none of the demographic variables significantly impacted performance scores (P > 0.05). However, the positive unstandardized coefficient for gender (B = 0.060) suggests a potential association worth exploring in future research. This finding aligns with the notion that gender dynamics within healthcare settings may influence professional interactions and patient advocacy efforts. Moreover, the absence of significant differences based on marital status contradicts some studies that have linked marital status to job satisfaction and burnout levels among healthcare professionals.

For instance, previous research has shown that married individuals often report higher job satisfaction compared to their single counterparts (23-25). However, our findings suggest that marital status may not directly influence performance in safeguarding patients' rights within this specific context. The moderate performance level observed in this study underscores the necessity for targeted educational programs and policy interventions aimed at enhancing healthcare professionals' understanding and implementation of patient rights. As healthcare systems become increasingly complex due to technological advancements and evolving patient needs, it is crucial to

Variables	Unstandardized Coefficients		Standardized Coefficients		D Value
	В	SE	Beta	i	r-value
Gender	0.060	0.034	0.181	7.392	0.083
Marital status	-0.615	0.368	-0.168	1.749	0.098
Occupation	-0.029	0.091	-0.046	-1.668	0.755
Educational status	-0.123	0.194	-0.096	-0.313	0.528

Abbreviation: SE, standard error.

^a Significant level < 0.05.

ensure that all healthcare providers are adequately trained to uphold ethical standards related to patient care.

This study had several limitations: First, the relatively small sample size and the focus on a single healthcare setting might limit generalizability. Second, while the study collected demographic data (age, gender, marital status, occupation, education), it did not explore other potentially influential factors such as years of experience, specific training related to patient rights, or institutional policies that might impact performance. Future research could benefit from larger, multi-center studies to validate these findings across diverse healthcare environments. These findings underscore the need for comprehensive, targeted interventions to enhance healthcare professionals' competencies in protecting and promoting patients' rights across all demographic groups.

5.1. Conclusions

The lack of significant differences in performance across demographic groups highlights the importance of developing comprehensive training programs that are universally applicable. These programs should focus on enhancing competencies in safeguarding patients' rights, regardless of healthcare professionals' gender, marital status, occupation, or educational background. Additionally, exploring other factors that might influence performance, such as work environment or training experiences, could provide further insights into improving patient care.

Acknowledgements

The authors thank the patients for their cooperation.

Footnotes

Authors' Contribution: Study concept and design: K. D., S. B., and S. M.; Analysis and interpretation of data: H. M. and S. B.; Drafting of the manuscript: M. M. and M. A.; Critical revision of the manuscript for important intellectual content: H. M., Z. K., M. A., and S. B.; Statistical analysis: H. M.

Conflict of Interests Statement: Authors declare there are no conflicts of interest regarding the publication of this article.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: This study design was approved by the Ethical Committee of Guilan University of Medical Sciences, Rasht, Iran (IR.GUMS.REC.1402.172).

Funding/Support: This study was not supported by grant.

Informed Consent: Informed consent was obtained from each participant prior to data collection.

References

- Rosen MA, DiazGranados D, Dietz AS, Benishek LE, Thompson D, Pronovost PJ, et al. Teamwork in healthcare: Key discoveries enabling safer, high-quality care. *Am Psychol.* 2018;73(4):433-50. [PubMed ID: 29792459]. [PubMed Central ID: PMC6361117]. https://doi.org/10.1037/amp0000298.
- Varkey B. Principles of Clinical Ethics and Their Application to Practice. *Med Princ Pract*. 2021;**30**(1):17-28. [PubMed ID: 32498071]. [PubMed Central ID: PMC7923912]. https://doi.org/10.1159/000509119.
- 3. Young M, Wagner A. *Medical Ethics*. USA: StatPearls; 2025. [PubMed ID: 30570982].
- Bazzano LA, Durant J, Brantley PR. A Modern History of Informed Consent and the Role of Key Information. *Ochsner J.* 2021;21(1):81-5.

[PubMed ID: 33828429]. [PubMed Central ID: PMC7993430]. https://doi.org/10.31486/t0j.19.0105.

- Buka P. Essential Law and Ethics in Nursing. Oxfordshire, England: Routledge; 2020. https://doi.org/10.4324/9780429292187.
- Haddad LM, Geiger RA. Nursing Ethical Considerations. USA: StatPearls; 2025. [PubMed ID: 30252310].
- Thacharodi A, Singh P, Meenatchi R, Tawfeeq Ahmed ZH, Kumar RRS, V N, et al. Revolutionizing healthcare and medicine: The impact of modern technologies for a healthier future-A comprehensive review. *Health Care Sci.* 2024;3(5):329-49. [PubMed ID: 39479277]. [PubMed Central ID: PMC11520245]. https://doi.org/10.1002/hcs2.115.
- Rodwin MA. Conflicts of interest in medicine: should we contract, conserve, or expand the traditional definition and scope of regulation. J Health Care. 2018;21:157.
- Bester J, Cole CM, Kodish E. The Limits of Informed Consent for an Overwhelmed Patient: Clinicians' Role in Protecting Patients and Preventing Overwhelm. AMA J Ethics. 2016;18(9):869-86. [PubMed ID: 27669132]. https://doi.org/10.1001/journalofethics.2016.18.9.peer2-1609.
- Thirumoorthy T. Conflicts of interest in medicine: understanding the concepts to preserve the integrity of professional judgement and promote trust in the profession. *Singapore Med J.* 2023;64(2):121-6. [PubMed ID: 36814176]. [PubMed Central ID: PMC10071851]. https://doi.org/10.4103/singaporemedj.SMJ-2022-217.
- Edgman-Levitan S, Schoenbaum SC. Patient-centered care: achieving higher quality by designing care through the patient's eyes. *Isr J Health Policy Res.* 2021;10(1):21. [PubMed ID: 33673875]. [PubMed Central ID: PMC7934513]. https://doi.org/10.1186/s13584-021-00459-9.
- Kwame A, Petrucka PM. A literature-based study of patient-centered care and communication in nurse-patient interactions: barriers, facilitators, and the way forward. *BMC Nurs.* 2021;**20**(1):158. [PubMed ID: 34479560]. [PubMed Central ID: PMC8414690]. https://doi.org/10.1186/s12912-021-00684-2.
- Bamforth K, Rae P, Maben J, Lloyd H, Pearce S. Perceptions of healthcare professionals' psychological wellbeing at work and the link to patients' experiences of care: A scoping review. *Int J Nurs Stud Adv.* 2023;**5**:100148. [PubMed ID: 38746580]. [PubMed Central ID: PMC11080414]. https://doi.org/10.1016/j.ijnsa.2023.100148.
- Ferreira DC, Vieira I, Pedro MI, Caldas P, Varela M. Patient Satisfaction with Healthcare Services and the Techniques Used for its Assessment: A Systematic Literature Review and a Bibliometric Analysis. *Healthcare (Basel)*. 2023;11(5). [PubMed ID: 36900644]. [PubMed Central ID: PMC10001171]. https://doi.org/10.3390/healthcare11050639.
- Molina-Mula J, Gallo-Estrada J. Impact of Nurse-Patient Relationship on Quality of Care and Patient Autonomy in Decision-Making. Int J

Environ Res Public Health. 2020;17(3). [PubMed ID: 32013108]. [PubMed Central ID: PMC7036952]. https://doi.org/10.3390/ijerph17030835.

- Kwame A, Petrucka PM. Universal healthcare coverage, patients' rights, and nurse-patient communication: a critical review of the evidence. *BMC Nurs*. 2022;21(1):54. [PubMed ID: 35255908]. [PubMed Central ID: PMC8900414]. https://doi.org/10.1186/s12912-022-00833-1.
- Al-Wathinani AM, Barten DG, Alsahli H, Alhamid A, Alghamdi W, Alqahtani W, et al. The Right to Refuse: Understanding Healthcare Providers' Perspectives on Patient Autonomy in Emergency Care. *Healthcare (Basel)*. 2023;11(12). [PubMed ID: 37372874]. [PubMed Central ID: PMC10297854]. https://doi.org/10.3390/healthcare11121756.
- Hall MA, Bobinski WJ, MA Orentlicher D. Health care law and ethics. USA: Aspen Pub; 2003.
- Moosavi S, Mousavi MS, Ahmadi A, Mardani A, Parsapoor A, Gooshki ES. Respecting patients' rights in hospitals: patients' and health-care workers' perspectives. J Med Ethics History Med. 2023;16. https://doi.org/10.18502/jmehm.v16i13.14308.
- Renukappa S, Mudiyi P, Suresh S, Abdalla W, Subbarao C. Evaluation of challenges for adoption of smart healthcare strategies. *Smart Health*. 2022;26. https://doi.org/10.1016/j.smhl.2022.100330.
- Alraimi A, Shelke PA. Nurses' Level of Awareness of Patients' Rights in hospitals. ASK Pub. 2023;2(3). https://doi.org/10.21203/rs.3.rs-3682473/v1.
- Czajkowska M, Janik A, Zborowska K, Plinta R, Brzek A, Skrzypulec-Plinta V. Knowledge and opinions of patients and medical staff about patients' rights. *Ginekol Pol.* 2021;92(7):491-7. [PubMed ID: 33757155]. https://doi.org/10.5603/GP.a2021.0014.
- Chen YH, Lou SZ, Yang CW, Tang HM, Lee CH, Jong GP. Effect of Marriage on Burnout among Healthcare Workers during the COVID-19 Pandemic. Int J Environ Res Public Health. 2022;19(23). [PubMed ID: 36497885]. [PubMed Central ID: PMC9737389]. https://doi.org/10.3390/ijerph192315811.
- Peng J, Wu WH, Doolan G, Choudhury N, Mehta P, Khatun A, et al. Marital Status and Gender Differences as Key Determinants of COVID-19 Impact on Wellbeing, Job Satisfaction and Resilience in Health Care Workers and Staff Working in Academia in the UK During the First Wave of the Pandemic. *Front Public Health*. 2022;**10**:928107. [PubMed ID: 35832284]. [PubMed Central ID: PMC9271694]. https://doi.org/10.3389/fpubh.2022.928107.
- Singh T, Kaur M, Verma M, Kumar R. Job satisfaction among health care providers: A cross-sectional study in public health facilities of Punjab, India. J Family Med Prim Care. 2019;8(10):3268-75. [PubMed ID: 31742154]. [PubMed Central ID: PMC6857391]. https://doi.org/10.4103/jfmpc.jfmpc_600_19.