



Optimization of Fiber Dyeing with Sour Tea (*Hibiscus sabdariffa*) to Achieve Red to Brown Shades and Investigate Antimicrobial Properties

Seyed Mahmoud Tabatabaei Hanzaei¹, Mohammad Khajeh Mehrizi², Marziye Rezaei^{3,*}

¹ Assistant Professor, Department of Carpet, Textile & Fashion Design, Faculty of Art, Science and Arts University, Yazd, Iran

² Department of Textile Engineering, Yazd University, Yazd, Iran

³ Department of Carpet, Faculty of Art and Architecture, University of Sistan and Baluchestan, Zahedan, Iran

*Corresponding Author: Department of Carpet, Faculty of Art and Architecture, University of Sistan and Baluchestan, Zahedan, Iran. Email: m.rezaei@arts.usb.ac.ir

Received: 22 December, 2024; Revised: 24 December, 2024; Accepted: 25 December, 2024

Abstract

Background: In recent years, the use of chemical dyes in the textile industry has caused numerous environmental problems. Consequently, there has been growing interest in natural dyes, particularly those derived from plants, as alternatives. The search for new dye sources, including plant-based options, has become a focal point.

Objectives: This study investigated the potential use of the sour tea plant (*Hibiscus sabdariffa*) as a dye for silk products.

Methods: To dye silk goods, it is necessary to clean the silk yarn to remove waste materials such as grease, wax, thorns, shavings, and dirt before the dyeing process. For this purpose, 2 g/L of non-ionic textile soap was used for washing. The silk yarn was soaked and washed in a bath containing 2 g/L of detergent at a liquor ratio (L:R) of 1:50 for 30 minutes at 60°C. Initially, the yarns were introduced into the bath at 30°C, and the temperature was gradually increased to 60°C, where they were washed for 30 minutes. The antimicrobial properties were assessed using the disk diffusion method, measuring the diameter of the inhibition zones against bacteria.

Results: The results of the color difference analysis of the samples revealed that the greatest color difference was associated with the dry dye. In this type of dyeing, both in the metachrome and prechrome methods, the oxalic acid variable produced a greater color difference compared to acetic acid. Conversely, in dyeing using the dye extract method, acetic acid resulted in a greater color difference. In the method using the extract of dandelion root, the largest color difference was observed in both the prechrome and metachrome methods. The findings also indicated that the diameter of the inhibition zone for the sour tea extract at a concentration of 50 mg/mL was largest against *Staphylococcus aureus* (6 mm), while the smallest inhibition zone diameter at the same concentration was observed against *Enterococcus faecalis* (4 mm).

Conclusions: The results of this study demonstrated that the sour tea plant possesses effective color-generating and antimicrobial properties.

Keywords: Sour Tea, Dyeing, Silk, Antimicrobial Properties

1. Background

Until the mid-19th century, when synthetic dyes became prevalent, all dyes were derived from natural sources. Before their application in textile dyeing, natural dyes were used for body painting, cosmetics, and coloring handicrafts such as pottery and baskets. Over 15,000 years ago, our ancestors utilized natural pigments to decorate cave walls.

Significant advancements in dyes, chemicals, and dyeing processes have occurred within the textile industry. However, this industry has a substantial

environmental impact, particularly in terms of water and soil pollution. During the dyeing process, chemicals are released into the environment, contaminating water and soil, as well as emitting harmful gases into the atmosphere. Many of the dyes and chemicals used in these processes are toxic and non-biodegradable, leading to adverse changes in the natural environment.

These disruptions to natural resources contribute to altered air flows, increased heat, and elevated atmospheric temperatures. The concept of a green or environmentally friendly approach is not new in the textile industry; it aligns with the broader goal of

environmental protection. Science and technology have made significant efforts to develop eco-friendly processes aimed at minimizing environmental harm.

Recently, there has been renewed interest in using natural dyes in textile dyeing. This resurgence is driven by stringent regulations imposed by many countries in response to the allergic and toxic effects associated with synthetic dyes. It is widely believed that natural dyes are more environmentally friendly than synthetic alternatives. Natural dyes are characterized by better biodegradability and a reduced environmental footprint.

Hibiscus sabdariffa L., belonging to the Malvaceae family, is a small tropical annual shrub native to Africa and widely distributed in Southeast Asia and Central America (1, 2). Commonly referred to as hibiscus, *H. sabdariffa* is internationally recognized for its diverse uses. Many parts of the plant are utilized and prescribed in traditional medicine in countries such as those in Africa, India, Mexico, Brazil, China, and Iran (3).

The leaves, consumed as a vegetable, exhibit diuretic, antiseptic, digestive, purgative, sedative, and astringent properties (4). The calyces are traditionally used to manage high blood pressure, treat liver disorders, and serve as a diuretic, digestive aid, and sedative (5, 6). Compared to other parts of the plant, the seeds are less frequently mentioned in traditional medicine. However, they are roasted and consumed as food and are traditionally used as a demulcent, diuretic, laxative, and tonic (3).

2. Methods

2.1. Steps for Conducting Experiments

2.1.1. Washing Silk Items

To prepare silk goods for dyeing, it is necessary to remove waste materials such as grease, wax, thorns, shavings, and dirt. For this purpose, 2 g/L of non-ionic textile soap was used to wash the silk yarn.

The silk yarns were soaked and washed in a bath containing a solution of 2 g/L of detergent at a liquor ratio (L:R) of 1:50 for 30 minutes at a temperature of 60°C. The items were introduced into the bath at an initial temperature of 30°C, which was gradually raised to 60°C, and the washing process was carried out for 30 minutes at this temperature.

2.2. How to Extract

To extract the dye from the sour tea plant, 50 g of the plant material were boiled in 1 L of water until the

volume was reduced by 50%. To maintain consistent dyeing conditions in both the extract and dry dye states, and due to the presence of natural acids in the plant, the use of any chemicals during the extraction process was avoided.

2.3. Dyeing Method

Silk goods were dyed using two methods: Prechrome and metachrome, in both dry dye and extract dye forms. To preserve the standard conditions and account for the sensitivity of silk fiber, low-concentration acid and near-neutral conditions (pH 5) with a mordant concentration of 7% were used. The amount of dye for all samples was standardized:

- For the dry dye method: Thirty percent of the dry dye by weight.

- For the extract dye method: Thirty percent of the bath volume, equivalent to 150 g/L of extract, was used.

In the prechrome dyeing process, the silk samples were dyed in a dye bath with a L:R of 1:50 for 45 minutes. The dyestuff was introduced into the dye bath at a L:R of 1:40 at an initial temperature of 40°C and left for 10 minutes. The silk samples were then introduced into the bath at the same temperature (40°C) and gradually heated to boiling over 30 minutes. The dyeing process was continued at boiling temperature for 60 minutes.

3. Results

3.1. Dyed Samples

Figure 1 illustrates the samples dyed using the extraction method and the dry method. As evident from the Table 1, the samples dyed with the dye extract appear darker than those dyed with the dry dye. Notably, in the samples dyed using the dry dye under the prechrome and metachrome A methods, half of the samples displayed negative A values. This indicates that some samples in this method tended toward a green hue. In contrast, none of the samples dyed using the extraction method exhibited negative A values. It is important to note that the negative A values in the dry method were associated with samples treated with aluminum sulfate and iron sulfate.

3.2. Antibacterial Results of the Medicinal Plant Sour Tea on Standard Bacteria

This study demonstrated that the diameter of the inhibitory zone of sour tea extract at a concentration of 50 mg/mL was largest against *Staphylococcus aureus* (6 mm) and smallest against *Enterococcus faecalis* (4 mm). Furthermore, at a concentration of 100 mg/mL, the sour

Table 1. Inhibition Zone Diameter of Sour Tea Plant Extract on Standard Bacteria at Two Different Concentrations (mm)

Bacterial Strain	Concentration 50 mg/mL	Concentration 100 mg/mL
<i>Staphylococcus aureus</i>	6	10
<i>Streptococcus pyogenes</i>	3.4	7
<i>Streptococcus pneumoniae</i>	5	8
<i>Enterococcus faecalis</i>	4	6









































Dry pre-chrome	Dry metachrome	Pre-chrome extraction	Metachrome extraction	Code	
				AaSa	1
				AaH	2
				AaM	3
				AaSc	4
				AaSm	5
				AoS	6
				AoH	7
				AoM	8
				AoS	9
				AoSMP	10

Figure 1. Samples dyed by extraction and dry methods

tea extract exhibited the largest inhibitory zone diameter against *S. aureus* (10 mm), as shown in [Table 1](#) and [Figure 2](#).

4. Discussion

In general, the results of the color difference analysis of the samples indicated that the greatest color difference was associated with the dry dye. In this type of dyeing, both in the metachrome and prechrome methods, the oxalic acid variable resulted in a greater color difference compared to acetic acid. Conversely, in

the dyeing process using the dye extract method, acetic acid produced a greater color difference.

For dyeing with the extract of the toothed thorn, the greatest color difference was observed in both the prechrome and metachrome methods. Meanwhile, in the samples dyed with dry dyes under the prechrome and metachrome methods, half of the samples exhibited negative A values, indicating a tendency toward a green hue. In contrast, none of the samples dyed using the extraction method showed negative A values. It is noteworthy that the negative A values in




Figure 2. Photograph showing the bacterial inhibitory zones of sour tea extract at concentrations of 50 and 100 mg/mL.

these samples were related to those dyed with aluminum sulfate and iron sulfate.

In another study that synthesized nanoparticles using sour tea extract and examined their antimicrobial activity, the results revealed strong inhibitory effects against bacteria. Specifically, the nanoparticles demonstrated high efficacy against *Escherichia coli*, *S. aureus*, *S. intermedius*, *Proteus mirabilis*, and *Pseudomonas aeruginosa*, with inhibitory concentration (IC_{50}) values ranging from 0.05 ± 0.15 to 4.64 ± 0.05 mg/mL (7).

In another study, the results indicated that the crude metabolites of the methanolic extract of *H. sabdariffa* (flowers) demonstrated relatively significant antimicrobial activity against gram-positive pathogens. Specifically, the diameter of the inhibition zone against *S. aureus* was 13 mm at a concentration of 5 mg/mL, while the inhibition zone against *E. coli* at the same concentration was 11 mm (8).

Abass et al. investigated the antimicrobial effect of Red Roselle (*H. sabdariffa*) against various types of oral bacteria. The study revealed that at a concentration of 100 mg, the diameter of the inhibition zone against

Streptococcus mutans was 27.6 mm, against *S. aureus* was 22.3 mm, and against *E. faecalis* was 22.4 mm (9).

In another study, the effect of the aqueous extract of *H. sabdariffa* on *E. coli* was evaluated. Using a standard microdilution assay, the extract showed a minimum inhibitory concentration (MIC) of 6.5 mg/mL against the growth of enteropathogenic *E. coli* (EPEC). A time-kill kinetics assay demonstrated significant bactericidal activity at 24 hours with a concentration of 25 mg/mL. Furthermore, the extract effectively prevented basal induction (10).

Another study on the antibacterial properties of the methanolic extract of *H. sabdariffa* calyces revealed the presence of effective antibacterial agents. These agents exhibited significant inhibition zones against both gram-negative and gram-positive bacteria tested, surpassing the effects of penicillin, which showed weak or no activity, and competing with gentamicin (11).

A comparative study examined the antimicrobial effects of an aqueous extract of red rose calyx (RE), chlorhexidine (CH), amoxicillin-clavulanic acid (ACA), tetracycline (Tet), and metronidazole (Met) on *S. mutans*,

S. aureus, and *E. faecalis*. The results showed that at a dilution of 25 mg/mL, the inhibition zone diameters for *S. mutans*, *S. aureus*, and *E. faecalis* were 9.1 mm, 7.5 mm, and 8 mm, respectively (9).

In another study, the antimicrobial activity of sour tea extract was tested against *S. mitis* and *S. oralis*. The extract inhibited 29.1% of *S. mitis* and 63.23% of *S. oralis* bacteria (12).

Finally, the study by Marquez-Rodriguez et al. investigated the in vitro antibacterial activity of sour tea phenolic extract and its potential application in beef preservation. The results showed that the MICs against *E. coli*, *Salmonella enterica*, *S. aureus*, *Listeria monocytogenes*, and *Bacillus cereus* were 300, 300, 200, 200, and 200 mg/mL, respectively (13).

4.1. Conclusions

The results demonstrated that the sour tea plant possesses significant antimicrobial and color-producing properties.

Acknowledgements

The authors of this article express their gratitude to all the professors who contributed to the collection of data and the writing of this article.

Footnotes

Authors' Contribution: Study concept and design: M. T. and M. K.; Analysis and interpretation of data: M. R. All authors have read and approved the final manuscript.

Conflict of Interests Statement: The authors declared no conflicts of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Funding/Support: This manuscript received no financial support.

References

- Morales Cabrera M, Hernandez Morales J, Leyva Ruelas G, Salinas Moreno Y, Soto Rojas L, Castro Rosas J. Influence of variety and extraction solvent on antibacterial activity of roselle (*Hibiscus sabdariffa* L.) calyces. *J Med Plant Res*. 2013;7(31):2319-22. <https://doi.org/10.5897/jmpr12.1242>.
- Voon HC, Bhat R, Rusul G. Flower Extracts and Their Essential Oils as Potential Antimicrobial Agents for Food Uses and Pharmaceutical Applications. *Compr Rev Food Sci Food Saf*. 2011;11(1):34-55. <https://doi.org/10.1111/j.1541-4337.2011.00169.x>.
- Da-Costa-Rocha I, Bonnlaender B, Sievers H, Pischel I, Heinrich M. *Hibiscus sabdariffa* L. - a phytochemical and pharmacological review. *Food Chem*. 2014;165:424-43. [PubMed ID: 25038696]. <https://doi.org/10.1016/j.foodchem.2014.05.002>.
- Obouayeba AP, Nazaire Djyh B, Diabate S, Djaman J, Jean David N, Koné M, et al. Phytochemical and antioxidant activity of Roselle (*Hibiscus Sabdariffa* L.) petal extracts. *Res J Pharm Biol Chem Sci*. 2014;5(2):1453-65.
- Wang CJ, Wang JM, Lin WL, Chu CY, Chou FP, Tseng TH. Protective effect of Hibiscus anthocyanins against tert-butyl hydroperoxide-induced hepatic toxicity in rats. *Food Chem Toxicol*. 2000;38(5):411-6. [PubMed ID: 10762726]. [https://doi.org/10.1016/S0278-6915\(00\)00011-9](https://doi.org/10.1016/S0278-6915(00)00011-9).
- Fullerton M, Khatiwada J, Johnson JU, Davis S, Williams LL. Determination of antimicrobial activity of sorrel (*Hibiscus sabdariffa*) on *Escherichia coli* O157:H7 isolated from food, veterinary, and clinical samples. *J Med Food*. 2011;14(9):950-6. [PubMed ID: 21548802]. [PubMed Central ID: PMC3157304]. <https://doi.org/10.1089/jmf.2010.0200>.
- Chongwilaikasem N, Sithisarn P, Rojsanga P, Sithisarn P. Green extraction and partial purification of roselle (*Hibiscus sabdariffa* L.) extracts with high amounts of phytochemicals and in vitro antioxidant and antibacterial effects. *J Food Sci*. 2024;89(12):8819-35. [PubMed ID: 39437315]. [PubMed Central ID: PMC11673433]. <https://doi.org/10.1111/1750-3841.17418>.
- Venkatesan K, Venkatesan S, Manivannan N. Antibacterial Activity of *Hibiscus sabdariffa* (Rosella) Using Methanolic Extract. *J Pharm Bioallied Sci*. 2024;16(Suppl 2):S1191-4. [PubMed ID: 38882855]. [PubMed Central ID: PMC11174337]. https://doi.org/10.4103/jpbs.jpbs_527_23.
- Abass AA, Al-Magsoosi MJN, Kadhim WA, Mustafa R, Ibrahim SA, Aljdaimi AI, et al. Antimicrobial effect of Red Roselle (*Hibiscus Sabdariffa*) against different types of oral bacteria. *J Med Life*. 2022;15(1):89-97. [PubMed ID: 35186141]. [PubMed Central ID: PMC8852637]. <https://doi.org/10.25122/jml-2021-0184>.
- Mohamed-Salem R, Rodriguez Fernandez C, Nieto-Pelegrin E, Conde-Valentin B, Rumbero A, Martinez-Quiles N. Aqueous extract of *Hibiscus sabdariffa* inhibits pedestal induction by enteropathogenic *E. coli* and promotes bacterial filamentation in vitro. *PLoS One*. 2019;14(3). e0213580. [PubMed ID: 30849110]. [PubMed Central ID: PMC6407759]. <https://doi.org/10.1371/journal.pone.0213580>.
- Abdallah EM. Antibacterial efficiency of the Sudanese Roselle (*Hibiscus sabdariffa* L.), a famous beverage from Sudanese folk medicine. *J Intercult Ethnopharmacol*. 2016;5(2):186-90. [PubMed ID: 27104041]. [PubMed Central ID: PMC4835995]. <https://doi.org/10.5455/jice.20160320022623>.
- Salah R, Haleem AM, Mazin H. Potential effect of Roselle (*Hibiscus sabdariffa*) ethanol extract against *Streptococcus mitis* and *Streptococcus oralis*. *J Herb Med*. 2023;41. <https://doi.org/10.1016/j.hermed.2023.100710>.
- Marquez-Rodriguez AS, Nevarez-Baca S, Lerma-Hernandez JC, Hernandez-Ochoa LR, Nevarez-Moorillon GV, Gutierrez-Mendez N, et al. In Vitro Antibacterial Activity of *Hibiscus sabdariffa* L. Phenolic Extract and Its In Situ Application on Shelf-Life of Beef Meat. *Foods*. 2020;9(8). [PubMed ID: 32784385]. [PubMed Central ID: PMC7464790]. <https://doi.org/10.3390/foods9081080>.